



Polish botanical and mycological studies of the Antarctic terrestrial and fresh water ecosystems in 1977–2009: An overview

Piotr KÖHLER and Maria OLECH

*Zakład Badań i Dokumentacji Polarnej im. Prof. Z. Czeppego, Instytut Botaniki,
Uniwersytet Jagielloński, ul. Kopernika 27, 31-501 Kraków, Poland
<piotr.kohler@uj.edu.pl> <maria.olech@uj.edu.pl>*

Abstract: This paper recapitulates Polish botanical and mycological research on terrestrial and freshwater Antarctic ecosystems carried out between 1977 and 2009. The main results are briefly summarized. The references encompass nearly 200 papers on floristics, taxonomy, biogeography, ecology, cytology, biochemistry, physiology and genetics of lichens, mosses, fungi, algae and vascular plants inhabiting soils, rocks and inland waters in the Antarctic.

Key words: Antarctic, plants, fungi.

Introduction

The Polish Antarctic *H. Arctowski* Station was opened on February 26th, 1977, on the shores of Admiralty Bay, King George Island, the South Shetland Islands. By 2009 as many as 33 Polish expeditions had worked at the Station. The date of opening of the Polish Polar Station on King George Island and the conclusion of the 4th International Polar Year constitute the temporal frame of this paper.

The Polish research program was linked with numerous international programs coordinated by the Scientific Committee on Antarctic Research (SCAR), including BIOTAS, RiSCC, ClicOPEN and Aliens in Antarctica. Polish botanical and mycological studies covered areas such as taxonomy, biogeography and ecology, including the following topics: plant adaptation to polar conditions, plant succession on the forefields of retreating glaciers, and changes in ecosystems caused by human activity.

Nearly 200 papers on plants and fungi of the terrestrial and freshwater ecosystems of the Antarctic have been published by Polish scientists. The majority of

Pol. Polar Res. 32 (2): 157–174, 2011

these studies were devoted to lichenology and bryology, given the dominance of their representatives in the Antarctic. Among the most eminent achievements are several monographs and the description of many plant and fungal taxa new to science, as well as the discovery of numerous species hitherto not reported from the Antarctic. The present paper supplements the earlier overview by Rakusa-Suszczewski *et al.* (1998).

Floristics, taxonomy and biogeography

The purpose of the floristic research was to create an inventory of taxa present in particular sites and study areas, as well as their geographical distribution (Ochyra 1983).

The studies started with the discovery of the lichen *Hypogymnia lugubris* (Pers.) Krog in the Admiralty Bay area on King George Island (Lindsay and Ochyra 1982). The lichen biota in this area of King George Island was then thoroughly studied and a total of 110 mainly bipolar species were found and new localities were recorded (Olech 1989a). As many as 61 species found in the Admiralty Bay area had not been observed before in that area, 35 were new for the South Shetland Islands, 15 were hitherto not recorded in the Antarctic region and 6 were recorded for the first time in the Southern Hemisphere. Several taxa formerly identified as an Arctic-alpine element were also recorded; apparently, these were bipolar species. An interesting lichen species, *Japewia tornoenensis* (Nyl.) Tønsberg, new to the Antarctic and to the Southern Hemisphere, was found on King George Island and Livingston Island (Olech 1991a). The lichen biota of SSSI no. 8 in the Admiralty Bay area was studied (Myrcha *et al.* 1991).

In the South Shetlands area, namely King George I., Livingston I., Greenwich I., and Deception I., the occurrence of 24 *Lecanora* species was recorded, including the first discovery of an Antarctic endemic, *Lecanora sverdrupiana* Øvstedal in Maritime Antarctic (Śliwa and Olech 2002). At least 22 lichen species of the genus *Caloplaca* have been recorded in the South Shetland Islands. Of these, almost 30% have a bipolar or cosmopolitan range. On the other hand, it is thought that maritime species typical of islands have probably evolved *in situ* (Søchting and Olech 1995). The lichenobiotas in the area of Bunger Oasis (Olech 1989b) and Schirmacher Oasis in continental Antarctica have been studied.

Olech (1989c) prepared a vegetation map of Livingston Island with an inventory of the area before the founding of the Spanish polar station.

In the area of SSSI no. 8 of King George Island, 9 *Umbilicaria* species were recorded; one of them, *Umbilicaria umbilicarioides* (Stein) Krog *et* Swinscow, was found for the first time in the Antarctic (Krzewicka and Smykla 2004).

Lichenological study was also performed in the Lions Rump protected area (SSSI no. 34, currently known as ASPA no. 151); 104 lichen species have been re-

corded there. Several ecological groups of lichens were identified (Olech 1994) and distribution maps of various lichen species and of plant communities of the Lions Rump ASPA were produced.

Critical species lists of Antarctic lichens and lichenicolous fungi have been published by Olech (2001) and afterwards, as a result of long-term research, a monograph on the King George Island lichen biota was prepared, describing a total of 294 lichenized fungal species (approximately 77% of all hitherto known Antarctic lichens), including two species new for the Antarctic, seven species new for the South Shetland Islands and 68 species new for King George I. (Olech 2004).

Macrofungi (Macromycetes) collected on King George I. and Livingston I. were investigated in detail. Four species were recognized; one of them, *Arrhenia salina* (Høiland) Bon et Courtecuisse, had not been previously recorded in the Antarctic (Gumińska *et al.* 1994). Soil mycoflora, especially of the genus *Penicillium*, was studied in the vicinity of the *Arctowski* Station. Depending on the environment the percentage of the *Penicillium* fungi in the general population of fungal biota varied from 8.5 to 90% (Czarnecki and Białasiewicz 1987; Zabawski 1995; Białasiewicz and Czarnecki 1999; Alias *et al.* 2008).

Polish bryological studies in the Antarctic produced 15 papers on the floristics and phytogeography of the Antarctic bryophyte species (Ochyra and Bell 1984; Ochyra 1986; Ochyra and Newton 1986; Ochi and Ochyra 1986; Ochyra and Ochi 1986; Ochyra and Lewis-Smith 1996, 1999; Bednarek-Ochyra *et al.* 1999; Ochyra *et al.* 2000, 2008a; Ochyra and Zander 2002; Ochyra and Tyshchenko 2006; Seppelt and Ochyra 2008; Ochyra and Singh 2008; Li *et al.* 2009).

An outline of the phytogeography of Antarctic liverworts has been published by Ochyra and Váňa (1989b), as a supplement to their paper on the taxonomy of 11 liverworts from King George Island (Ochyra and Váňa 1989a). An annotated guide to Antarctic bryophytes exsiccata was also published (Ochyra *et al.* 1986), as well as a paper on the history of the Antarctic moss flora research (Ochyra *et al.* 1998).

Two liverwort species, *Hygrolembidium ventrosum* (Mitt.) Grolle and *Scapania obcordata* (Berggr.) S. Arnell, were recorded for the first time in the Antarctic and the latter had only been known from the Arctic. Papers solving numerous detailed taxonomic questions concerning particular species of mosses were also published by Ochyra (1987, 1990, 1993a, b, c, 1996b, c, 1997, 1998c, 1999a, b, c, 2004a, b, d), Ochyra and Lewis-Smith (1998) and Ochyra and Zander (2007). In addition, new species of mosses were described, namely *Ditrichum gemmiferum* (Ochyra and Lewis-Smith 1998), *D. lewis-smithii* (Ochyra 1996a), *Schistidium deceptionense* (Ochyra *et al.* 2003), *S. halinae* (Ochyra 1998a), *S. leptoneurum* (Ochyra 2004c), *S. lewis-smithii* (Ochyra 2003) and *S. steerei* (Ochyra 1987).

Soil and freshwater cyanobacteria and algae, especially diatoms, were studied by Zabawski and Piasecki (1981), Kawecka and Olech (1993, 2003, 2004), Luścińska and Kyć (1993), Kawecka *et al.* (1996, 1998), Starmach (1995), Mrozińska *et al.* (1998b, 2007), and Noga and Olech (2004). Diatoms dominated

the freshwater algal communities (some 90 taxa) and were mostly eurytopic and cosmopolitan species.

Polish bryological studies performed in the South Shetland Islands area were summarized in a monograph of King George Island mosses (Ochyra 1998b) where taxonomic, phytogeographic and ecological analyses of 61 moss species were presented. A monograph, *The liverwort flora of Antarctica*, has been published by Bednarek-Ochyra *et al.* in 2000; it was the first critical survey of this group of plants on this continent. In this book 27 Antarctic liverwort species are discussed; three taxa, namely *Pachyglossa spagazziniana* (C. Massal.) Herzog and Grolle var. *exilis* Herzog and Grolle, *P. fissa* (Mitt.) Herzog and Grolle, and *Scapania gamundiae* R. M. Schust. were recorded for the first time in this part of the world. This monograph presents detailed morphological and anatomical characteristics of all the species; taxonomic and nomenclatural issues are discussed and maps presenting their distribution in the Antarctic are included.

Another impressive monograph, *The illustrated moss flora of Antarctica*, has been published by Ochyra *et al.* (2008b). All species of Bryophyta hitherto recorded in the Antarctic are treated there. The descriptions of 111 species of mosses are accompanied by detailed drawings and distribution maps; their reproductive biology and ecology in the Antarctic is also thoroughly discussed.

Polish studies on the taxonomy and systematics of Antarctic lichenicolous fungi were initiated in late 1980s. In the material collected between 1986 and 1996 by several expeditions to *Arctowski* Station and in 1988/1989 to *Bunger Oasis* in continental Antarctica, 65 species of lichenicolous fungi were identified; out of them numerous taxa were recorded from the Antarctic for the first time (3 genera and 31 species). Among others there were two new species described: *Dactylospora dobrowolskii* Olech *et* Alstrup and *Octospora arctowskii* Olech *et* Mleczko (Ascomycota) named in honour of two famous Polish scientists, A.B. Dobrowolski and H. Arctowski, the participants of the *Belgica* expedition (1897-99) (Olech and Alstrup 1990, 1995, 1996; Olech and Mleczko 2000).

Taxonomic studies on lichenized fungi resulted in the description of numerous new species, namely *Caloplaca buelliae* Olech *et* Söchting, *C. iomma* Olech *et* Söchting, *C. psoromatis* Olech *et* Söchting and *C. siphonospora* Olech *et* Söchting, *C. scolecomarginata* Söchting *et* Olech, *C. frigida* Söchting, *Bryoria forsteri* Olech *et* Bystrek, *Bacidia subcoprodes* Olech *et* Czarnota and *B. chrysocolla* Olech, Czarnota *et* Llop (Olech and Söchting 1993; Söchting and Olech 2000; Olech and Bystrek 2004; Olech and Czarnota 2009). In several papers (*e.g.* Śliwa and Olech 2002) difficult taxonomic problems within the lichen genus *Lecanora* were discussed.

Taxonomic and chemotaxonomic revision of *Cladonia* species from King George Island was undertaken. Fourteen species were found there including *Cladonia asahinae* J.W. Thomson, recorded for the first time (Osyczka and Olech 2004). A key for the identification of species belonging to the genus *Cladonia* was

also provided (Osyczka and Olech 2005b). Chemotaxonomic revision of the genus *Tephromela* (six species) was also published (Osyczka and Olech 2005a).

Lichenological material collected from the South Shetland Islands and from Schirmacher Oasis and Bunger Oasis on continental Antarctica led to the publication of new data on four species within the lichen genus *Lepraria* (Osyczka *et al.* 2009).

Several papers published by Polish scientists have been devoted to the taxonomy of algae. One new genus, one new species and several new varieties were described (Massalski *et al.* 1995, 1999b; Mrozińska *et al.* 1998a; Olech *et al.* 1998b; Kostikov *et al.* 2003).

Morphology, anatomy, cytology, embryology, and biochemistry

Karyological and cytological studies on Antarctic liverworts were carried out by Ochyra *et al.* (1982) and on mosses by Kuta *et al.* (1982) and Przywara *et al.* (1984). Karyological analysis of 15 species was performed and, in several cases, the number of chromosomes was given for the first time. Rhizoid gemmae in a moss *Ditrichum brotherusii* were also described (Ochyra 1996d).

Morphological studies covered both flowering plants and algae. Anatomical, morphological and functional variability of *Deschampsia antarctica* from King George Island was studied. Significant differences between particular populations from various habitats were considered to be a result of the lack of interspecific competition in these habitats (Barcikowski *et al.* 2001, 2003; Chwedorzewska *et al.* 2008a). Anatomical studies on the ultrastructure of the leaves of *D. antarctica* and *Colobanthus quitensis* published by Giełwanowska (2005), Giełwanowska and Szczuka (2005), and Giełwanowska *et al.* (2005b, 2008a, b) helped to explain the adaptations of both species to extreme habitat conditions and their anatomical reaction to abiotic stress factors. New ultrastructural features were found in the mesophyll cells (Giełwanowska and Szczuka 2005). The morphology of pollen of *D. antarctica* and *C. quitensis* has been studied by Sadowska (1998), Giełwanowska *et al.* (2008c) and Szczuka *et al.* (2008).

Morphological diversity of some groups of algae and blue-green algae from King George Island was studied by Kawecka *et al.* (1996) (Bacillariophyta), Mrozińska *et al.* (1998b) (Chrysophyta), Massalski *et al.* (1994, 2001) (Chlorophyta), and Massalski *et al.* (1999a, b) (Cyanobacteria). The cell ultrastructure of *Xanthonema* (Xanthophyta) was analyzed in detail. Peculiar processes during mitosis and cytokinesis in this species were recorded by Massalski *et al.* (2009).

Biochemical studies on Antarctic mosses and the flowering plants, *D. antarctica* and *C. quitensis*, were studied by Czczuga *et al.* (1982, 1984) and Piotrowicz-Cieślak *et al.* (2005), who investigated the content of carotenoids and carbohydrates in those plants. The carotenoid content in the thalli of different Antarctic lichen species was examined by Xavier-Filho *et al.* (1986), Czczuga *et al.* (1986, 1996),

Czczuga and Xavier-Filho (1987), Czczuga and Olech (1989) and by Czczuga and Koch (1991). Lipids and insoluble carbohydrates in Antarctic lichens were also studied by Giełwanowska *et al.* (2008b) while the chemical composition of some dominating plants in the maritime Antarctic tundra was investigated by Fabiszewski and Wojtuń (2000).

Physiology and reproduction

Changes in the chlorophyll content in selected moss species were investigated by Barcikowski and Loro (1999).

Many papers by Polish physiologists concerned the grass *D. antarctica*. Its response to temperature stress was studied by Bystrzejewska (2001), Giełwanowska (2003) and Bystrzejewska-Piotrowska and Urban (2009). The response of this grass to the concentration of biogenic substances in its habitats was investigated by Nędzarek and Chwedorzewska (2004). It was demonstrated that *D. antarctica* is highly tolerant to the extreme diversity of nitrogen and phosphorus concentrations both in soil and water. Pollen growth and pollination types in *C. quitensis* were studied by Giełwanowska *et al.* (2006, 2007) and Szczuka *et al.* (2008), while the biology and generative reproduction of *D. antarctica* was studied by Giełwanowska *et al.* (2005a).

The production of enzymes in hyphal fungi was investigated by Kasieczka-Burnecka *et al.* (2005); pectinolytic enzymes and tannases were found in these organisms. Ecophysiological studies of lichens were conducted by Schroeter *et al.* (1995). They examined the influence of microclimatic factors on primary production in *Usnea antarctica*. A mechanism of two-stage hydration/dehydration of the lichen thallus, extremely important during weather changes accompanied by frost, was discovered. The two-stage hydration mechanism is present only in fruticose lichens and was not observed in those crustose lichens forming a flat crust on the substratum (Harańczyk *et al.* 2009). The mechanism of resistance of lichens to low temperatures was also studied in detail. It appeared that the tightly bound water in lichen thalli usually does not freeze. It was found also that, when the temperature drops down, free water which would otherwise freeze in the thallus of a lichen turns into non-freezing bound water. This is possible due to the formation of a gel-like structure which traps the water (Harańczyk *et al.* 1998a, b, c, 2000a, b, 2001, 2003a, b, c, 2006, 2008, 2009).

Ecology

The first Polish studies on plant communities in the area near to the *Arctowski* Station were performed in 1979/1980 (Furmańczyk and Ochyra 1982).

Terrestrial biotopes of the coastal Admiralty Bay ecosystem were also described in detail by Zarzycki (1993). The influence of wind on the spreading of diaspores in this region was also investigated (Hołdyński *et al.* 2003). The problem of the introduction of an alien grass species, *Poa annua* L. was studied by Olech (1996b) and Chwedorzewska (2008). High variability and high genetic diversity suggest that this grass originated from many different places, whereas its survival and growth in the Antarctic are possible due to the favourable conditions created by global climate warming.

Plant communities of King George Island, of South Shetland Islands and the Antarctic as a whole were described and discussed in the papers by Ochyra (1984) and Olech (1993, 1998a, 2002).

The growth rate and biomass production of *Deschampsia antarctica* in the Admiralty Bay area was studied by Barcikowski *et al.* (1999). The biomass of some species of mosses occurring on King George Island was studied by Barcikowski and Gurtowska (1999).

Diatom communities inhabiting streams and stagnant water bodies were also thoroughly investigated. Long term observations of the structure of these communities showed that they are changing, reflecting the unstable environmental conditions which are the result of the temperature rise and desiccation caused by climatic change (Kawecka and Olech 1993, 2004; Kawecka *et al.* 1998; Ochwanowski and Pocięcha 2005).

The ecology of algae, especially their colonization of glacial moraines as well as the influence of penguin colonies upon algae distribution, was analyzed by Mrozińska *et al.* (1998a, 2007).

Plant communities in abandoned penguin rookeries of King George Island and, in general, the influence of penguin guano on particular plant species, especially *Deschampsia antarctica*, were thoroughly studied by Tatur and Myrcha (1989), Olech (1990, 1996a, b, 1998a), Tatur *et al.* (1997), Pisarek *et al.* (2003), Chwedorzewska *et al.* (2004), Smykla (2005) and Smykla *et al.* (2006, 2007). The role of ornithogenic soils in the functioning of polar ecosystems was also discussed in the papers by Krywult *et al.* (2003), Barcikowski *et al.* (2005) and Smykla (2008). The colonization of deglaciated areas by plant communities was studied by Olech (1996b) and Olech and Massalski (2001). Plant biomass and seasonal changes in the concentration of some organic compounds, principally the chlorophyll, in mosses and vascular plants, were analyzed by Barcikowski and Luścińska (2001).

Human influence on the Antarctic environment was also studied, especially the trace element content, including heavy metals (particularly lead), in the thalli of lichens growing in the vicinity of polar stations (Olech 1991b, 1996a, 1997; Olech *et al.* 1993, 1998a, 2000; Osyczka *et al.* 2003, 2007; Smykla *et al.* 2005). The pollution of the Antarctic environment was also studied by means of the analysis of the amount of radionuclides in lichens and mosses (Schuch *et al.* 1993; Godoy *et al.* 1998; Mietelski *et al.* 2000, 2008; Gaca *et al.* 2003). In comparison

with other organisms occurring on King George Island the concentration of radionuclides was highest in lichens (Mietelski *et al.* 2008).

The initial stages of flora and plant-cover synanthropization were monitored by Olech (1994, 1998b). Special attention was paid to the invasive plants (Olech 1994, 1998b; Chwedorzewska *et al.* 2008b; Chwedorzewska 2009).

The study of samples from the soil from Coulman Island produced interesting information on propagules of the potential colonists of the Antarctic region (Lewis-Smith and Ochyra 2006).

Peat-forming vegetation of King George Island of extreme floristic poverty was studied by Fabiszewski and Wojtuń (1993, 1997). Using C-14 dating these authors demonstrated that these peat-banks developed some 4000 years ago.

Palynological (aerobiological) study was performed along a transect from Antarctica to Poland (4th March – 13th April 1990), from which a valuable collection of lichen propagules was sampled (Harmata and Olech 1991).

Genetics

Genetic research on *Deschampsia antarctica* was recently undertaken. Intra-specific genetic variability and affinity between the populations from various islands of the South Shetlands archipelago, from islands neighbouring the Antarctic Peninsula and from the Falkland Islands was studied. A low genetic variability was found (Chwedorzewska and Nędzarek 2005; Kyryachenko *et al.* 2005; Chwedorzewska 2006; Chwedorzewska and Bednarek 2008).

Genetic studies of algae were also carried out by Rybalka *et al.* (2009). The endemism and genotypic diversity of the Antarctic algae of the family *Tribo-nemataceae* (*Xanthophyceae*) were analyzed. It was found that Antarctic populations differ from those inhabiting moderate climatic zones. Currently-identified morphospecies do not reflect the actual biodiversity within this group.

Acknowledgements. — We would like to thank three anonymous reviewers for their very careful and inspiring reviews and to Dr Roger Bamber (ARTOO), who kindly corrected English through the manuscript.

References

- ALIAS S.A., OMAR S. and SMYKLA J. 2008. Preliminary data on soil microfungi from Beaufort Island, Ross Sea, Antarctica. *In*: A. Anton, A.A. Samah and C.M. Wong (eds) *From the tropics to the Poles*. Academy of Sciences of Malaysia, Kota Kinabalu, Sabah: 65–69.
- BARCİKOWSKI A. and GURTOWSKA J. 1999. Studies on the biomass of selected species of Antarctic mosses on King George Island, South Shetland Islands, Antarctica. *Polish Polar Research* 20: 283–290.

- BARCIKOWSKI A. and LORO P.M. 1999. Changes in chlorophyll content throughout the year in selected species of mosses on King George Island, South Shetland Islands, maritime Antarctic. *Polish Polar Research* 20: 291–299.
- BARCIKOWSKI A. and LUŚCIŃSKA M. 2001. Plant biomass structure of Antarctic terrestrial communities in relation to energy flow. *Folia Facultatis Scientiarum Naturalium Universitatis Masarykianae Brunensis, Geographia* 25: 65–72.
- BARCIKOWSKI A., CZAPLEWSKA J., GIELWANOWSKA I., LORO P., SMYKLA J. and ZARZYCKI K. 2001. *Deschampsia antarctica* (Poaceae) – the only native grass from Antarctica. In: L. Frey (ed.) *Studies on grasses in Poland*. Kraków: 367–377.
- BARCIKOWSKI A., CZAPLEWSKA J., LORO P., ŁYSZKIEWICZ A., SMYKLA J. and WOJCIECHOWSKA A. 2003. Ecological variability of *Deschampsia antarctica* in the area of Admiralty Bay (King George Island, Maritime Antarctic). In: L. Frey (ed.) *Problems of grass biology*. Kraków: 383–396.
- BARCIKOWSKI A., ŁYSZKIEWICZ A., LORO P., REKTORIS L., SMYKLA J., WINCENCIAK A. and ZUBEL P. 2005. Keystone species and ecosystems functioning: the role of penguin colonies in differentiation of the terrestrial vegetation in the Maritime Antarctic. *Ecological Questions* 6: 117–128.
- BARCIKOWSKI A., ŁYŻWIŃSKA R. and ZARZYCKI K. 1999. Growth rate and biomass production of *Deschampsia antarctica* Desv. in the Admiralty Bay region, South Shetland Islands, Antarctica. *Polish Polar Research* 20: 301–311.
- BEDNAREK-OCHYRA H., LEWIS-SMITH R.I., and OCHYRA R. 1999. The genus *Plagiothecium* (Plagiotheciaceae, Bryopsida) in Antarctica. *Bryobrothera* 5: 55–64.
- BEDNAREK-OCHYRA H., VÁŇA J., OCHYRA R. and LEWIS-SMITH R.I. 2000. *The liverwort flora of Antarctica*. Polish Academy of Sciences, Institute of Botany, Cracow: xvi + 236 pp.
- BIAŁASIEWICZ D. and CZARNECKI B. 1999. Microfungi in the aerosphere of the *Arctowski* Polar Station. *Polish Polar Research* 20: 319–324.
- BYSTRZEJSKA G. 2001. Photosynthetic temperature response of Antarctic plant *Deschampsia antarctica* and of temperate region plant *Deschampsia caespitosa*. *Polish Journal of Ecology* 49: 215–219.
- BYSTRZEJSKA-PIOTROWSKA G. and URBAN P.L. 2009. Tufted hairgrass (*Deschampsia caespitosa*) exhibits a lower photosynthetic plasticity than Antarctic hairgrass (*D. antarctica*). *Journal of Integrative Plant Biology* 51: 593–603.
- CHWEDORZEWSKA K.J. 2006. Preliminary genetic study on species from genus *Deschampsia* from Antarctic (King George I.) and Arctic (Spitsbergen). *Polar Biosciences* 19: 142–147.
- CHWEDORZEWSKA K.J. 2008. *Poa annua* L. in Antarctic: searching for the source of introduction. *Polar Biology* 31: 263–268.
- CHWEDORZEWSKA K.J. 2009. Terrestrial Antarctic ecosystems in the changing world: an overview. *Polish Polar Research* 30: 263–276.
- CHWEDORZEWSKA K.J. and BEDNAREK P.T. 2008. Genetic variability in the Antarctic hairgrass *Deschampsia antarctica* Desv. from Maritime Antarctic and Subantarctic sites. *Polish Journal of Ecology* 56: 209–216.
- CHWEDORZEWSKA K.J. and NĘDZAREK A. 2005. Preliminary genetic study on *Deschampsia antarctica*, and their related species from genus *Deschampsia*. In: W. Prus-Głowacki and E.M. Pawlaczyk (eds) *Variability and evolution – new perspectives*. Uniwersytet im. A. Mickiewicza w Poznaniu, Seria Biologiczna 72, Poznań: 149–157.
- CHWEDORZEWSKA K.J., BEDNAREK P.T. and PUCHALSKI J. 2004. Molecular variation of Antarctic grass *Deschampsia antarctica* Desv. from King George Island (Antarctica). *Acta Societatis Botanicorum Poloniae* 73: 23–29.
- CHWEDORZEWSKA K.J., GIELWANOWSKA I., SZCZUKA E. and BOCHENEK A. 2008a. High anatomical and low genetic diversity in *Deschampsia antarctica* Desv. from King George Island (the Antarctic). *Polish Polar Research* 29: 377–386.

- CHWEDORZEWSKA K.J., OLECH M. and KORCZAK M. 2008b. *Obcy w Antarktyce*. In: A. Kowalska, A. Lachota, H. Marszałek and J. Pereyma (eds) *Środowisko przyrodnicze obszarów polarnych*. Wrocław: 197–201.
- CZARNECKI B. and BIAŁASIEWICZ D. 1987. Fungi as a component of the aerosphere in the H. Arctowski Polar Station and its vicinity (King George Island, South Shetland Islands). *Polish Polar Research* 8: 153–158.
- CZECZUGA B. and KOCH P. 1991. Investigaciones en carotenoides de líquenes. XXXIV. Diferencias en el contenido de carotenoides en talos de *Xanthoria candelaria* Arnold, de la Antártica y de Concepción, Chile. *Instituto Antártico Chileno, Serie Científica* 41: 107–111.
- CZECZUGA B. and OLECH M. 1989. Investigations on carotenoids in lichens. XXIV. Further studies of carotenoids in lichens of the Antarctica. *Serie Científica Instituto Nacional Antártico Chileno* 39: 91–96.
- CZECZUGA B. and XAVIER-FILHO L.X. 1987. Investigations on carotenoids in lichens. X. Luteoxantin and apo-12'-violaxanthal in lichens from Antarctica. *Instituto Antártico Chileno, Serie Científica* 36: 151–155.
- CZECZUGA B., GUTKOWSKI R. and CZERPAK R. 1982. Investigations of carotenoids in Embryophyta. II. Musci from the Antarctic. *Nova Hedwigia* 36: 695–701.
- CZECZUGA B., GUTKOWSKI R. and CZERPAK R. 1984. Studies on the carotenoids of the Embryophyta. VII. Representatives of the families Gramineae and Caryophyllaceae from King George Island (South Shetland Islands). *Polish Polar Research* 5: 113–120.
- CZECZUGA B., GUTKOWSKI R. and CZERPAK R. 1986. Investigations on carotenoids in lichens. V. Lichens from Antarctica. *Polish Polar Research* 7: 295–303.
- CZECZUGA B., INOUE M. and UPRETI D.K. 1996. Carotenoids in lichens from the Antarctic. *Nankyoku Shiryo (Antarctic Record)* 40: 247–257.
- FABISZEWSKI J. and WOJTUŃ B. 1993. Peat-forming vegetation. In: S. Rakusa-Suszczewski (ed.) *The Maritime Antarctic coastal ecosystem of Admiralty Bay*. Department of Antarctic Biology, Polish Academy of Sciences, Warsaw: 189–195.
- FABISZEWSKI J. and WOJTUŃ B. 1997. The occurrence and development of peat mounds on King George Island (Maritime Antarctic). *Acta Societatis Botanicorum Poloniae* 66: 223–229.
- FABISZEWSKI J. and WOJTUŃ B. 2000. Chemical composition of some dominating plants in the maritime Antarctic tundra (King George Island). *Bibliotheca Lichenologica* 75: 79–91.
- FURMAŃCZYK K. and OCHYRA R. 1982. Plant communities of the Admiralty Bay region (King George Island, South Shetland Islands, Antarctic) I. Jasnorzewski Gardens. *Polish Polar Research* 3: 25–39.
- GACA P., MIETELSKI J. W. and OLECH M.A. 2003. Radioactive contamination of lichens and mosses collected in Antarctica. In: M.A. Olech (ed.) *XXIX International Polar Symposium*. Kraków: 39–45.
- GIELWANOWSKA I. 2003. *Deschampsia antarctica* responses to abiotic stress factors. *Acta Physiologiae Plantarum* 25: 61–62.
- GIELWANOWSKA I. 2005. *Specific developmental characters of Antarctic vascular plants Colobanthus quitensis (Kunth) Bartl. and Deschampsia antarctica Desv.* Wydawnictwo Uniwersytetu Warmińsko-Mazurskiego, Olsztyn: 175 pp.
- GIELWANOWSKA I. and SZCZUKA E. 2005. New ultrastructural features of organelles in leaf cells of *Deschampsia antarctica* Desv. *Polar Biology* 28: 951–955 [in Polish].
- GIELWANOWSKA I., BOCHENEK A. and LORO P.M. 2005a. Biology of generative reproduction of *Deschampsia antarctica*. In: L. Frey (ed.) *Biology of grasses*. Polish Academy of Sciences, Kraków: 181–195.
- GIELWANOWSKA I., BOCHENEK A. and LORO P.M. 2008a. Anatomical responses of *Colobanthus quitensis* (Kunth) Bartl. and *Deschampsia antarctica* Desv. to abiotic stress factors. *Ecological Questions* 9: 45–56.

- GIEŁWANOWSKA I., BOCHENEK A. and SZCZUKA E. 2007. Development of the pollen in the Antarctic flowering plant *Colobanthus quitensis* (Kunth) Bartl. *Acta Agrobotanica* 60: 3–8.
- GIEŁWANOWSKA I., GOJŁO E., GÓRECKI R. and OLECH A.M. 2008b. Lipids and insoluble carbohydrates in thallus cells of the Antarctica lichens. *Acta Biochimica Polonica* 55, Supplement 3: 224.
- GIEŁWANOWSKA I., SZCZUKA E. and BOCHENEK A. 2006. Pollination in the Antarctic flowering plant *Colobanthus quitensis* (Kunth) Bartl. *Acta Agrobotanica* 59: 123–131.
- GIEŁWANOWSKA I., SZCZUKA E., BEDNARA J. and GÓRECKI R. 2005b. Anatomical features and ultrastructure of *Deschampsia antarctica* (Poaceae) leaves from different growing habitats. *Annals of Botany* 96: 1109–1119.
- GIEŁWANOWSKA I., SZCZUKA E., PIDEK A., SETA A. and DOMACIUK M. 2008c. The unique features and ultrastructure of the microspore and pollen grain of the Antarctic dicotyledonous plant *Colobanthus quitensis* (Kunth) Bartl. *Sovremennyj nauchnyj vestnik* 27: 46–54.
- GODOY J.M., SCHUCH L.A., NORDEMANN D.J.R., REIS V.R.G., RAMALHO M., RECIO J.C., BRITO R.R.A. and OLECH M.A. 1998. ^{137}Cs , $^{226,228}\text{Ra}$, ^{210}Pb and ^{40}K concentrations in Antarctic soil, sediment and selected moss and lichen samples. *Journal of Environmental Radioactivity* 41: 33–45.
- GUMIŃSKA B., HEINRICH Z. and OLECH M. 1994. Macromycetes of the South Shetland Islands (Antarctica). *Polish Polar Research* 15: 103–109.
- HARAŃCZYK H., BACIOR M. and OLECH M.A. 2008. Deep dehydration of *Umbilicaria aprina* thalli observed by proton NMR and sorption isotherm. *Antarctic Science* 20: 527–535.
- HARAŃCZYK H., BACIOR M. and OLECH M. 2009. Deep dehydration of Antarctic lichen *Leptogium puberulum* observed by NMR and Sorption isotherm. *Acta Physica Polonica A* 115: 516–520.
- HARAŃCZYK H., GAŹDZIŃSKI S. and OLECH M. 1998a. Freezing protection mechanism in *Cladonia mitis* as observed by proton magnetic relaxation. *Lichenologist* 30: 417–431.
- HARAŃCZYK H., GAŹDZIŃSKI S. and OLECH M. 1998b. Initial stages of lichen hydration observed by proton magnetic relaxation. *New Phytologist* 138: 192–202.
- HARAŃCZYK H., GAŹDZIŃSKI S. and OLECH M. 2000a. Freezing protection mechanism in *Cladonia mitis* as observed by proton magnetic relaxation. *Bibliotheca Lichenologica* 75: 265–274.
- HARAŃCZYK H., GAŹDZIŃSKI S. and OLECH M. 2000b. Low temperature effect on the thallus of *Cladonia mitis* as observed by proton spin-lattice relaxation. *Molecular Physics Reports* 29: 135–138.
- HARAŃCZYK H., GRANDJEAN J. and OLECH M. 1998c. The freezing of water bound in thallus of *Cladonia mitis* as observed by ^1H NMR+. In: *Materiały XXX ogólnopolskiego seminarium na temat magnetycznego rezonansu jądrowego i jego zastosowań*. Kraków: 276–279.
- HARAŃCZYK H., GRANDJEAN J. and OLECH M. 2001. Low temperature effect in D_2O -hydrated Antarctic lichen *Himantormia lugubris* as observed by H-NMR. *Molecular Physics Reports* 33: 220–224.
- HARAŃCZYK H., GRANDJEAN J. and OLECH M. 2003a. Freezing of water bound in lichen thallus as observed by ^1H NMR. I. Freezing of loosely bound water in *Cladonia mitis* at different hydration levels. *Colloids and Surfaces, B: Biointerfaces* 28: 239–249.
- HARAŃCZYK H., GRANDJEAN J., OLECH M. and MICHALIK M. 2003b. Freezing of water bound in lichen thallus as observed by H NMR. II. Freezing protection mechanisms in a cosmopolitan lichen *Cladonia mitis* and in Antarctic lichen species at different hydration levels. *Colloids and Surfaces, B: Biointerfaces* 28: 251–260.
- HARAŃCZYK H., LIGĘZOWSKA A. and OLECH M.A. 2003c. Desiccation resistance of the lichen *Turgidosculum complicatulum* and its photobiont *Prasiola crispa* by proton magnetic relaxation, sorption kinetics and sorption isotherm. In: M.A. Olech (ed.) *XXIX International Polar Symposium*. Kraków: 51–56.
- HARAŃCZYK H., PIETRZYK A., LEJA A. and OLECH M.A. 2006. Bound water structure on the surfaces of *Usnea antarctica* as observed by NMR and sorption isotherm. *Acta Physica Polonica, Ser. A* 109 (3): 411–416.

- HARMATA K. and OLECH M. 1991. Transect for aerobiological studies from Antarctica to Poland. *Grana* 30: 458–463.
- HOLDYŃSKI C., LORO P.M. and PISAREK W. 2003. Wind dispersal of *Deschampsia antarctica* diaspores at the vicinity of the Arctowski Polar Station. In: M.A. Olech (ed.) *XXIX International Polar Symposium*. Kraków: 57–60.
- KASIECZKA-BURNECKA M., KALINOWSKA H., KUC K. and TURKIEWICZ M. 2005. Produkcja enzymów pektynolitycznych i tannaz przez antarktyczne grzyby strzępkowe. In: M. Józwiak and R. Kozłowski (eds) *Polish Polar Studies. XXXI Polar Symposium*. Kielce: 60–68.
- KAWECKA B. and OLECH M. 1993. Diatom communities in the Vanishing and Ornithologist Creek, King George Island, South Shetlands, Antarctica. *Hydrobiologia* 269/270: 327–333.
- KAWECKA B. and OLECH M.A. 2003. Long-term observations of diatom communities structure of Vanishing and Ornithologists creeks (King George Island, South Shetlands Islands, Maritime Antarctica). In: M.A. Olech (ed.) *XXIX International Polar Symposium*. Kraków: 67–69.
- KAWECKA B. and OLECH M. 2004. Diatom diversity of streams in Finnish Lapland and maritime Antarctica. In: M. Poulin (ed.) *Seventeenth International Diatom Symposium 2002, Ottawa, Canada*. Biopress Limited, Bristol: 161–186.
- KAWECKA B., OLECH M. and NOWOGRODZKA-ZAGÓRSKA M. 1996. Morphological variability of the diatom *Luticola muticopsis* (van Heurck) D.G. Mann in the inland waters of King George Island, South Shetland Islands, Antarctica. *Polish Polar Research* 17: 143–150.
- KAWECKA B., OLECH M., NOWOGRODZKA-ZAGÓRSKA M. and WOJTUŃ B. 1998. Diatom communities in small water bodies at H. Arctowski Polish Antarctic Station, King George Island, South Shetland Islands, Antarctica. *Polar Biology* 19: 183–192.
- KOSTIKOV I., MASSALSKI A. and OLECH M.A. 2003. Taxonomical and ecological studies on the pioneer soil algae from deglaciated areas of Maritime Antarctica. In: M.A. Olech (ed.) *XXIX International Polar Symposium*. Kraków: 71–73.
- KRYWULT M., SMYKLA J. and WINCENCIAK A. 2003. Influence of ornithogenic fertilization on nitrogen metabolism of the Antarctic vegetation. In: M.A. Olech (ed.) *XXIX International Polar Symposium*. Kraków: 79–84.
- KRZEWICKA B. and SMYKLA J. 2004. The lichen genus *Umbilicaria* from the neighbourhood of Admiralty Bay (King George Island, maritime Antarctic), with a proposed new key to all Antarctic taxa. *Polar Biology* 28: 15–25.
- KUTA E., OCHYRA R. and PRZYWARA L. 1982. Karyological studies on Antarctic mosses. I. *The Bryologist* 85: 131–138.
- KYRYACHENKO S.S., KOZERETSKA I.A., RAKUSA-SUSZCZEWSKI S. 2005. The genetic and molecular biological enigma of *Deschampsia antarctica* in Antarctica. *Cytology and Genetics* 39: 75–80.
- LEWIS-SMITH R.I. and OCHYRA R. 2006. High altitude Antarctic soil propagule bank yields an exotic moss and potential colonist. *Journal of the Hattori Botanical Laboratory* 100: 325–331.
- LI S.-P., OCHYRA R., WU P.-C., SEPPELT R.D., CAI M.-H., WANG H.-Y. and LI C.-S. 2009. *Drepanocladus longifolius* (Amblystegiaceae), and addition to the moss flora of King George Island, South Shetland Islands, with a review of Antarctic benthic mosses. *Polar Biology* 32: 1415–1425.
- LINDSAY D.C. and OCHYRA R. 1982. The lichen genus *Hypogymnia* (Nyl.) Nyl. in the Antarctic. *Fragmenta Floristica et Geobotanica* 28: 413–416.
- LUŚCIŃSKA M. and KYĆ A. 1993. Algae inhabiting creeks of the region of “H. Arctowski” Polish Antarctic Station, King George Island, South Shetlands. *Polish Polar Research* 14: 393–405.
- MASSALSKI A., KOSTIKOV I., OLECH M. and HOFFMANN L. 2009. Mitosis, cytokinesis and multinuclearity in a *Xanthonema* (Xanthophyta) isolated from Antarctic. *European Journal of Phycology* 44: 263–275.

- MASSALSKI A., MROZIŃSKA T. and OLECH M. 1994. Ultrastructure of *Lobosphaera reniformis* (Watanabe) Komárek et Fott (= Chlorellales) from King George Island, South Shetland Islands, Antarctica. *Acta Societatis Botanicorum Poloniae* 63: 205–210.
- MASSALSKI A., MROZIŃSKA T. and OLECH M. 1995. *Lobococcus irregularis* (Boye-Pet.) Reisigl var. *antarcticus* var. nov. (Chlorellales, Chlorophyta) from King George Island, South Shetland Islands, Antarctica, and its ultrastructure. *Nova Hedwigia* 61: 199–206.
- MASSALSKI A., MROZIŃSKA T. and OLECH M. 1999a. Ultrastructure of selected Cyanophyta/Cyanobacteria from King George Island, Antarctica. *Algological Studies* 94: 249–259.
- MASSALSKI A., MROZIŃSKA T. and OLECH M. 1999b. Ultrastructures of Antarctic algae. *Pseudosiderocelopsis antarctica* gen. et sp. nov. (Chlorophyta). *Algological Studies* 92: 1–10.
- MASSALSKI A., MROZIŃSKA T. and OLECH M. 2001. Ultrastructural observations on five pioneer soil algae from ice denuded areas (King George Island, West Antarctica). *Polar Bioscience* 14: 61–70.
- MIETELSKI J.W., GACA P. and OLECH M.A. 2000. Radioactive contamination of lichens and mosses collected in South Shetlands and Antarctic Peninsula. *Journal of Radioanalytical and Nuclear Chemistry* 245: 527–537.
- MIETELSKI J.W., OLECH M.A., SOBIECH-MATURA K., HOWARD B.J., GACA P., Zwolak M., Błażej S. and Tomankiewicz E. 2008. ^{137}Cs , ^{40}K , ^{238}Pu , $^{239+240}\text{Pu}$ and ^{90}Sr in biological samples from King George Island (Southern Shetlands) in Antarctica. *Polar Biology* 31: 1081–1089.
- MROZIŃSKA T., CZERWIK-MARCINKOWSKA J. and SMYKLA J. 2007. Desmids and associating algae of terrestrial small water bodies in the Admiralty Bay area (King George Island, Maritime Antarctic). *Oceanological and Hydrobiological Studies* 36 Supplement 1: 1–10.
- MROZIŃSKA T., OLECH M. and MASSALSKI A. 1998a. Algae of ponds and a stream on moraines of Ecology Glacier (King George Island, South Shetland Islands, Antarctica). *Nova Hedwigia* 67: 169–188.
- MROZIŃSKA T., OLECH M. and MASSALSKI A. 1998b. Cysts of Chrysophyceae from King George Island (South Shetland Islands, Antarctica). *Polish Polar Research* 19: 205–210.
- MYRCHA A., OCHYRA R. and TATUR A. 1991. Site of Special Scientific Interest No. 8 – western shores of Admiralty Bay, King George Island, South Shetland Islands. In: R.Z. Klekowski and K.W. Opalinski (eds) *The first Polish-Soviet Antarctic Symposium*. Institute of Ecology Publishing Office, Dziekanów Leśny: 157–168.
- NĘDZAREK A. and CHWEDORZEWSKA K.J. 2004. Nutrients content in water supplying chosen sites of Antarctic grass *Deschampsia antarctica* Desv. (King George Island, Antarctica). In: A. Styszyńska and A.A. Marsz (eds) *Polish Polar Studies*. Gdynia: 263–270.
- NOGA T. and OLECH M.A. 2004. Diatom communities in Moss Creek (King George Island, South Shetland Islands, Antarctica) in two summer seasons: 1995/96 and 2001/02. *Oceanological and Hydrobiological Studies* 33: 103–120.
- OCHI H. and OCHYRA R. 1986 (1985). *Bryum dichotomum* Hedw., a species new to the Antarctic. *Lindbergia* 11: 157–160.
- OCHWANOWSKI P. and POCHYCHA A. 2005. The impact of abiotic factors on diatom density dynamics in the freshwater Uncle Lake near the Henryk Arctowski Polish Antarctic Station during the austral summer. *Oceanological and Hydrobiological Studies* 34 Supplement 3: 257–267.
- OCHYRA R. 1983. Świat roślinny Antarktyki. *Kosmos* 32: 277–292.
- OCHYRA R. 1984. Zbiorowiska roślinne Antarktyki. *Kosmos* 33: 43–56.
- OCHYRA R. 1986 (1985). On the Antarctic species of the family Orthotrichaceae. *Lindbergia* 11: 141–146.
- OCHYRA R. 1987. *Schistidium steerei* (Grimmiaceae), a remarkable new species from the Antarctic, with a note on *S. obtusifolium*. *Memoirs of the New York Botanical Garden* 45: 607–614.
- OCHYRA R. 1990. The discovery of the South Georgian endemic species *Schistidium urnulaceum* (C. Muell.) B.G. Bell (Musci: Grimmiaceae) in the Antarctic. *Polish Polar Research* 11: 133–146.

- OCHYRA R. 1993a. Antipodal mosses: I. A revision of the genus *Holodontium* (Seligeriaceae). *Fragmenta Floristica et Geobotanica* 38: 75–98.
- OCHYRA R. 1993b. *Grimmia plagiopodia* (Musci, Grimmiaceae) in the Southern Hemisphere. *Fragmenta Floristica et Geobotanica* 38: 21–27.
- OCHYRA R. 1993c. The taxonomic status of *Blindia skottsbergii* (Musci, Seligeriaceae). *Fragmenta Floristica et Geobotanica* 38: 503–508.
- OCHYRA R. 1996a. *Ditrichum lewis-smithii* (Ditrichaceae, Bryopsida), a new species from Antarctica. *Annales Botanici Fennici* 33: 303–309.
- OCHYRA R. 1996b. Nomenclatural notes on mosses: 10. *Andreaea gainii* and *A. parallela* (Andreaeaceae) from Antarctica. *Fragmenta Floristica et Geobotanica* 41: 479–480.
- OCHYRA R. 1996c. One more synonym of *Holodontium strictum* (Musci, Seligeriaceae). *Fragmenta Floristica et Geobotanica* 41: 1011–1014.
- OCHYRA R. 1996d. Rhizoid gemmae in *Ditrichum brotherusii* (Musci, Ditrichaceae) from Antarctica. *Fragmenta Floristica et Geobotanica* 41: 1019–1021.
- OCHYRA R. 1997. Nomenclatural note on mosses: 15. Typification of *Polytrichum subpiliferum* (Polytrichaceae). *Fragmenta Floristica et Geobotanica* 42: 175–176.
- OCHYRA R. 1998a. *Schistidium halinae* (Bryopsida, Grimmiaceae), a new moss species from the Antarctic. *Annales Botanici Fennici* 35: 267–273.
- OCHYRA R. 1998b. *The moss flora of King George Island, Antarctica*. Polish Academy of Sciences, Institute of Botany, Cracow: xxiv + 278 pp.
- OCHYRA R. 1998c. The taxonomic status of *Brachythecium turquetii* (Musci, Brachytheciaceae). *Fragmenta Floristica et Geobotanica* 43: 287–289.
- OCHYRA R. 1999a. Antipodal mosses: IX. *Platydictya* (Bryopsida, Hypnaceae). *Annales Botanici Fennici* 36: 51–58.
- OCHYRA R. 1999b. *Ditrichum austrogeorgicum*: a synonym of *D. hyalinum* (Ditrichaceae, Bryopsida). *Cryptogamie, Bryologie* 20: 247–253.
- OCHYRA R. 1999c. The identities of some forms of Antarctic mosses. *Fragmenta Floristica et Geobotanica* 44: 511–514.
- OCHYRA R. 2003. *Schistidium lewis-smithii* (Bryopsida, Grimmiaceae), a new species from the maritime Antarctic. *Nova Hedwigia* 77: 363–372.
- OCHYRA R. 2004a. Antipodal mosses: XIV. On the taxonomic status of *Grimmia lawiana* (Bryopsida: Grimmiaceae) from the continental Antarctic. *Polish Polar Research* 25: 111–122.
- OCHYRA R. 2004b. Antipodal mosses: XV. Taxonomy and distribution of *Schistidium cupulare* (Bryopsida: Grimmiaceae). *Polish Polar Research* 25: 123–133.
- OCHYRA R. 2004c. *Schistidium leptoneurum* species nova from the South Shetland Islands, Antarctica. *Cryptogamie, Bryologie* 25: 125–130.
- OCHYRA R. 2004d. *Tortula hoppeana*, the correct name for *Desmatodon latifolius* in *Tortula* (Bryopsida, Pottiaceae). *The Bryologist* 107: 497–500.
- OCHYRA R. and BELL B.G. 1984. A record of *Schistidium falcatum* (Bryophyta: Musci) from the Antarctic. *British Antarctic Survey Bulletin* 64: 77–79.
- OCHYRA R. and LEWIS-SMITH R.I. 1996. Antipodal mosses: VI. *Stegonia latifolia* (Musci, Pottiaceae) in the Antarctic. *Fragmenta Floristica et Geobotanica* 41: 985–994.
- OCHYRA R. and LEWIS-SMITH R.I. 1998. Antarctic species in the genus *Ditrichum* (Ditrichaceae, Bryopsida), with a description of *D. gemmiferum* sp. nov. *Annales Botanici Fennici* 35: 33–53.
- OCHYRA R. and LEWIS-SMITH R.I. 1999. *Meesia uliginosa* Hedw. (Musci, Meesiaceae) in Antarctica. *Cryptogamie, Bryologie* 20: 5–10.
- OCHYRA R. and NEWTON M.E. 1986 (1985). The taxonomy and distribution of *Dicranella cardotii* (R. Br. ter.) Dix., an addition to the moss flora of Antarctica. *Lindbergia* 11: 94–98.
- OCHYRA R. and OCHI H. 1986. New or otherwise interesting species of the genus *Bryum* (Musci, Bryaceae) in the Antarctic. *Acta Botanica Hungarica* 32: 209–219.

- OCHYRA R. and SINGH S.M. 2008. Three remarkable moss records from Dronning Maud Land, continental Antarctica. *Nova Hedwigia* 86: 497–506.
- OCHYRA R. and TYSHCHENKO O. 2006. *Leptobryum pyriforme* (Hedw.) Wilson. Antarctic Peninsula. *Journal of Bryology* 28: 151–152.
- OCHYRA R. and VÁŇA J. 1989a. The hepatics of King George Island, South Shetland Islands, Antarctica, with particular reference to the Admiralty Bay region. *Polish Polar Research* 10: 183–210.
- OCHYRA R. and VÁŇA J. 1989b. The hepatics reported from the Antarctic and an outline of their phytogeography. *Polish Polar Research* 10: 211–229.
- OCHYRA R. and ZANDER R.H. 2002. The genera *Didymodon* and *Bryoerythrophyllum* (Pottiaceae) in Antarctica. *Journal of Bryology* 24: 33–44.
- OCHYRA R. and ZANDER R.H. 2007. Czy *Tortula lithophila* jest konspecyficzna z *Sarconeurum glaciale* (Bryopsida: Pottiaceae)? *Fragmenta Floristica et Geobotanica ser. Polonica* 14: 209–212.
- OCHYRA R., BEDNAREK-OCHYRA H. and LEWIS-SMITH R.I. 1998. 170 years of research of the Antarctic moss flora. In: P. Głowacki and J. Bednarek (eds) *Polish Polar Studies. 25th International Polar Symposium*. Warszawa: 159–177.
- OCHYRA R., BEDNAREK-OCHYRA H. and LEWIS-SMITH R.I. 2003. *Schistidium deceptionense*, another new endemic moss species from the Antarctic. *The Bryologist* 106: 569–574.
- OCHYRA R., BEDNAREK-OCHYRA H. and LEWIS-SMITH R.I. 2008a. New and rare moss species from the Antarctic. *Nova Hedwigia* 87: 457–477.
- OCHYRA R., BEDNAREK-OCHYRA H., ARTS T. and LEWIS-SMITH R.I. 2000. Occurrence of the neotropical moss *Dicranella hilariana* (Mont.) Mitt. in the Antarctic. *Tropical Bryology* 18: 153–160.
- OCHYRA R., LEWIS-SMITH R.I. and BEDNAREK-OCHYRA H. 2008b. *The illustrated moss flora of Antarctica*. Cambridge University Press, Cambridge: xvii + 685 pp.
- OCHYRA R., PRZYWARA L. and KUTA E. 1982. Karyological studies on some Antarctic liverworts. *Journal of Bryology* 12: 259–263.
- OCHYRA R., VITT D.H. and HORTON D.G. 1986. An annotated guide to Bryophyta Antarctica Exsiccata. *Cryptogamie. Bryologie, lichénologie* 7: 53–62.
- OLECH M. 1989a. Lichens from the Admiralty Bay region, King George Island (South Shetland Islands, Antarctica). *Acta Societatis Botanicorum Poloniae* 58: 493–512.
- OLECH M. 1989b. Preliminary botanical studies at Bunger Oasis, East Antarctica. *Polish Polar Research* 10: 605–609.
- OLECH M. 1989c. Preliminary botanical studies in Johnsons Dock area (Livingston, Antarctica). *Bulletin of the Polish Academy of Sciences, Biological Sciences* 37: 223–230.
- OLECH M. 1990. Preliminary studies on ornithocoprophilous lichens of the Arctic and Antarctic regions. *Proceedings of the NIPR Symposium on Polar Biology* 3: 218–223.
- OLECH M. 1991a. *Japewia tornoenis* (Lichenes), a species new to the Antarctic. *Polish Botanical Studies* 2: 217–218.
- OLECH M. 1991b. Preliminary observations on the content of heavy metals in thalli of *Usnea anatarctica* Du Rietz (Lichenes) in the vicinity of the “H. Arctowski” Polish Antarctic Station. *Polish Polar Research* 12: 129–131.
- OLECH M. 1993. Lower plants. In: S. Rakusa-Suszczewski (ed.) *The Maritime Antarctic Coastal Ecosystem of Admiralty Bay*. Department of Antarctic Biology, Polish Academy of Sciences, Warsaw: 173–179.
- OLECH M. 1994. Lichenological assessment of the Cape Lions Rump, King George Island, South Shetland Islands; a baseline for monitoring biological changes. *Polish Polar Research* 15: 111–130.

- OLECH M. 1996a. Human impact on terrestrial ecosystems in West Antarctica. *Proceedings of the NIPR Symposium on Polar Biology* 9: 299–306.
- OLECH M. 1996b. Plant colonization and community development on the Ecology glacier moraines. In: M.A. Olech (ed.) *Proceedings of the International Workshop "Antarctic Research on Taxonomy and Ecology of Algae"*, Cracow, Poland – 1st to 3rd October 1996. Institute of Botany, Jagiellonian University, Cracow: 15–17.
- OLECH M. 1997. Polskie osiągnięcia w dziedzinie ochrony przyrody regionów polarnych. *Chrońmy przyrodę ojczystą* 53: 27–34.
- OLECH M. 1998a. Tundra ecosystems of Antarctica. In: A. Gaździcki and K. Jażdżewski (eds) *Polar ecosystems*. *Kosmos* 47: 569–578.
- OLECH M. 1998b. Synanthropization of the flora of Antarctica: an issue. *Phytocoenosis* 10: 269–273.
- OLECH M. 2001. *Annotated checklist of Antarctic lichens and lichenicolous fungi*. The Institute of Botany of the Jagiellonian University, Kraków: 145 pp.
- OLECH M. 2002. Plant communities on King George Island. *Ecological Studies* 154: 215–231.
- OLECH M. 2004. *Lichens of King George Island, Antarctica*. The Institute of Botany of the Jagiellonian University, Kraków: 393 pp.
- OLECH M. and ALSTRUP V. 1990. *Thelocarpon cyaneum* sp. nov. *Nordic Journal of Botany* 9: 575–576.
- OLECH M. and ALSTRUP V. 1995. Lichenicolous fungi from polar regions. In: J. Pereyma and J. Piasecki (eds) *XXII Polar Symposium*. Wrocław-Książ: 119–122.
- OLECH M. and ALSTRUP V. 1996. *Dactylospora dobrowolskii* sp. nov. and additions to the flora of lichens and lichenicolous fungi of Bunge Oasis, East Antarctica. *Polish Polar Research* 17: 165–168.
- OLECH M. and BYSTREK J. 2004. *Bryoria forsteri* (lichenized Ascomycotina), a new species from Antarctica. *Acta Societatis Botanicorum Poloniae* 73: 151–153.
- OLECH M. and CZARNOŃA P. 2009. Two new *Bacidia* (Ramalinaceae, lichenized Ascomycota) from Antarctica. *Polish Polar Research* 30: 339–346.
- OLECH M. and MASSALSKI A. 2001. Plant colonization and community development in the Sphinx Glacier forefield. *Folia Facultatis Scientiarum Naturalium Universitatis Masarykianae Brunensis, Geographia* 25: 111–119.
- OLECH M. and MLECZKO P. 2000. New species and new combination in the fungal genus *Octospora* from Antarctica. *Acta Societatis Botanicorum Poloniae* 69: 277–283.
- OLECH M. and SÖCHTING U. 1993. Four new species of *Caloplaca* from Antarctica. *Lichenologist* 25: 261–269.
- OLECH M., KWIATEK W.M. and DUTKIEWICZ E.M. 1998a. Lead pollution in the Antarctic region. *X-Ray Spectrometry* 27: 232–235.
- OLECH M., MROZIŃSKA T. and MASSALSKI A. 1998b. Algae of South Shetland Islands (West Antarctica). In: *Wyprawy Geograficzne na Spitsbergen, IV Zjazd Geomorfologów Polskich, UMCS Lublin 3–6 czerwca 1998*. Lublin: 225–236.
- OLECH M., OSY CZKA P. and DUTKIEWICZ E.M. 2000. Local environmental pollution with heavy metals in the Admiralty Bay region (South Shetland, Antarctica). In: M. Grześ, K.R. Lankauf and I. Sobota (eds) *Polish Polar Studies. The 27th International Polar Symposium*. Toruń: 99–103.
- OLECH M., SZYMCZYK S. and KAJFOSZ J. 1993. Lokalne zanieczyszczenia ołowiem środowiska w rejonie Antarktyki. *Prace Mineralogiczne* 83: 51–54.
- OSY CZKA P. and OLECH M. 2004. Taxa of the genus *Cladonia* (lichenized fungi) in the region of King George Island (South Shetlands, Antarctica). In: A. Styszyńska and A.A. Marsz (eds) *Polish Polar Studies. XXX Międzynarodowe Sympozjum Polarne*. Gdynia: 285–291.
- OSY CZKA P. and OLECH M. 2005a. Species of the lichen genus *Tephromela* M. Choisy in the Antarctic Region. In: M. Józwiak and R. Kozłowski (eds) *XXXI Polar Symposium with participation visitors from foreign countries, Kielce 12–14 September 2005, Résumé*. Kraków: 102–104.

- OSYCZKA P. and OLECH M. 2005b. The lichen genus *Cladonia* of King George Island, South Shetland Islands, Antarctica. *Polish Polar Research* 26: 107–123.
- OSYCZKA P., DUTKIEWICZ E.M. and OLECH M.A. 2003. PIXE analysis of trace elements in Antarctic lichens *Usnea antarctica*, *Usnea aurantiaco-atra* and mosses *Sanionia uncinata*. In: M.A. Olech (ed.) *XXIX International Polar Symposium*. Kraków: 97–102.
- OSYCZKA P., DUTKIEWICZ E.M. and OLECH M. 2007. Trace elements concentrations in selected moss and lichen species collected within Antarctic research stations. *Polish Journal of Ecology* 55: 39–48.
- OSYCZKA P., KUKWA M. and OLECH M. 2009. Notes on the lichen genus *Lepraria* from maritime (South Shetlands) and continental (Schirmacher and Bunger Oases) Antarctica. *Polar Biology* doi 10.1007/s00300-009-0738-7.
- PIOTROWICZ-CIEŚLAK A., GIELWANOWSKA I., BOCHENEK A., LORO P. and GÓRECKI R. 2005. Carbohydrates in *Colobanthus quitensis* and *Deschampsia antarctica*. *Acta Societatis Botanicorum Poloniae* 74: 209–217.
- PISAREK W., HOŁDYŃSKI C. and LORO P.M. 2003. Vegetation diversity in the Moss Creek Valley (the King George Island, South Shetland Islands, Antarctic). In: M.A. Olech (ed.) *XXIX International Polar Symposium*. Kraków: 103–108.
- PRZYWARA L., KUTA E. and OCHYRA R. 1984. Cytological studies on Antarctic mosses. II. *Journal of the Hattori Botanical Laboratory* 57: 127–137.
- RAKUSA-SUSZCZEWSKI S., JAŹDŹEWSKI K., MYRCHA A. and OLECH M. 1998. Biological and ecological studies carried out at the Polish Antarctic Station *Henryk Arctowski*, 1977–1997. *Polish Polar Research* 19: 37–60.
- RYBALKA N., ANDERSEN R.A., KOSTIKOV I., MOHR K.I., MASSALSKI A., OLECH M. and FRIEDL T. 2009. Testing for endemism, genotypic diversity and species concept in Antarctic terrestrial microalgae of the Tribonemataceae (Stramenopiles, Xanthophyceae). *Environmental Microbiology* 11: 554–565.
- SADOWSKA A. 1998. Pollen morphology of two angiospermous plants from Antarctica – *Colobanthus quitensis* and *Deschampsia antarctica*. *Grana* 37: 58–62.
- SCHROETER B., OLECH M., KAPPEN L. and HEITLAND W. 1995. Ecophysiological investigations of *Usnea antarctica* in the maritime Antarctic. I. Annual microclimatic conditions and potential primary production. *Antarctic Science* 7: 251–260.
- SCHUCH L.A., GODOY J.M., NORDEMANN D.J.R., REIS V.R.G., RAMALHO M., RECIO J.C., BRITO R.R.A. and OLECH M.A. 1993. Radioactivity levels in Antarctic samples. In: P. Strand and E. Holm (eds) *Environmental radioactivity in the Arctic and Antarctic*. Østerås: 211–214.
- SEPPELT R.D. and OCHYRA R. 2008. Moss amongst the ice – the forests of Antarctica. *Field Bryology* 94: 39–43.
- SMYKLA J. 2005. Rola kolonii pingwinów w kształtowaniu składu gatunkowego oraz wzorców przestrzennego zróżnicowania zbiorowisk roślinnych w Antarktyce Morskiej. In: M. Józwiak and R. Kozłowski (eds) *Polish Polar Studies. XXXI Polar Symposium, Kielce 2005*. Kielce: 153–165.
- SMYKLA J. 2008. Strategie życiowe roślin w gradiencie oddziaływania kolonii pingwinów – próba weryfikacji modeli teoretycznych. In: A. Kowalska, A. Lachota, H. Marszałek and J. Pereyma (eds) *Środowisko przyrodnicze obszarów polarnych*. Wydział Nauk o Ziemi i Kształtowania Środowiska Uniwersytetu Wrocławskiego, Wrocław: 208–217.
- SMYKLA J., SZAREK-GWIAZDA E. and KRZEWICKA B. 2005. Trace elements in the lichens *Usnea aurantiaco-atra* and *Usnea antarctica* from the vicinity of Uruguay's Artigas Research Station on King George Island, Maritime Antarctic. *Polish Botanical Studies* 19: 49–57.
- SMYKLA J., WOLEK J. and BARCIKOWSKI A. 2007. Zonation of vegetation related to penguin rookeries on King George Island, Maritime Antarctic. *Arctic, Antarctic and Alpine Research* 39: 143–151.

- SMYKLA J., WOŁEK J., BARCIKOWSKI A. and LORO P. 2006. Vegetation patterns around penguin rookeries at Admiralty Bay, King George Island, Maritime Antarctica: preliminary results. *Polish Botanical Studies* 22: 449–458.
- SØCHTING U. and OLECH M. 1995. The lichen genus *Caloplaca* in polar regions. *Lichenologist* 27: 463–471.
- SØCHTING U. and OLECH M. 2000. *Caloplaca scolecomarginata* spec. nova and *Caloplaca frigida* spec. nova, two new lichen species from Antarctica. *Bibliotheca Lichenologia* 75: 19–26.
- STARMACH K. 1995. Freshwater algae of the Thala Hills oasis (Enderby Land, East Antarctica). *Polish Polar Research* 16: 113–148.
- SZCZUKA E., GIELWANOWSKA I., PIDEK I.A., SETA A., DOMACIUK M. and KOŁODZIEJSKI W. 2008. Pollen of the Antarctic plants *Colobanthus quitensis* and *Deschampsia antarctica* and its representation in moss polsters. *Annales Universitatis Mariae Curie-Skłodowska, Lublin – Polonia, sectio C* 63: 63–70.
- ŚLIWA L. and OLECH M. 2002. Notes on species of *Lecanora* (lichenized Ascomycotina) from the Antarctic. *Polish Polar Research* 23: 117–133.
- TATUR A. and MYRCHA A. 1989. Soils and vegetation in abandoned penguin rookeries (Maritime Antarctic). *NIPR Symposium on Polar Biology* 2: 181–189.
- TATUR A., MYRCHA A. and NIEGODZISZ J. 1997. Formation of abandoned penguin rookery ecosystems in the maritime Antarctic. *Polar Biology* 17: 405–417.
- XAVIER-FILHO L., KOŁAKOWSKA A. and VINCENTE C. 1986. Carotenoid composition of *Caloplaca regalis* (Vain.) Zahlbr. (Lichenes) and of penguin faeces at King George Island (Antarctica). *Polish Polar Research* 7: 289–294.
- ZABAWSKI J. 1995. Fungi of *Penicillium* genus in several soil of King George Island (South Shetlands, the Antarctic). In: J. Pereyma and J. Piasecki (eds) *XXII Polar Symposium*. Wrocław-Książ: 143–149.
- ZABAWSKI J. and PIASECKI J. 1981. Studia nad mikroflorą glebową Wyspy King George (Południowe Szetlandy). In: *Materiały Ogólnopolskiego Seminarium nt. Mikrobiologiczne przemiany związków azotowych w różnych warunkach ekologicznych, Puławy 3–5 V 1981*. Instytut Uprawy Nawożenia i Gleboznawstwa Puławy 2: 131–138.
- ZARZYCKI K. 1993. Vascular plants and terrestrial biotopes. In: S. Rakusa-Suszczewski (ed.) *The Maritime Antarctic coastal ecosystem of Admiralty Bay*. Department of Antarctic Biology, Polish Academy of Sciences, Warsaw: 181–187.

Received 30 September 2010

Accepted 3 March 2011