

CONVERGENCE IN CAPACITY OF THE PLURICOMPLEX GREEN FUNCTION

BY RAFAŁ CZYŻ

Abstract. In this paper we prove that if Ω is a bounded hyperconvex domain in \mathbb{C}^n and if $\Omega \ni z_j \rightarrow \partial\Omega$, $j \rightarrow \infty$, then the pluricomplex Green function $g_\Omega(z_j, \cdot)$ tends to 0 in capacity, as $j \rightarrow \infty$.

A bounded open connected set $\Omega \subset \mathbb{C}^n$ is called hyperconvex if there exists negative plurisubharmonic function $\psi \in PSH(\Omega)$ such that $\{z \in \Omega : \psi(z) < c\} \subset\subset \Omega$ for all $c < 0$. Such ψ is called an exhaustion function for Ω . It was proved in [6] that for every hyperconvex domain there exists smooth exhaustion function ψ such that $\lim_{z \rightarrow \zeta} \psi(z) = 0$, for all $\zeta \in \partial\Omega$.

Let Ω be a bounded hyperconvex domain in \mathbb{C}^n . Let $z \in \Omega$. Recall that the pluricomplex Green function with the pole at z is defined as follows

$$g_\Omega(z, w) = \sup\{u(w) : u \in PSH(\Omega), u \leq 0, |u(\xi) - \log|\xi - z|| \leq C \text{ near } z\}.$$

It is well known that $g_\Omega(z, \cdot) \in PSH(\Omega) \cap \mathcal{C}(\Omega \setminus \{z\})$, $g_\Omega(z, w) = 0$ for $w \in \partial\Omega$ and $(dd^c g_\Omega(z, \cdot))^n = (2\pi)^n \delta_z$, where δ_z is the Dirac measure at z (see [7]). Carlehed, Cegrell and Wikstöm proved in [4] that for every $z_0 \in \partial\Omega$ there exists a pluripolar set $E \subset \Omega$ such that

$$\limsup_{z \rightarrow z_0} g_\Omega(z, w) = 0,$$

for every $w \in \Omega \setminus E$. Blocki and Pflug proved in [3] that if $\Omega \ni z_j \rightarrow \partial\Omega$ then $g_\Omega(z_j, \cdot) \rightarrow 0$ in L^p for every $1 \leq p < +\infty$, as $j \rightarrow \infty$. By $z_j \rightarrow \partial\Omega$ we mean that $\text{dist}(z_j, \partial\Omega) \rightarrow 0$. This result was used in [3] to show Bergman completeness of the hyperconvex domain. Herbort proved in [5] that if a

1991 *Mathematics Subject Classification.* 32U35; 32W20.

Key words and phrases. Pluricomplex Green functions, complex Monge–Ampère operator.

Partially supported by KBN grant no 1 P03A 037 26.

bounded hyperconvex domain $\Omega \subset \mathbb{C}^n$ admits a Hoelder continuous exhaustion function then the pluricomplex Green function $g_\Omega(z_j, \cdot)$ tends to zero uniformly on compact subsets of Ω if the pole $z_j \rightarrow z_0 \in \partial\Omega$. We prove the following theorem.

THEOREM 1. *Let Ω be a bounded hyperconvex domain in \mathbb{C}^n and let $\Omega \ni z_j \rightarrow \partial\Omega$, $j \rightarrow \infty$. Then $g_\Omega(z_j, \cdot) \rightarrow 0$ in capacity as $j \rightarrow \infty$.*

First let us recall the definition of the relative capacity and of convergence in capacity.

DEFINITION 2. The relative capacity of the Borel set $E \subset \Omega \subset \mathbb{C}^n$ with respect Ω is defined in [1]

$$\text{cap}(E, \Omega) = \sup \left\{ \int_E (dd^c u)^n : u \in PSH(\Omega), -1 \leq u \leq 0 \right\}.$$

DEFINITION 3. Let $u_j, u \in PSH(\Omega)$. We say that a sequence u_j converges to u in capacity if for any $\epsilon > 0$ and $K \subset\subset \Omega$

$$\lim_{j \rightarrow \infty} \text{cap}(K \cap \{|u_j - u| > \epsilon\}) = 0.$$

REMARK. Convergence in capacity is stronger then convergence in L^p since the Lebesgue measure $(d\lambda)$ is dominated by the relative capacity, i.e. there exists constant $C(n, \Omega) > 0$ depends only on n and Ω such that

$$\text{cap}(E) \geq C(n, \Omega)\lambda(E).$$

To prove the last inequality observe that there exist constants $C_1, C_2 > 0$ depending only on Ω such that $-1 \leq C_1|z|^2 - C_2 \leq 0$ on Ω and $(dd^c(C_1|z|^2 - C_2))^n = 4^n n! C_1^n d\lambda$. Therefore the above inequality holds with $C(n, \Omega) = 4^n n! C_1^n$. Observe also that uniform convergence on compact sets is stronger then convergence in capacity, since the following inequality holds

$$\text{cap}(K \cap \{|u_j - u| > \epsilon\}) \leq \epsilon^{-1} \text{cap}(K) \sup_K |u_j - u|.$$

To prove Theorem 1 we will need the following lemma proved in [2].

LEMMA 4. *Let Ω be a bounded domain \mathbb{C}^n . Assume that u, v are bounded negative plurisubharmonic functions such that $\lim_{z \rightarrow \zeta} v(z) = 0$, for all $\zeta \in \partial\Omega$. Then*

$$\int_\Omega (-v)^n (dd^c u)^n \leq n! (\sup_\Omega |u|)^{n-1} \int_\Omega (-u) (dd^c v)^n.$$

PROOF OF THEOREM 1. Let us denote $u_j = g_\Omega(z_j, \cdot)$. Suppose that u_j does not converge in capacity to 0, $j \rightarrow \infty$. Then for some $\epsilon > 0$ and $K \subset\subset \Omega$

there exist a subsequence u_{j_k} , and constants $c > 0$ and $N > 0$ such that for $j_k \geq N$ we have

$$(1) \quad \text{cap}(K \cap \{-u_{j_k} > \epsilon\}) \geq c.$$

From the definition of capacity there exists $v \in PSH(\Omega)$ such that $-1 \leq v \leq 0$ and

$$(2) \quad \int_{K \cap \{-u_{j_k} > \epsilon\}} (dd^c v)^n \geq \frac{c}{2}.$$

Now we will show that $u_j \rightarrow 0$ on K in $L^n((dd^c v)^n)$. Since Ω is hyperconvex then there exist ψ a continuous exhaustion function for Ω and a constant $A > 0$ such that $A\psi < v$ on K . Define the following bounded plurisubharmonic function $\varphi = \max(A\psi, v)$. Then $\lim_{z \rightarrow \zeta} \varphi(z) = 0$, for all $\zeta \in \partial\Omega$ and

$$(dd^c \varphi)^n \geq \chi_K (dd^c v)^n,$$

where χ_K is the characteristic function of the set K . Observe that φ is an exhaustion function for Ω , which implies that $\varphi(z_j) \rightarrow 0$ if $\text{dist}(z_j, \partial\Omega) \rightarrow 0$.

Using the monotone convergence theorem and Lemma 4 we get

$$\begin{aligned} \int_K (-u_j)^n (dd^c v)^n &= \int_{\Omega} (-u_j)^n (dd^c \varphi)^n = \lim_{k \rightarrow +\infty} \int_{\Omega} (-\max(u_j, -k))^n (dd^c \varphi)^n \\ &\leq n! (\sup_{\Omega} |\varphi|)^{n-1} \lim_{k \rightarrow +\infty} \int_{\Omega} |\varphi| (dd^c \max(u_j, -k))^n = n! (2\pi)^n (\sup_{\Omega} |\varphi|)^{n-1} |\varphi(z_j)|, \end{aligned}$$

which means that $u_j \rightarrow 0$ on K in $L^n((dd^c v)^n)$, since $\varphi(z_j) \rightarrow 0$, as $j \rightarrow \infty$.

Observe that inequality (2) implies that

$$\frac{c}{2} \leq \int_{K \cap \{-u_{j_k} > \epsilon\}} (dd^c v)^n \leq \epsilon^{-n} \int_K (-u_{j_k})^n (dd^c v)^n,$$

which is impossible since $u_{j_k} \rightarrow 0$ on K in $L^n((dd^c v)^n)$. This means that $u_j \rightarrow 0$ in capacity as $j \rightarrow \infty$. The proof is finished. \square

Now we recall the definition of the multipolar Green function introduced by Lelong [8]. Let $A = \{(z^{(1)}, \nu^{(1)}), \dots, (z^{(m)}, \nu^{(m)})\}$ be a finite subset of $\Omega \times \mathbb{R}_+$. Let

$$g_{\Omega}(A, w) = \sup\{u(w) : u \in \mathcal{L}_A, u \leq 0\},$$

where \mathcal{L}_A denotes the family of plurisubharmonic functions on Ω having a logarithmic pole with weight $\nu^{(k)}$ at $w^{(k)}$, for $k = 1, \dots, m$, i.e.

$$\mathcal{L}_A = \{u \in PSH(\Omega) : |u(\xi) - \nu^{(j)} \log |\xi - z^{(j)}|| \leq C_j \text{ near } z^{(j)}, 1 \leq j \leq m\}.$$

We show that it is possible to generalize Theorem 1 for the multipolar Green function.

COROLLARY 5. *Let Ω be a bounded hyperconvex domain in \mathbb{C}^n and let $A_j = \{(z_j^{(1)}, \nu^{(1)}), \dots, (z_j^{(m)}, \nu^{(m)})\}$ be a subset of $\Omega \times \mathbb{R}_+$, for $j = 1, 2, \dots$, such that $\Omega \ni z_j^{(k)} \rightarrow \partial\Omega$, $j \rightarrow \infty$ for all $k = 1, \dots, m$. Then $g_\Omega(A_j, \cdot) \rightarrow 0$ in capacity as $j \rightarrow \infty$.*

PROOF. Directly from the definition of the multipolar Green function we have

$$\sum_{k=1}^m \nu^{(k)} g_\Omega(z_j^{(k)}, \cdot) \leq g_\Omega(A_j, \cdot) \leq 0.$$

By Theorem 1 we have that $g_\Omega(z_j^{(k)}, \cdot) \rightarrow 0$ in capacity as $j \rightarrow \infty$ for all $k = 1, \dots, m$, so also $g_\Omega(A_j, \cdot) \rightarrow 0$ in capacity as $j \rightarrow \infty$. This ends the proof. \square

References

1. Bedford E., Taylor B.A., *A new capacity for plurisubharmonic functions*, Acta Math., **149** (1982), 1–40.
2. Blocki Z., *Estimates for the complex Monge–Ampère operator*, Bull. Pol. Acad. Sci. Math., **41** (1993), 151–157.
3. Blocki Z., Pflug P., *Hyperconvexity and Bergman completeness*, Nagoya Math. J., **151** (1998), 221–225.
4. Carlehed M., Cegrell U., Wikström F., *Jensen measures, hyperconvexity and boundary behavior of the pluricomplex Green function*, Ann. Polon. Math., **41** (1999), 87–103.
5. Herbot G., *The pluricomplex Green function on pseudoconvex domains with a smooth boundary*, Int. J. Math., **11** (2000), 509–522.
6. Kerzman N., Rosay J.P., *Fonctions plurisousharmoniques d'exhaustion bornées et domaines tauts*, Math. Ann., **257** (1981), 171–184.
7. Klimek M., *Pluripotential theory*, London Mathematical Society Monographs, Clarendon Press, Oxford – New York – Tokyo, 1991.
8. Lelong P., *Fonction de Green pluricomplexe et lemme de Schwarz dans les espaces de Banach*, J. Math. Pures Appl., **68** (1989), 319–347.

Received December 1, 2005

Jagiellonian University
Institute of Mathematics
ul. Reymonta 4
30-059 Kraków
Poland
e-mail: rafal.czyz@im.uj.edu.pl