



Saponins as cytotoxic agents: an update (2010–2018). Part I—steroidal saponins

Danuta Sobolewska · Agnieszka Galanty · Karolina Grabowska · Justyna Makowska-Wąs · Dagmara Wróbel-Biedrawa · Irma Podolak



Received: 5 November 2019 / Accepted: 28 January 2020 / Published online: 22 February 2020
© The Author(s) 2020

Abstract Steroidal saponins are a group of glycosides widely distributed among monocotyledonous families. They exert a wide spectrum of biological effects including cytotoxic and antitumor properties which are the most studied. This review is an update of our previous paper—*Saponins as cytotoxic agents* (Podolak et al. in *Phytochem Rev* 9:425–474, 2010) and covers studies that were since published (2010–2018). In this paper we refer to steroidal saponins presenting results of cytotoxicity studies, mechanisms of action and structure–activity relationships.

Keywords Cytotoxicity · Steroidal glycosides · Mechanisms of action

Introduction

Steroidal saponins are an important group of glycosidic plant metabolites. They are mainly distributed among monocotyledonous families: Amarillidaceae (*Agapanthus*, *Allium*), Asparagaceae (*Agave*, *Anemarrhena*, *Asparagus*, *Convallaria*, *Hosta*, *Nolina*, *Ophiopogon*, *Ornithogalum*, *Polygonatum*, *Ruscus*, *Sansevieria*, *Tupistra*, *Yucca*), Costaceae (*Costus*), Dioscoreaceae (*Dioscorea*), Liliaceae (*Fritillaria*, *Lilium*), Melanthiaceae (*Paris*), Smilacaceae (*Smilax*). Although it is uncommon, steroidal saponins can also be found in some dicotyledonous angiosperms, such as: Fabaceae (*Trigonella*), Zygophyllaceae (*Tribulus*, *Zygophyllum*), Solanaceae (*Solanum*, *Lycopersicon*, *Capsicum*), Asteraceae (*Vernonia*), and Plantaginaceae (*Digitalis*) (Faizal and Geelen 2013; Rahman et al. 2017; Lanzotti 2005; Sobolewska et al. 2016; Tang et al. 2013; Wang et al. 2018). Moreover, these compounds have been identified in starfish and marine sponges (Ivanchina et al. 2011; Barnett et al. 1988; Regaldo et al. 2010).

Structurally, steroidal saponins are distinguished by the nature of the aglycone part. Sapogenins are polycyclic 27-C-compounds which can be divided into three distinct groups: spirostane, furostane, and open-chain (cholestane) compounds (Challinor and De Voss 2013). Some authors distinguish iso-spirostane-type saponins—possessing an equatorial oriented (hydroxy)methyl on F ring *versus* spirostane-type with an axial oriented C-27 group (Tian et al. 2017).

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s11101-020-09661-0>) contains supplementary material, which is available to authorized users.

D. Sobolewska (✉) · A. Galanty · K. Grabowska · J. Makowska-Wąs · D. Wróbel-Biedrawa · I. Podolak
Department of Pharmacognosy, Jagiellonian University, Medical College, 9 Medyczna Street, Kraków, Poland
e-mail: dsobolew@cm-uj.krakow.pl

Furthermore, spirosolane-type glycoalkaloids in which a nitrogen atom is incorporated in the steroid aglycone at the heterocyclic oxygen site (e.g. in solasodine) are sometimes included in the group of steroidal saponins. The sugar residue of steroidal saponins consists of one to three straight or branched sugar chains, which are composed usually of β -D-glucopyranosyl (Glc), α -L-rhamnopyranosyl (Rha), β -D-galactopyranosyl (Gal), β -L-arabinofuranosyl (Ara), β -D-xylopyranosyl (Xyl), β -D-fucopyranosyl (Fuc), β -D-mannopyranosyl (Man), or β -D-quinovopyranosyl (Qui) residues.

Since many years spirostanol saponins, such as e.g. disogenin or hecogenin, have been valued by pharmaceutical industry and used as substrates in the production of steroid hormones and drugs. Also, medicinal properties of saponin containing plants are well known. Some of most prominent examples include *Ruscus aculeatus*, which is used as vasoprotective agent, or *Tribulus terrestris*, found in many products dedicated to fertility stimulation in men (Masullo et al. 2016; Salgado et al. 2017). Steroidal saponins are a research target of many scientist groups. Numerous published reports have confirmed that these compounds exert a wide spectrum of pharmacological activities, including antimicrobial, anti-inflammatory, cardioprotective, cAMP phosphodiesterase inhibitory, or anti-adipogenic (Sohn et al. 2006; Huang et al. 2013; Tang et al. 2015; Ning et al. 2010; Nakamura et al. 1993; Poudel et al. 2014).

One of the activities that is especially widely explored is cytotoxic effect (Podolak et al. 2010; Böttger et al. 2012). The search for potential new chemotherapeutics within natural sources is obviously triggered by a growing need to provide effective treatment to counteract cancer. Results of studies on in vitro and in vivo cytotoxicity of steroidal saponins indicate that these compounds provide an interesting research target. In our previous review the results of experimental studies on cytotoxicity of saponins, both triterpene and steroidal, covering the period from 2005 to 2009 have been summarized (Podolak et al. 2010). Since then, a vast number of new experimental data have appeared in literature. This issue is however scarcely reviewed. Several papers that discussed biological activities of compounds found in a particular genus, like e.g. *Allium* or *Smilax*, (Sobolewska et al. 2016; Tian et al. 2017), referred also to their cytotoxic effects, but there are virtually

none reviews focused entirely on this activity despite a growing number of reports with experimental data. Some more general aspects were tackled by Xu et al. (2016) who discussed anticancer saponins from Chinese plants. In a recent paper by Zhao et al. (2018), advances in antitumor potential of steroidal saponins have been focused on the mechanisms of action, and included examples of saponinogenins and saponins, as well as some other compounds such as a cardiac glycoside—bufalin or cucurbitacins.

Taking into account a large number of experimental data referring to cytotoxicity of saponins, that have been published since our previous review, we decided to divide this update into two parts, each dedicated to the one of the distinct structural groups, that is triterpene and steroidal compounds.

Thus, in the current review, we present an update on the cytotoxic activity of steroidal saponins and saponinogenins covering recent studies from the period of 2010 to 2018. Discussion of structure–activity data and mechanisms of action is also provided, together with a selection of most promising compounds with a potential for future development as anticancer chemotherapeutics.

The literature search was conducted in the following electronic databases: SCOPUS, EMBASE and MEDLINE/PubMed. The keywords used were: steroidal saponins, steroidal saponinogenins, cancer, cytotoxicity.

Since 2010 year in vitro cytotoxicity studies have been performed on different human and animal cell cancer and normal lines, including:

- *Human cancer cell lines* Breast: BT-549, MCF-7, MDA-MB-231, MDA-MB-435, MDA-MB-468, SK-BR-3; bone: 143-B, HOS; cervix: HeLa, Caski, KB, SiHa; colon: COLO, DLD-1, HT-29, HCT 116, HCT-15, CaCo-2, SW480, SW620, W480, LOVO; esophagus: KYSE 510; gingival: Ca9-22; glioblastoma: SF-268, SF-295, U251, U87MG; larynx: Hep2; leukemia: CCRF-CEM, HL-60, Jurkat, K562; liver: HLE, Hep3B, HepG2, HuH-7, C3A, BEL-7402, BEL-7403, BEL-7404, MHCC97-L, SMMC-7721, SMMC-7221, SNU-387, WRL; lung: 95D, A549, LAC, NCI-H1299, NCI-H446, NCI-H460, SK-MES-1; melanoma: A375, A375.S2, MM96L, SK-MEL, SK-MEL-2, WM-115; neuroblastoma: IMR-32, LA-N-2, NB-69; ovary: HO-8910PM, OVCAR-8, SK-OV-3;

- pancreas: BxPC-3, PANC-1; pharynx: 5-8F, CNE; prostate: DU145, PC-3; sarcoma: MG-63, Rh1; stomach: BGC-823, SGC-7901, SGC-7901/DDP [cisplatin (DDP)-resistant], HIF1 α -knockdown BGC-823 (hypoxia-mimic sensitive), MGC-803; urinary bladder: ECV-304;
- *Animal cancer cell lines* Breast: EMT6; glioblastoma: C6; leukemia: Baf3-WT; lung: LL2; colon: C26; melanoma: B16; sarcoma: WEHI-164, J-774;
 - *Human normal cell lines* Fibroblasts: HFF, NFF, Hs68; keratinocytes: HaCaT; kidney embryonic: HEK293; lung epithelial: MRS-5; vein endothelial: EA.hy926, HUVEC;
 - *Animal normal cell lines* Cardiomyoblasts: H9c2; epidermal: JB6 P⁺Cl-41; fibroblasts: 3T3; kidney epithelial: LLC-PK1; kidney fibroblasts: VERO.

The results of these studies have been summarized in Table 1.

Based on data published in years 2010–2018 it may be concluded that out of 284 substances that are included in the current review, a vast majority, that is 96.8%, were pure single compounds, both saponins and sapogenins, either structurally novel or known previously. A graph representing a number of tested substances and a number of reports published in the time scope covered by this review (2010–2018) is shown in Fig. 1.

Cytotoxicity studies were performed on animal and human cell line models, with significant predominance of the latter, which accounted to 92.7% of all assays. The effects of steroidal saponins/sapogenins against human colon, breast and liver cancers have been most widely studied, accounting to 17.9%, 16.5% and 16% of all assays on human cell lines, respectively. A graph showing the share of experiments on specific types of tumors and normal cell lines in the total number of tests performed on human cell lines is shown in Fig. 2.

The largest number of substances was analysed against following cell lines: HepG2—human hepatocellular carcinoma, MCF-7—human breast adenocarcinoma, and A549—human lung adenocarcinoma cells, which constituted 27.8%, 27.4% and 23.5% of the pool of substances under study, respectively. Tests in which normal cell lines were included in the study accounted for only 4.4% of all assays conducted on human cell lines.

The most preferred method used was the MTT assay. In most cases (80.4% of all assays) IC₅₀ values for analysed saponins were compared with a positive control. Well-known anticancer drugs such as doxorubicin, cisplatin and paclitaxel were most frequently used as reference substances. Other compounds with anticancer activity were chosen definitely less often and these include: actinomycin D, adriamycin, beta-L-(–)-dioxalane-cytidine (–)-OddC, camptothecin, elipticin, etoposide, 5-FU, mitamycin C, mitoxantrone, nimustine (ACNU), podophyllotoxin, staurosporine, tamoxifen, and troxacitabine. In one study resveratrol, which is not an approved anticancer drug, served as the control (Shen et al. 2012).

In the majority of cases steroidal saponins were less active than the control substances. However, there were some noticeable examples of compounds which displayed cytotoxic effect higher than the reference drug. Three saponins isolated from *Dracaena cambodiana* dragon's blood that are glycosides of diosgenin, pennogenin and spirost-5,25(27)-dien-1 β ,3 β -diol, exerted stronger cytotoxic activity (IC₅₀: 1.27 μ M, 5.09 μ M, 4.77 μ M, respectively) against K-562 cells than paclitaxel (IC₅₀: 5.98 μ M), while a pennogenin glycoside showed higher cytotoxic effect on BEL-7402 than paclitaxel (IC₅₀: 1.13 μ M and 3.75 μ M respectively) (Shen et al. 2014). Results obtained by Teponno et al. showed that a well known steroidal glycoside—dioscin—diosgenin 3-*O*- α -L-Rha-(1 \rightarrow 4)-[α -L-Rha-(1 \rightarrow 2)]- β -D-Glc (for the purpose of the study isolated from *Dracaena viridiflora*) had cytotoxic activity against Jurkat, Caco-2, SK-OV-3, and A549 cells (IC₅₀: 1.70 \pm 0.38 μ g ml⁻¹, 2.58 \pm 0.21 μ g ml⁻¹, 1.90 \pm 0.86 μ g ml⁻¹, and 0.42 \pm 0.15 μ g ml⁻¹, respectively) comparable to doxorubicin used as positive control (IC₅₀: 0.61 \pm 0.04 μ g ml⁻¹, 2.32 \pm 1.04 μ g ml⁻¹, 0.84 \pm 0.08 μ g ml⁻¹, and 1.15 \pm 0.84 μ g ml⁻¹, respectively) (Teponno et al. 2017). Another diosgenin derivative named SAP-1016 (diosgenin 3-*O*- β -D-Xyl-(1 \rightarrow 3)- β -D-Glc-(1 \rightarrow 4)-[α -L-Rha-(1 \rightarrow 2)]- β -D-Glc) which was found in the fruits and roots of *Balanites aegyptiaca* showed potent antiproliferative activity against MCF-7 and HT-29 cancer cells (IC₅₀: 2.4 \pm 0.35 and 3.3 \pm 0.19 μ M, respectively) higher than dioscin (IC₅₀: 3.1 \pm 0.39 μ M and 4.9 \pm 0.32 μ M, respectively) and cisplatin (IC₅₀: 30.3 \pm 0.33 μ M and 40.2 \pm 0.44 μ M, respectively) (Beit-Yannai et al. 2011).

Table 1 Cytotoxic steroidal saponins/sapogenins (2010–2018)

Source	Compound	Cell line	Concentration	Assay	References
<i>Agave sisalana</i> leaves	Hecogenin 3- <i>O</i> - β -D-Glc-(1 \rightarrow 2)-[β -D-Xyl-(1 \rightarrow 3)]- β -D-Glc-(1 \rightarrow 4)- β -D-Gal	NCI-H460	IC ₅₀ (μ M) 5.3 \pm 1.8	Methylene blue dye assay	Chen et al. (2011b)
		MCF-7	11.9 \pm 2.6		
		SF-268	4.0 \pm 2.2		
	Hecogenin 3- <i>O</i> - α -L-Rha-(1 \rightarrow 3)- β -D-Xyl-(1 \rightarrow 2)-[β -D-Xyl-(1 \rightarrow 3)]- β -D-Glc-(1 \rightarrow 3)]- β -D-Glc-(1 \rightarrow 4)- β -D-Gal	NCI-H460	6.5 \pm 1.1		
		MCF-7	9.5 \pm 4.8		
		SF-268	8.2 \pm 1.2		
	Polianthoside E	NCI-H460	> 20		
		MCF-7	> 20		
		SF-268	7.5 \pm 1.4		
	Neotigogenin 3- <i>O</i> - α -L-Rha-(1 \rightarrow 4)- β -D-Glc-(1 \rightarrow 2)-[β -D-Xyl-(1 \rightarrow 3)]- β -D-Glc-(1 \rightarrow 3)]- β -D-Glc-(1 \rightarrow 4)- β -D-Gal	NCI-H460	3.8 \pm 2.7		
		MCF-7	1.2 \pm 0.1		
		SF-268	1.5 \pm 0.8		
	Actinomycin D (control)	NCI-H460	2.6 \pm 1.6		
		MCF-7	31.1 \pm 2.9		
SF-268		7.5 \pm 5.2			
<i>Allium flavum</i> whole plant	Yuccagenin 3- <i>O</i> - β -D-Xyl-(1 \rightarrow 3)-[β -D-Gal-(1 \rightarrow 2)]- <i>O</i> - β -D-Gal-(1 \rightarrow 4)- <i>O</i> - β -D-Gal	SW480	IC ₅₀ (μ M) 14.3	XTT assay	Rezgui et al. (2014)
		SW480	14		
		SW480	18.1		
		SW480	1.47		
		SW480	1.47		
<i>Allium nigrum</i> bulbs	Nigrosides A1/A2	HCT 116	IC ₅₀ (μ M) 47.8	MTT assay	Jabrane et al. (2011)
		HT-29	70.8		
	Aginoside/turoside A	HCT 116	1.59		
		HT-29	1.09		
	25(R,S)-5 α -spirostan-2 α ,3 β ,6 β -trio1 3- <i>O</i> - β -D-Glc-(1 \rightarrow 2)- <i>O</i> -[4- <i>O</i> -(3S)-3-hydroxy-3-methylglutaryl]- β -D-Xyl-(1 \rightarrow 3)]- <i>O</i> - β -D-Glc-(1 \rightarrow 4)- β -D-Gal	HCT 116	3.45		
		HT-29	2.82		
	Paclitaxel (control)	HCT 116	0.00321		
		HT-29	0.0014		

Table 1 continued

Source	Compound	Cell line	Concentration	Assay	References	
<i>Allium schoenoprasum</i> whole plant			IC ₅₀ (μM)	MTT assay	Timité et al. (2013)	
		(25R)-5α-spirostane-3β,11α-diol 3-O-β-D-Glc-(1 → 3)-[β-D-Glc-(1 → 4)]-O-β-D-Gal	HCT 116	8.45		
			HT-29	8.64		
		Laxogenin 3-O-α-L-Rha-(1 → 2)-O-β-D-Glc	HCT 116	> 100		
			HT-29	> 100		
		Deltonin	HCT 116	0.4		
			HT-29	0.75		
		Deltoside	HCT 116	1.58		
			HT-29	1.56		
		Paclitaxel (control)	HCT 116	0.00275		
	HT-29	0.00206				
<i>Allium vavilovii</i> bulbs			IC ₅₀ (μg ml ⁻¹)	MTT assay	Zolfaghari et al. (2013)	
		Vavilosides A1/A2	J-774	5.1		
			WEHI-164	4.7		
		Vavilosides B1/B2	J-774	3.5		
			WEHI-164	3.1		
		Ascalonicosides A1/A2	J-774	4		
	WEHI-164	3.7				
<i>Anemarrhena asphodeloides</i> rhizomes			IC ₅₀ (μM)	SRB assay	Kang et al. (2011)	
		Timosaponin AIII	HCT-15	6.1		
			HCT 116	5.5		
			HT-29	10.3		
			SW480	13.1		
			SW620	11.1		
			normal lung epithelial (MRS-5) cells	> 50		
			fibroblast (Hs68) cells	> 50		
<i>Anemarrhena asphodeloides</i> rhizomes			IC ₅₀ (μM)	SRB assay	Guo et al. (2015)	
		Timosaponin BI	HT-29	14.3		
			HeLa	12.29		
			MDA-MB-468	4.5		
		Timosaponin BII	BEL-7402	2.01		
			HT-29	1.65		
			MDA-MB-468	5.5		
		Timosaponin AIII	BEL-7402	1.65		

Table 1 continued

Source	Compound	Cell line	Concentration	Assay	References				
<i>Anemarrhena asphodeloides</i> rhizomes	Anemarsaponin F Schidigerasaponin F2 Doxorubicin (control)	HT-29	2.2	MTT assay	Yang et al. (2017)				
		HeLa	9.63						
		MDA-MB-468	1.6						
		HT-29	4.04						
		HT-29	9.42						
		BEL-7402	0.3						
		HT-29	0.46						
		HeLa	6.91						
		MDA-MB-468	0.28						
		IC ₅₀ (μM)							
<i>Anemarrhena asphodeloides</i> rhizomes	Anemarsaponin R Timosaponin E1 Doxorubicin (control)	HepG2	43.90 ± 3.36	MTT assay	Zhang et al. (2017)				
		SGC7901	57.90 ± 2.88						
		HepG2	8.20 ± 1.25						
		SGC7901	6.25 ± 2.18						
		IC ₅₀ (μM)							
		<i>Anemarrhena asphodeloides</i> rhizomes	Schidigerasaponin F2			MCF-7	98 ± 8.98	MTT assay	Yang et al. (2018)
						SW480	97.02 ± 14.99		
						HepG2	> 100		
						SGC7901	> 100		
						Anemarsaponin F	MCF-7		
SW480	5.56 ± 1.50								
HepG2	11.73 ± 1.24								
SGC7901	8.18 ± 0.26								
Timosaponin AI	MCF-7			6.83 ± 1.99					
	SW480			4.17 ± 0.72					
	HepG2	7.83 ± 1.72							
	SGC7901	4.38 ± 0.50							
Timosaponin AIII (control)	MCF-7	3.34 ± 1.10							
	SW480	2.94 ± 1.05							
	HepG2	4.96 ± 0.93							
	SGC7901	12.15 ± 1.36							
<i>Anemarrhena asphodeloides</i> rhizomes	Aneglycoside A	HepG2	38.4 ± 2.4	MTT assay	Yang et al. (2018)				
		HeLa	29.7 ± 01.9						
		SGC7901	> 100						
		Aneglycoside B	HepG2			41.8 ± 3.5			
			HeLa			34.2 ± 3.6			
			SGC7901			> 100			
		Timosaponin U	HepG2			61.8 ± 4.1			

Table 1 continued

Source	Compound	Cell line	Concentration	Assay	References				
<i>Archaster typicus</i> starfish	Doxorubicin (control)	HeLa	39.7 ± 3.7	MTS assay	Kicha et al. (2010)				
		SGC7901	44.5 ± 2.0						
		HepG2	8.4 ± 2.2						
		HeLa	9.0 ± 1.4						
		SGC7901	6.7 ± 1.8						
	Archasteroside A	HeLa	24						
		JB6 P+ Cl41	37						
	Archasteroside B	HeLa	14						
		JB6 P+ Cl41	18						
	Regularoside A	HeLa	110						
JB6P+Cl41		> 50							
<i>Asparagus filicinus</i> roots	Filiasparsoside A Filiasparsoside B Filiasparsoside C Filiasparsoside E Filiasparsoside F Filiasparsoside G Asparagusin A Aspafilioside A Aspafilioside B Staurosporine (control)	MDA-MB-231	19.8 ± 1.3	MTT assay	Wu et al. (2010)				
		MDA-MB-231	> 50						
		MDA-MB-231	3.4 ± 0.2						
		MDA-MB-231	> 50						
		MDA-MB-231	> 50						
		MDA-MB-231	> 50						
		MDA-MB-231	> 50						
		MDA-MB-231	6.6 ± 0.3						
		MDA-MB-231	5.3 ± 0.4						
		MDA-MB-231	0.0145 ± 0.0004						
		<i>Aspidistra elatior</i> rhizomes	(25R)-26-O-β-D-Glc-furost-5,20-dien-3β,26-diol-3-O-β-D-Glc (1 → 2)-[β-D-Glc-(1 → 3)]-β-D-Glc-(1 → 4)-β-D-Gal			A549	3.8	MTT assay	Zuo et al. (2018)
						Caski	7.2		
						HepG2	8.2		
MCF-7	10.7								
Aspidsaponin A	A549			5.1					
	Caski			8.6					
	HepG2			11.1					
Adriamycin (control)	MCF-7			13.8					
	A549			1.4					
	Caski			1.5					
	HepG2	0.7							
<i>Avena sativa</i> bran	Avenacoside B	MCF-7	1.7	MTT assay	Yang et al. (2016b)				
		HCT 116	175.3						

Table 1 continued

Source	Compound	Cell line	Concentration	Assay	References
<i>Balanites aegyptiaca</i> fruits and roots			IC ₅₀ (μM)	MTT assay	Beit-Yannai et al. (2011)
			SAP-1016	MCF-7	2.4 ± 0.35
				HT-29	3.3 ± 0.19
				HFF	2.1 ± 0.16
			SAP-884	MCF-7	4.3 ± 0.18
				HT-29	7.6 ± 0.17
				HFF	5.2 ± 0.32
			KE-1046	MCF-7	5.3 ± 0.26
				HT-29	10.4 ± 0.11
				HFF	10.3 ± 0.18
			KE-1064	MCF-7	5.1 ± 0.28
				HT-29	7.8 ± 0.32
				HFF	7.4 ± 0.20
			Diosgenin	MCF-7	28.1 ± 0.52
				HT-29	30.6 ± 0.33
				HFF	20.7 ± 0.45
			Dioscin (control)	MCF-7	3.1 ± 0.39
				HT-29	4.9 ± 0.32
				HFF	2.8 ± 0.19
			Cisplatin (control)	MCF-7	30.3 ± 0.33
				HT-29	40.2 ± 0.44
	HFF	20.6 ± 0.30			
<i>Bletilla striata</i> roots			IC ₅₀ (μM)	SRB assay	Park et al. (2014)
			Bletilnoside A	A549	4.56 ± 0.29
				SK-OV-3	4.00 ± 0.06
				SK-MEL-2	3.98 ± 0.16
				HCT-15	5.08 ± 0.51
			Bletilnoside B	A549	8.79 ± 1.01
				SK-OV-3	8.08 ± 0.83
				SK-MEL-2	5.29 ± 0.34
				HCT-15	9.29 ± 1.23
			3-O-β-D-Glc-3-Epiruscogenin	A549	12.10 ± 0.40
				SK-OV-3	11.80 ± 0.28
				SK-MEL-2	11.55 ± 0.27
				HCT-15	11.00 ± 0.23
			Doxorubicin (control)	A549	0.0035 ± 0.0025
				SK-OV-3	0.0037 ± 0.0022
	SK-MEL-2	0.0009 ± 0.0001			
	HCT-15	0.1574 ± 0.0569			

Table 1 continued

Source	Compound	Cell line	Concentration	Assay	References	
<i>Bletilla striata</i> roots	(1 α ,3 α)-1- <i>O</i> -[(β -D-Xyl-(1 \rightarrow 2)- α -L-Rha)]-3- <i>O</i> -D-Glc-5 α -spirostan	A549	12.3	MTT assay	Wang and Meng (2015)	
		BGC-823	15.9			
		HepG2	14.3			
		HL-60	17			
		MCF-7	15.1			
		SMMC-7721	14.7			
		W480	17.1			
		(1 α ,3 α)-1- <i>O</i> -[(β -D-Xyl-(1 \rightarrow 2)- α -L-Rha)oxy]-3- <i>O</i> -D-Glc-25(27)-ene-5 α -spirostan	A549			12.7
			BGC-823			12.2
			HepG2			12.8
			HL-60			13.8
			MCF-7			11.3
			SMMC-7721			11.7
		(1 α ,3 α)-1- <i>O</i> -[(β -D-Xyl-(1 \rightarrow 2)- α -L-Rha)oxy]-epiruscogenin	W480			18.3
	A549		24.3			
	BGC-823		29.4			
	HepG2		30.1			
	HL-60		31.1			
	MCF-7		30.4			
	(1 α ,3 α)-1- <i>O</i> -[(β -D-Xyl-(1 \rightarrow 2)- α -L-Rha)oxy]-epineoruscogenin	SMMC-7721	29.7			
		W480	29.1			
		A549	29.7			
		BGC-823	29.6			
		HepG2	29.4			
		HL-60	29.4			
		MCF-7	27.1			
		SMMC-7721	30.1			
	Bletilinoside A	W480	24.9			
A549		76.3				
BGC-823		68.7				
HepG2		66.9				
HL-60		72.3				
MCF-7		76.2				
SMMC-7721		70.8				
3- <i>O</i> - β -D-Glc-3-epi-neoruscogenin	W480	69.4				
	A549	31.9				
	BGC-823	31.2				
	HepG2	30.7				
	HL-60	32.2				
	MCF-7	28.1				
	SMMC-7721	29.9				
W480	27.6					

Table 1 continued

Source	Compound	Cell line	Concentration	Assay	References
<i>Cestrum laevigatum</i> leaves	(25R)-Spirost-5-ene-3 β ,26 β -diol 3- <i>O</i> - α -L-Rha-(1 \rightarrow 4)- α -L-Rha-(1 \rightarrow 4)-[(1 \rightarrow 2)- α -L-Rha]- β -D-Glc	HL-60	6.5 (5.2–8.1)	MTT assay	Ribeiro et al. (2016a)
		OVCAR-8	10.3 (5.4–19.9)		
		HCT 116	10.1 (4.5–23.0)		
		SF-295	7.7 (4.2–14.1)		
	(25R)-Spirost-6-ene-3 β ,5 β -diol 3- <i>O</i> - α -L-Rha-(1 \rightarrow 4)- α -L-Rha-(1 \rightarrow 4)-[(1 \rightarrow 2)- α -L-Rha]- β -D-Glc	HL-60	7.3 (6.7–7.9)		
		OVCAR-8	15.3 (11.9–19.6)		
		HCT 116	11.4 (9.6–13.5)		
	Diosgenin 3- <i>O</i> - α -L-Rha-(1 \rightarrow 4)- α -L-Rha-(1 \rightarrow 4)- β -D-Glc	HL-60	> 25		
		OVCAR-8	> 25		
		HCT 116	> 25		
	Chonglouoside SL-5	HL-60	8.2 (7.4–9.1)		
		OVCAR-8	10.8 (9.4–12.4)		
		HCT 116	8.6 (7.6–9.9)		
		SF-295	6.9 (5.6–8.4)		
	Paris saponin Pb	HL-60	0.6 (0.4–0.7)		
		OVCAR-8	2.4 (1.9–2.9)		
		HCT 116	1.01 (0.74–1.37)		
	Doxorubicin (control)	HL-60	0.02 (0.01–0.02)		
		OVCAR-8	0.3 (0.2–0.3)		
HCT 116		0.1 (0.1–0.2)			
SF-295		0.2 (0.2–0.3)			
<i>Cestrum laevigatum</i> stems and roots	(25R,S)-5 α -spirostan-2 α ,3 β -diol 3- <i>O</i> - β -D-Glc-(1 \rightarrow 4)- β -D-Gal-(1 \rightarrow 4)- β -D-Gal	HL-60	2.22 (1.55–3.17)	MTT assay	Ribeiro et al. (2016b)
		OVCAR-8	10.80 (9.51–2.27)		
		HCT 116	7.27 (5.93–8.90)		
		SF-295	6.88 (4.49–10.56)		
	(25R,S)-5 α -spirostan-2 α ,3 β -diol 3- <i>O</i> - β -D-Glc-(1 \rightarrow 2)- α -L-Rha-(1 \rightarrow 4)- β -D-Gal	HL-60	7.28 (6.68–7.95)		
		OVCAR-8	15.30 (11.91–19.64)		
		HCT 116	11.41 (9.63–13.51)		
	(25R,S)-5 α -spirostan-2 α ,3 β -diol 3- <i>O</i> - β -D-Gal	SF-295	12.90 (10.78–15.43)		
		HL-60	16.68 (11.85–23.49)		
		OVCAR-8	11.30 (9.21–13.87)		
	Doxorubicin (control)	HCT 116	16.50 (14.3–19.1)		
		SF-295	> 25		
		HL-60	0.02 (0.01–0.02)		
		OVCAR-8	0.26 (0.17–0.3)		
		HCT 116	0.12 (0.09–0.17)		

Table 1 continued

Source	Compound	Cell line	Concentration	Assay	References	
<i>Cestrum parqui</i> leaves	Parquiroside	SF-295	0.24 (0.02–0.27)	CCK-8 assay	Mosad et al. (2017)	
		IC ₅₀ (μM)				
		HeLa	7.7 ± 1.5			
		HepG2	7.2 ± 1.4			
		MCF-7	14.1 ± 4.5			
		U87	3.3 ± 0.63			
		Cisplatin (control)	HeLa			39.2 ± 8.2
			HepG2			14.6 ± 5.9
			MCF-7			7.3 ± 1.3
			U87			23.0 ± 5.6
<i>Chlorophytum deistelianum</i> aerial parts	Chlorodeistelianoside A	IC ₅₀ (μM)		XTT assay	Tabopda et al. (2016)	
		SW480	> 22.12			
		H9c2	> 22.12			
		Chlorodeistelianoside C	SW480			> 24.12
			H9c2			> 24.12
		(25R)-3β-[(β-D-Glc-(1 → 2)-[β-D-Xyl-(1 → 3)]-β-D-Glc-(1 → 4)-β-D-Gal)oxy]-5α-spirostan-12-one	SW480			9.13 ± 0.41
		Solanigroside G	H9c2			8.25 ± 1.16
			SW480			10.07 ± 0.61
		F-gitonin	H9c2			9.57 ± 0.21
			SW480			9.45 ± 0.58
Polianthoside D	H9c2	9.82 ± 0.30				
	SW480	> 24.43				
	H9c2	> 24.43				
	SW480	> 24.43				
<i>Chlorophytum laxum</i> roots	(25R)-26-[[β-D-Glc]oxy]-22α-methoxy-5α-furostan-3β-yl β-D-Glc-(1 → 2)-[β-D-Xyl-(1 → 3)]-β-D-Glc-(1 → 4)-β-D-Gal	SW480	> 24.43	CKK-8 assay	Chu et al. (2018)	
	H9c2	> 24.43				
<i>Chlorophytum orchidastrum</i> roots	25-R-Spirosta-3,5-dien-12β-ol	5-8F	24.8	MTT assay	Acharya et al. (2010)	
	Diosgenin	5-8F	41.9			
<i>Chlorophytum orchidastrum</i> roots	Orchidastroside A	IC ₅₀ (μM)		MTT assay	Acharya et al. (2010)	
		HCT 116	1.6			
		HT-29	1.5			
		Orchidastroside C	HCT 116			1.35
			HT-29			3.6
		Orchidastroside D	HCT 116			2.19
			HT-29			9.15
		Orchidastroside F	HCT 116			2.12
			HT-29			8.87

Table 1 continued

Source	Compound	Cell line	Concentration	Assay	References			
<i>Cordyline fruticosa</i> leaves	Paclitaxel (control)	HCT 116	2.4 (nM)	MTT assay	Fouedjou et al. (2014)			
		HT-29	2.1					
	Fruticoside H		IC ₅₀ (μM)					
		MDA-MB231	69.68					
		A375	37.83					
	Fruticoside I							
		HCT 116	39.8					
		MDA-MB231	50.45					
	Fruticoside J							
		A375	46.59					
		HCT 116	59.97					
	Cisplatin (control)							
MDA-MB231		> 200						
A375		> 200						
HCT 116		> 200						
<i>Costus speciosus</i> tuber	Paclitaxel (control)	MDA-MB231	7.28	MTT assay	Selim and Al Jaouni (2015)			
		A375	0.62					
	Diosgenin	HCT 116	4.97					
			IC ₅₀ (μg ml ⁻¹)					
		HepG2	32.62					
	Paclitaxel (control)	HL-60	22.98					
		MCF-7	11.03					
		HepG2	0.48					
	<i>Cynanchum paniculatum</i> roots	Paclitaxel (control)	HL-60			0.78	SRB assay	Kim et al. (2013)
			MCF-7			0.61		
Cynanside A			IC ₅₀ (μM)					
		A549	> 30					
		SK-OV-3	> 30					
Cynanside B								
		SK-MEL-2	26.55					
		HCT-15	> 30					
		A549	> 30					
		SK-OV-3	> 30					
Doxorubicin (control)								
	SK-MEL-2	17.36						
	HCT-15	> 30						
	A549	0.029						
<i>Datura metel</i> whole plant	Doxorubicin (control)	SK-OV-3	0.036	SRB assay	Mai et al. (2017)			
		SK-MEL-2	0.001					
	Cynanside B	HCT-15	2.041					
			Cell death (%)					

Table 1 continued

Source	Compound	Cell line	Concentration	Assay	References
	3- <i>O</i> -β-D-Xyl-(1 → 2)-α-L-Rha-(1 → 4)[α-L-Rha-(1 → 2)]-β-D-Glc (25R,26R)-spirost-5-en-3β-ol-26-acetamide	HepG2	4 (μg ml ⁻¹)	9.4 (%)	
		MCF-7	20	16.0	
		SK-MEL-2	100	34.9	
	Dioscoroside D	HepG2	4	12.7	
		MCF-7	20	14.1	
		SK-MEL-2	100	28.3	
	Meteloside D	HepG2	4	10.4	
		MCF-7	20	15.5	
		SK-MEL-2	100	30.0	
	Meteloside E	HepG2	4	14.1	
		MCF-7	20	16.1	
		SK-MEL-2	100	25.8	
	Camptothecin (control)	HepG2	4	6.9	
		MCF-7	20	15.5	
		SK-MEL-2	100	35.5	
			4	16.4	
			20	19.9	
			100	31.9	
			4	18.5	
			20	28.6	
			100	38.8	
			4	11.4	
			20	27.1	
			100	44.4	
			4	5.7	
			20	22.5	
			100	44.2	
			4	13.3	
			20	28.6	
		100	41.2		
		4	11.1		
		20	28.4		
		100	48.3		
		4	14.3		
		20	18.7		
		100	41.6		
		4	72.5		
		20	97.0		
		4	76.8		
		20	96.9		
		4	68.1		

Table 1 continued

Source	Compound	Cell line	Concentration	Assay	References	
<i>Digitalis trojana</i> aerial parts			20	MTT assay	Kirmizibezkmez et al. (2014)	
			IC ₅₀ (μM)			84.9
	22- <i>O</i> -methylparvispinoside A	HT29	50.0 ± 0.90			
		MCF-7	50.0 ± 0.15			
	Parvispinoside	HT29	50.0 ± 0.20			
		MCF-7	36.5 ± 0.08			
		PC3	> 50			
	22- <i>O</i> -methylparvispinoside B	HT29	10.0 ± 0.25			
		MCF-7	46.0 ± 0.15			
		PC3	> 50			
<i>Dioscorea bulbifera</i> var. <i>sativa</i> flowers	Staurosporine (control)	HT29	1.2 ± 0.05	MTT assay	Tapondjou et al. (2013)	
		MCF-7	1.0 ± 0.01			
		PC3	1.5 ± 0.05			
			IC ₅₀ (μg ml ⁻¹)			
	Spiroconazol A	ECV-304	5.8			
<i>Dioscorea preussii</i> rhizomes	Pennogenin 3- <i>O</i> -α-L-Rha-(1 → 4)-α-L-Rha-(1 → 4)-[α-L-Rha-(1 → 2)]-β-D-Glc	ECV-304	8.5	MTT assay	Tabopda et al. (2014)	
	26- <i>O</i> -β-D-Glc-(25R)-5-en-furost-3β,17α,22α, 26-tetraol-3- <i>O</i> -α-L-Rha-(1 → 4)-α-L-Rha-(1 → 4)-[α-L-Rha-(1 → 2)]-β-D-Glc	ECV-304	14.3			
			IC ₅₀ (μM)			
	Diospreussinoside B	HCT 116	48.7			
		HT-29	31			
<i>Dioscorea zingiberensis</i> rhizomes	(25R)-17α-hydroxySpirost-5-en-3β-yl <i>O</i> -α-L-Rha-(1 → 4)- <i>O</i> -α-L-Rha-(1 → 4)-β-D-Glc	HCT 116	37.41	MTT assay	Tong et al. (2012)	
		HT-29	42.43			
	(25R)-17α-hydroxySpirost-5-en-3β-yl <i>O</i> -α-L-Rha-(1 → 2)- <i>O</i> -[<i>O</i> -α-L-Rha-(1 → 4)-α-L-Rha-(1 → 4)]-β-D-Glc	HCT 116	2.17			
		HT-29	1.64			
	Paclitaxel (control)	HCT 116	2.65 10 ⁻³			
		HT-29	2.29 10 ⁻³			
			IC ₅₀ (μM)			
Diosgenin	SK-OV-3	> 20				
	B16	> 20				
	LL2	> 20				
	C26	> 20				
	A549	> 20				
	HEK293	> 20				
Trillin	SK-OV-3	> 20				

Table 1 continued

Source	Compound	Cell line	Concentration	Assay	References
		B16	> 20		
		LL2	> 20		
		C26	18.74 ± 1.60		
		A549	> 20		
		HEK293	> 20		
	Diosgenin diglucoside	SK-OV-3	16.71 ± 0.84		
		B16	16.53 ± 0.28		
		LL2	18.02 ± 0.66		
		C26	14.51 ± 0.90		
		A549	18.86 ± 1.24		
		HEK293	> 20		
	Deltonin	SK-OV-3	3.15 ± 0.29		
		B16	4.88 ± 0.43		
		LL2	4.42 ± 0.77		
		C26	1.41 ± 0.51		
		A549	5.65 ± 0.82		
		HEK293	9.73 ± 0.85		
	Zingiberensis saponin	SK-OV-3	1.51 ± 0.53		
		B16	2.64 ± 0.49		
		LL2	2.37 ± 0.54		
		C26	0.81 ± 0.35		
		A549	2.13 ± 0.48		
		HEK293	4.15 ± 0.22		
	Protodeltonin	SK-OV-3	15.86 ± 0.55		
		B16	14.23 ± 1.60		
		LL2	15.58 ± 0.75		
		C26	12.54 ± 0.81		
		A549	14.82 ± 1.28		
		HEK293	> 20		
	Parvifloside	SK-OV-3	16.59 ± 0.72		
		B16	16.12 ± 0.90		
		LL2	14.82 ± 1.60		
		C26	13.83 ± 2.52		
		A549	14.36 ± 1.14		
		HEK293	> 20		
	Dioscin (control)	SK-OV-3	4.14 ± 0.80		
		B16	4.57 ± 0.61		
		LL2	5.03 ± 0.76		
		C26	2.81 ± 1.21		

Table 1 continued

Source	Compound	Cell line	Concentration	Assay	References		
<i>Dracaena cambodiana</i> resin	Doxorubicin (control)	A549	6.82 ± 1.55	MTT assay	Shen et al. (2014)		
		HEK293	6.62 ± 0.28				
		SK-OV-3	0.73 ± 0.35				
		B16	0.77 ± 0.28				
		LL2	0.67 ± 0.12				
		C26	0.50 ± 0.18				
		A549	1.05 ± 0.25				
	Diosgenin 3- <i>O</i> - α -L-Rha-(1 → 2)-[α -L-Rha-(1 → 3)]- β -D-Glc	HEK293	1.32 ± 0.52				
		IC ₅₀ (μ M)					
		K562	1.27				
		BEL-7402	4.72				
		SGC-7901	2.88				
		Pennogenin 3- <i>O</i> - α -L-Rha-(1 → 2)-[α -L-Rha-(1 → 3)]- β -D-Glc	K562			5.09	
			BEL-7402			1.13	
Spirost-5,25(27)-dien-1 β ,3 β -diol 1- <i>O</i> - α -L-Rha-(1 → 2)-[β -D-Xyl-(1 → 3)]- α -L-Ara	SGC-7901	3.39					
	K562	4.77					
	BEL-7402	6.44					
Paclitaxel (control)	SGC-7901	5.61					
	K562	5.98					
	BEL-7402	3.75					
	SGC-7901	1.88					
	<i>Dracaena deisteliiana</i> stem <i>Dracaena arborea</i> bark	Neuroscogenin 1- <i>O</i> - α -L-Rha-(1 → 2)-[β -D-Xyl-(1 → 3)]- α -L-Ara	HT-29	IC ₅₀ (μ M)	MTT assay	Kougan et al. (2010)	
Manioside A		HCT 116	values in the range 7.60–70.73				
		HT-29	1.67				
Spiroconazol A		HCT 116	2.04				
		HT-29	3.21				
Paclitaxel (control)		HCT 116	1.4				
		HT-29					
<i>Dracaena marginata</i> roots		Saponin fraction	HCT 116		Acid phosphatase assay		Ghaly et al. (2014)
			IC ₅₀ (μ g ml ⁻¹)				
		HepG2	13.4				
	MCF7	35					
	Methylprotogracillin	HepG2	29.8				
MCF7		> 50					

Table 1 continued

Source	Compound	Cell line	Concentration	Assay	References	
<i>Dracaena viridiflora</i> leaves	Methylprotodioscin	HepG2	29.8	MTT assay	Teponno et al. (2017)	
		MCF7	> 50			
	Adriamycin (control)	HepG2	6.9			
		MCF7	2.5			
	Trillin		Jurkat			22.36 ± 1.40
			Caco-2			36.49 ± 2.14
			SK-OV-3			64.78 ± 1.91
			A549			14.14 ± 0.10
	Prosapogenin A of dioscin		Jurkat			2.06 ± 0.12
			Caco-2			2.51 ± 0.32
			SK-OV-3			5.69 ± 0.88
			A549			2.11 ± 0.54
	Prosapogenin B of dioscin		Jurkat			21.74 ± 1.80
			Caco-2			13.72 ± 0.84
			SK-OV-3			62.33 ± 1.42
			A549			42.44 ± 1.60
	Dioscin		Jurkat			1.70 ± 0.38
			Caco-2			2.58 ± 0.21
			SK-OV-3			1.90 ± 0.86
A549			0.42 ± 0.15			
Methylprotodioscin		Jurkat	4.82 ± 0.33			
		Caco-2	16.13 ± 0.34			
		SK-OV-3	7.07 ± 0.39			
		A549	5.26 ± 0.29			
Doxorubicin (control)		Jurkat	0.61 ± 0.04			
		Caco-2	2.32 ± 1.04			
		SK-OV-3	0.84 ± 0.08			
		A549	1.15 ± 0.84			
<i>Fritillaria pallidiflora</i> bulbs			IC ₅₀ (μM)	MTT assay	Shen et al. (2012)	
			Pallidifloside D	C6		53.2 ± 3.2
			Polygonatoside B3	HeLa		75.8 ± 4.5
				C6		24.1 ± 1.7
			Polyphyllin V	HeLa		28.1 ± 3.9
				C6		10.3 ± 2.2
			Deltonin	HeLa		9.4 ± 1.1
				C6		5.1 ± 0.2
			Resveratrol (control)	HeLa		5.2 ± 0.9
				C6		24.8 ± 1.8
<i>Lilium longiflorum</i> bulbs	(22R,25R)-spirosol-5-en-3β-yl 3-O-α-L-Rha-(1 → 2)-β-D-Glc-(1 → 4)-β-D-Glc	3T3	IC ₅₀ (μM)	MTT assay	Esposito et al. (2013)	
			8.2			

Table 1 continued

Source	Compound	Cell line	Concentration	Assay	References
	(22R,25R)-spiroisol-5-en-3 β -yl 3-O- α -L-Rha-(1 \rightarrow 2)-[6-O-acetyl- β -D-Glc-(1 \rightarrow 4)]- β -D-Glc	3T3	25.8		
	(25R)-26-O-(β -D-Glc)-furost-5-en-3 β ,22 α ,26-triol 3-O- α -L-Rha-(1 \rightarrow 2)- β -D-Glc-(1 \rightarrow 4)- β -D-Glc	3T3	8.7		
	(25R)-26-O-(β -D-Glc)-furost-5-en-3 β ,22 α ,26-triol 3-O- α -L-Rha-(1 \rightarrow 2)- α -L-Ara-(1 \rightarrow 3)- β -D-Glc	3T3	<1.0		
	(25R)-26-O-(β -D-Glc)-furost-5-en-3 β ,22 α ,26-triol 3-O- α -L-Rha-(1 \rightarrow 2)- α -L-Xyl-(1 \rightarrow 3)- β -D-Glc	3T3	<1.0		
<i>Liriope graminifolia</i> tubers			IC ₅₀ (μ g ml ⁻¹)	MTT assay	Wang et al. (2011)
	Lirigramoside A	SMMC-7721	76.4 \pm 6.6		
		HeLa	26.1 \pm 4.4		
	Lirigramoside B	SMMC-7721	> 100		
		HeLa	18.6 \pm 3.6		
	1-O- β -D-Xyl-3-O- α -L-Rha-(25S)-ruscogenin	SMMC-7721	45.8 \pm 5.4		
		HeLa	13.3 \pm 3.0		
	3-O- α -L-Rha-1-O-sulfo-(25S)-ruscogenin	SMMC-7721	> 100		
		HeLa	40.6 \pm 6.4		
	Cisplatin (control)	SMMC-7721	12.8 \pm 4.8		
		HeLa	5.4 \pm 1.8		
<i>Liriope muscari</i> roots			IC ₅₀ (μ M)	MTT assay	Wu et al. (2017b)
	(25S)-Ruscogenin 1-O- β -D-Glc-(1 \rightarrow 2)-[β -D-Xyl-(1 \rightarrow 3)]- β -D-Glc	MDA-MB-435	15.99 \pm 1.03		
		95D	20.13 \pm 1.18		
		HepG2	49.68 \pm 1.57		
		HeLa	39.98 \pm 1.20		
		MCF-7	47.30 \pm 1.56		
		A549	36.35 \pm 1.39		
	(25R)-Ruscogenin 1-O- β -D-Glc-(1 \rightarrow 2)-[β -D-Xyl-(1 \rightarrow 3)]- β -D-Glc	MDA-MB-435	26.01 \pm 0.85		
		95D	30.00 \pm 0.51		
		HepG2	40.52 \pm 0.96		
		HeLa	33.42 \pm 1.39		
		MCF-7	39.12 \pm 1.02		
		A549	36.01 \pm 1.31		
	(25S)-Ruscogenin 1-O- β -D-Glc-(1 \rightarrow 2)-[β -D-Xyl-(1 \rightarrow 3)]- β -D-Xyl	MDA-MB-435	18.07 \pm 1.34		
		95D	25.67 \pm 0.41		
		HepG2	37.17 \pm 1.71		
		HeLa	21.58 \pm 1.42		
		MCF-7	45.82 \pm 1.44		
		A549	43.53 \pm 1.16		

Table 1 continued

Source	Compound	Cell line	Concentration	Assay	References
	(25R)-Ruscogenin 1- <i>O</i> - β -D-Glc-(1 \rightarrow 2)- [β -D-Xyl-(1 \rightarrow 3)]- β -D-Xyl	MDA-MB-435	17.68 \pm 2.50		
		95D	17.83 \pm 0.37		
		HepG2	29.48 \pm 1.64		
		HeLa	22.23 \pm 1.43		
		MCF-7	42.16 \pm 1.26		
	(25R)-Ruscogenin 1- <i>O</i> - α -L-Rha-(1 \rightarrow 2)- [β -D-Xyl-(1 \rightarrow 3)]- β -D-Glc	MDA-MB-435	19.63 \pm 0.76		
		95D	10.82 \pm 0.18		
		HepG2	15.26 \pm 1.29		
		A549	35.56 \pm 1.46		
	(25S)-Ruscogenin 1- <i>O</i> - β -D-Glc-(1 \rightarrow 2)- [α -L-Ara-(1 \rightarrow 3)]- β -D-Fuc	MDA-MB-435	16.34 \pm 0.60		
		95D	14.34 \pm 0.33		
		HepG2	27.10 \pm 0.84		
		HeLa	14.76 \pm 0.52		
		MCF-7	35.21 \pm 2.02		
	(25R)-Ruscogenin 1- <i>O</i> - β -D-Glc-(1 \rightarrow 2)- [α -L-Ara-(1 \rightarrow 3)]- β -D-Fuc	95D	22.15 \pm 1.41		
		HeLa	42.56 \pm 3.75		
	Neoruscogenin-1- <i>O</i> - β -D-Glc-(1 \rightarrow 2)-[β - D-Xyl-(1 \rightarrow 3)]- β -D-Xyl	MDA-MB-435	24.52 \pm 0.91		
		95D	36.12 \pm 1.08		
		HeLa	24.30 \pm 1.55		
	Neoruscogenin 1- <i>O</i> - α -L-Rha-l-(1 \rightarrow 2)- [β -D-Xyl-(1 \rightarrow 3)]- β -D-Glc	MDA-MB-435	17.54 \pm 1.39		
		95D	11.09 \pm 0.15		
	Neoruscogenin 1- <i>O</i> - β -D-Glc-(1 \rightarrow 2)-[β - D-Xyl-(1 \rightarrow 3)]- β -D-Fuc	MDA-MB-435	9.74 \pm 0.62		
		95D	10.64 \pm 0.21		
		HepG2	15.48 \pm 0.52		
		HeLa	11.02 \pm 0.42		
		MCF-7	10.02 \pm 0.73		
	(25R)-Ruscogenin 1- <i>O</i> - β -D-Glc-(1 \rightarrow 2)- [β -D-Xyl-(1 \rightarrow 3)]- β -D-Fuc	A549	21.25 \pm 1.42		
		MDA-MB-435	4.71 \pm 0.75		
		95D	11.62 \pm 2.00		
		HepG2	Not active		
		HeLa	26.36 \pm 2.01		
	(25S)-Ruscogenin 1- <i>O</i> - β -D-Glc-(1 \rightarrow 2)-[β -D-Xyl-(1 \rightarrow 3)]- β -D-Fuc	MCF-7	NA		
		A549	23.56 \pm 2.64		
		MDA-MB-435	5.91 \pm 0.27		
		95D	11.20 \pm 0.17		
		HepG2	12.76 \pm 0.74		
		HeLa	8.00 \pm 0.45		
		MCF-7	17.88 \pm 0.97		
		A549	8.226 \pm 0.78		

Table 1 continued

Source	Compound	Cell line	Concentration	Assay	References			
<i>Ophiopogon japonicus</i> roots	(25S)-Ruscogenin 1- <i>O</i> - α -L-Rha-(1 \rightarrow 2)-[β -D-Xyl-(1 \rightarrow 3)]- β -D-Glc	MDA-MB-435	9.75 \pm 0.34	MTT assay	Wu et al. (2018)			
		95D	19.58 \pm 0.67					
		HepG2	15.24 \pm 1.53					
		HeLa	14.03 \pm 0.61					
		MCF-7	16.30 \pm 0.73					
		A549	13.99 \pm 0.64					
	5-Fluorouracil (control)	MDA-MB-435	116.8 \pm 13.93					
		95D	83.55 \pm 10.66					
		HepG2	91.9 \pm 16.20					
		HeLa	251.3 \pm 19.93					
		MCF-7	568.3 \pm 54.37					
		A549	244.8 \pm 21.23					
	<i>Ophiopogon japonicus</i> tuberous roots	Pennogenin-3- <i>O</i> - α -L-Rha-(1 \rightarrow 2)-[β -D-Api-(1 \rightarrow 4)]- β -D-Glc	MDA-MB-435			1.90 \pm 0.17	MTT assay	Li et al. (2013)
			HepG2			1.69 \pm 0.18		
A549			4.39 \pm 0.37					
Pennogenin-3- <i>O</i> - α -L-Rha-(1 \rightarrow 2)-[β -D-Xyl-(1 \rightarrow 3)]-[β -D-Glc-(1 \rightarrow 4)]- β -D-Glc		MDA-MB-435	9.13 \pm 1.43					
		HepG2	21.18 \pm 1.87					
		A549	21.27 \pm 2.53					
(25R)-Ruscogenin-1- <i>O</i> - α -L-Rha-(1 \rightarrow 2)-[β -D-Xyl-(1 \rightarrow 3)]- α -L-Ara		MDA-MB-435	10.32 \pm 2.37					
		HepG2	NA (IC ₅₀ > 50 μ M)					
		A549	29.12 \pm 4.66					
5-FU (control)		MDA-MB-435	120 \pm 15.53					
		HepG2	87.3 \pm 12.10					
		A549	256.8 \pm 19.03					
<i>Ophiopogon japonicus</i> tuberous roots		Ophiopogonin Q	HepG2	24 h	2.88			
			HepG2	72 h	1.06			
	HLE		24 h	2.61				
	BEL-7402		24 h	3.59				
	BEL-7403		24 h	6.25				
	HeLa		24 h	2.74				
	Pennogenin 3- <i>O</i> -[2- <i>O</i> -acetyl- α -L-Rha-(1 \rightarrow 2)] [β -D-Xyl-(1 \rightarrow 4)]- β -D-Glc		HepG2	24 h	3.54			
			HepG2	72 h	1.60			
			HLE	24 h	3.63			
			BEL-7402	24 h	3.72			
		BEL-7403	24 h	12.28				

Table 1 continued

Source	Compound	Cell line	Concentration	Assay	References		
<i>Ophiopogon japonicus</i> tubers	Diosgenin 3- <i>O</i> -[2- <i>O</i> -acetyl- α -L-Rha-(1 \rightarrow 2)][β -D-Xyl-(1 \rightarrow 4)]- β -D-Glc	HeLa	24 h	4.26	MTT assay	Wang et al. (2017a)	
		HepG2	24 h	3.30			
		HepG2	72 h	1.49			
		HLE	24 h	1.49			
		BEL-7402	24 h	8.06			
		BEL-7403	24 h	5.13			
		HeLa	24 h	1.47			
		Sprengerinin C	HepG2	24 h			3.07
		HepG2	72 h	1.83			
		HLE	24 h	3.68			
	BEL-7402	24 h	8.13				
	BEL-7403	24 h	1.97				
	Pennogenin 3- <i>O</i> - α -L-Rha-(1 \rightarrow 2)-[β -D-Xyl-(1 \rightarrow 4)]- β -D-Glc	HeLa	24 h	1.74			
	HepG2	24 h	3.04				
	HepG2	72 h	1.71				
	HLE	24 h	3.30				
	BEL-7402	24 h	6.08				
	BEL-7403	24 h	5.14				
	Taxol (control)	HeLa	24 h	3.34			
	HepG2	24 h	33.3				
HepG2	72 h	0.251					
HLE a	24 h	1.95					
BEL-7402	24 h	5.92					
BEL-7403	24 h	11.84					
HeLa	24 h	3.10					
<i>Ophiopogon japonicus</i> fibrous roots	(25R)-Ruscogenin-3-yl α -L-Rha-(1 \rightarrow 2)-[β -D-Xyl-(1 \rightarrow 4)]- β -D-Glc		IC ₅₀ (μ M)		MTT assay	Duan et al. (2010)	
		Ophiopogonin D'	MG-63	3.09			
			SNU-387	3.63			
		Diosgenin 3- <i>O</i> -[2- <i>O</i> -acetyl- α -L-Rha-(1 \rightarrow 2)]- β -D-Xyl-(1 \rightarrow 3)- β -D-Glc	MG-63	1.9			
			SNU-387	0.76			
<i>Ophiopogon japonicus</i> fibrous roots	(25R)-Ruscogenin-3-yl α -L-Rha-(1 \rightarrow 2)-[β -D-Xyl-(1 \rightarrow 4)]- β -D-Glc	Cisplatin (control)	MG-63	11.31	MTT assay	Duan et al. (2010)	
			SNU-387	5.59			
			HeLa	9.14			
			HEp2	11.27			
		Diosgenin-3-yl 2- <i>O</i> -acetyl- α -L-Rha-(1 \rightarrow 2)-[β -D-Xyl-(1 \rightarrow 4)]- β -D-Glc	HeLa	10.77			
	HEp2	10.08					
Pennogenin-3-yl 2- <i>O</i> -acetyl- α -L-Rha-(1 \rightarrow 2)-[β -D-Xyl-(1 \rightarrow 4)]- β -D-Glc	HeLa	13.46					

Table 1 continued

Source	Compound	Cell line	Concentration	Assay	References
<i>Ophiopogon japonicus</i> fibrous roots		HEp2	13.32	MTT assay	Duan et al. (2018)
			IC ₅₀ (μg ml ⁻¹)		
	Fibrophiopogonin A	A375	201.1		
	Fibrophiopogonin B	A375	42.06		
		MCF-7	45.32		
	(25R)-26-[(<i>O</i> -β-D-Glc-(1 → 2)-β-D-Glc)]-22α-hydroxyfurost-5-ene-3- <i>O</i> -[α-L-Rha-(1 → 2)]-β-D-Glc	A375	63.43		
<i>Panicum turgidum</i> aerial parts			IC ₅₀ (μM)	Neutral red uptake assay	Zaki et al. (2017)
	Pennogenin 3β- <i>O</i> -α-L-Rha-(1 → 2)- <i>O</i> -[α-L-Rha-(1 → 4)]- <i>O</i> -α-L-Rha-(1 → 4)]- <i>O</i> -β-D-Glc	SK-MEL	0.47 ± 0.15		
		KB	1.6 ± 0.4		
		BT-549	0.59 ± 0.09		
		SK-OV-3	0.81 ± 0.11		
		VERO	1.5 ± 0.2		
		LLC-PK1	1.005 ± 0.105		
	Yamogenin 3β- <i>O</i> -α-L-Rha-(1 → 2)- <i>O</i> -[α-L-Rha-(1 → 4)]- <i>O</i> -β-D-Glc	SK-MEL	0.76 ± 0.04		
		KB	3.5 ± 1.5		
		BT-549	3.3 ± 1.2		
		SK-OV-3	1.24 ± 0.26		
		VERO	2.8 ± 1.7		
		LLC-PK1	3.15 ± 1.15		
	Yamogenin 3β- <i>O</i> -α-L-Rha-(1 → 2)- <i>O</i> -[α-L-Rha-(1 → 4)]- <i>O</i> -α-L-Rha-(1 → 4)]- <i>O</i> -β-D-Glc	SK-MEL	4.2 ± 1.3		
		KB	8.25 ± 3.25		
		BT-549	4.1 ± 1.9		
		SK-OV-3	3.35 ± 1.15		
		VERO	7.0 ± 3.8		
		LLC-PK1	3.7 ± 1.6		
	Pennogenin 3β- <i>O</i> -α-L-Rha-(1 → 2)- <i>O</i> -[α-L-Rha-(1 → 4)]- <i>O</i> -β-D-Glc	SK-MEL	0.295 ± 0.07		
		KB	1.0 ± 0.1		
		BT-549	1.55 ± 0.15		
SK-OV-3		0.765 ± 0.015			
VERO		0.5 ± 0.05			
	LLC-PK1	0.65 ± 0.05			
Doxorubicin (control)	SK-MEL	3.0 ± 0.78			
	KB	1.7 ± 0.0			
	BT-549	2.9 ± 1.4			
	SK-OV-3	3.3 ± 0.17			
	VERO	> 9			
	LLC-PK1	2.5 ± 0.9			

Table 1 continued

Source	Compound	Cell line	Concentration	Assay	References
<i>Paris polyphylla</i> rhizomes			IC ₅₀ (μM)	XTT assay	Kang et al. (2012)
	Parisyunnanoside G	CCRF-CEM	NA		
	Parisyunnanoside H	CCRF-CEM	NA		
	Parisyunnanoside I	CCRF-CEM	NA		
	Dichotomin	CCRF-CEM	0.59 ± 0.11		
	Pseudoproto-Pb	CCRF-CEM	6.52 ± 0.29		
	Parisyunnanoside A	CCRF-CEM	6.68 ± 0.22		
	Th	CCRF-CEM	5.15 ± 0.16		
	Paris saponin I	CCRF-CEM	1.23 ± 0.08		
	Protogracillin	CCRF-CEM	1.77 ± 0.14		
	Doxorubicin (control)	CCRF-CEM	2.14 ± 0.005		
<i>Paris polyphylla</i> var. <i>yunnanensis</i> rhizomes (Rs), leaves and stems (LSs)			IC ₅₀ (μg ml ⁻¹)	MTT assay	Qin et al. (2018)
	Total saponins Rs	HL-60	1.77		
		A-549	1.75		
		SM MC772	5.23		
		MCF-7	6.62		
		SW480	3.49		
	Total saponins LSs	HL-60	9.54		
		A-549	9.3		
		SM MC772	12.61		
		MCF-7	8.12		
		SW480	11.25		
	Cisplatin (control)	HL-60	0.87		
		A-549	6.48		
		SM MC772	3.77		
		MCF-7	6.4		
		SW480	4.18		
	<i>Paris polyphylla</i> var. <i>yunnanensis</i> rhizomes			IC ₅₀ (μM)	MTT assay
(3β,25R)-spirost-5-en-3-ol-3-O-β-D-Api-(1 → 3)-[α-L-Rha-(1 → 2)]-β-D-Glc		CNE	5.06 ± 1.42		
(3β,17α,25R)-spirost-5-ene-3,17-diol-3-O-β-D-Api-(1 → 3)-[α-L-Rha-(1 → 2)]-β-D-Glc		CNE	3.57 ± 1.05		
(3β,17α,25R)-spirost-5-ene-3,17-diol-3-O-β-D-Glc-(1 → 5)-α-L-Ara-(1 → 4)-[α-L-Rha-(1 → 2)]-β-D-Glc		CNE	9.50 ± 0.80		
(3β,17α,25R)-spirost-5-ene-3,17-diol-3-O-β-D-Xyl-(1 → 5)-α-L-Ara-(1 → 4)-β-D-Glc		CNE	188.55 ± 7.62		
(3β,17α,25R)-spirost-5-ene-3,17-diol-3-O-β-D-Xyl-(1 → 5)-α-L-Ara-(1 → 4)-β-D-Glc		CNE	134.38 ± 2.95		
(3β,25S)-spirost-5-ene-3,27-diol-3-O-α-L-Rha-(1 → 4)-α-L-Rha-(1 → 4)-[α-L-Rha-(1 → 2)]-β-D-Glc		CNE	35.58 ± 2.80		

Table 1 continued

Source	Compound	Cell line	Concentration	Assay	References
	(3 β ,7 β ,25R)-spirost-5-ene-3,7-diol-3- <i>O</i> - β -D-Glc-(1 \rightarrow 3)-[α -L-Rha-(1 \rightarrow 2)]- β -D-Glc	CNE	164.43 \pm 15.0		
	(3 β ,7 α ,25R)-spirost-5-ene-3,7-diol-3- <i>O</i> - α -L-Ara-(1 \rightarrow 4)- β -D-Glc	CNE	1.50 \pm 0.14		
	(3 β ,25R)-spirost-5-en-3-ol-3- <i>O</i> - β -D-Glc-(1 \rightarrow 4)- α -L-Rha-(1 \rightarrow 4)-[α -L-Rha-(1 \rightarrow 2)]- β -D-Glc	CNE	63.98 \pm 4.90		
	(3 β ,17 α ,25R)-spirost-5-ene-3,17-diol-3- <i>O</i> - α -L-Rha-(1 \rightarrow 4)- α -L-Rha-(1 \rightarrow 4)- β -D-Glc	CNE	2.51 \pm 0.42		
	(3 β ,25R)-spirost-5-ene-3-ol-3- <i>O</i> - α -L-Ara-(1 \rightarrow 4)-[α -L-Rha-(1 \rightarrow 2)]- β -D-Glc	CNE	7.28 \pm 1.10		
	(3 β ,25R)-spirost-5-ene-3-ol-3- <i>O</i> - β -D-Glc-(1 \rightarrow 3)-[α -L-Rha-(1 \rightarrow 2)]- β -D-Glc	CNE	95.98 \pm 0.65		
	(3 β ,17 α ,25R)-spirost-5-ene-3,17-diol-3- <i>O</i> - α -L-Ara-(1 \rightarrow 4)- β -D-Glc	CNE	5.92 \pm 0.83		
	(3 β ,17 α ,25R)-spirost-5-ene-3,17-diol-3- <i>O</i> - α -L-Rha-(1 \rightarrow 4)- α -L-Rha-(1 \rightarrow 4)-[α -L-Rha-(1 \rightarrow 2)]- β -D-Glc	CNE	50.46 \pm 2.90		
	(3 β ,25R)-3-hydroxyspirost-5-ene-7-one-3- <i>O</i> - α -L-Ara-(1 \rightarrow 4)-[α -L-Rha-(1 \rightarrow 2)]- β -D-Glc	CNE	23.73 \pm 1.53		
	Cisplatin (control)				
<i>Paris polyphylla</i> var. <i>yunnanensis</i> rhizomes			IC ₅₀ (μ M)	MTT assay	Wu et al. (2017a)
	(23S,24S)-spirost-5,25(27)-diene-1 β ,3 β ,21,23 α ,24 α -pentol-1- <i>O</i> -{ α -L-Rha-(1 \rightarrow 2)-[β -D-Xyl-(1 \rightarrow 3)]- β -D-Glc}-21- <i>O</i> - β -D-Gal-24- <i>O</i> - β -D-Gal	CNE	32.56		
	Parisyunnanoside I	CNE	33.1		
	Cisplatin (control)	CNE	9.35		
<i>Paris polyphylla</i> var. <i>yunnanensis</i> roots			IC ₅₀ (μ M)	MTT assay	Wu et al. (2012a)
	(3 β ,25R)-spirost-5-en-3-ol 3- <i>O</i> - α -L-Rha-(1 \rightarrow 2)- β -D-Glc	CNE	9.2 \pm 0.7		
	(3 β ,25R)-spirost-5-en-3-ol 3- <i>O</i> - β -D-Glc-(1 \rightarrow 6)-[α -L-Rha-(1 \rightarrow 2)]- β -D-Glc	CNE	52.9 \pm 3.7		
	(3 β ,25R)-spirost-5-en-3-ol 3- <i>O</i> - α -L-Rha-(1 \rightarrow 4)- α -L-Rha-(1 \rightarrow 4)-[α -L-Rha-(1 \rightarrow 2)]- β -D-Glc	CNE	4.7 \pm 1.1		
	(3 β ,17 α ,25R)-spirost-5-ene-3,17-diol 3- <i>O</i> - α -L-Rha-(1 \rightarrow 2)- β -D-Glc	CNE	11.1 \pm 4.7		
	(3 β ,17 α ,25R)-spirost-5-ene-3,17-diol 3- <i>O</i> - α -L-Ara-(1 \rightarrow 4)-[α -L-Rha-(1 \rightarrow 2)]- β -D-Glc	CNE	2.7 \pm 1.1		
	Cisplatin (control)	CNE	23.7 \pm 1.5		

Table 1 continued

Source	Compound	Cell line	Concentration	Assay	References
<i>Paris polyphylla</i> var. <i>yunnanensis</i> rhizomes			IC ₅₀ (μM)	MTT assay	Wen et al. (2015)
	Parisyunnanoside H	HEK293	0.9		
		HepG2	5.6		
	Paris saponin I	HEK293	1.8		
		HepG2	1.8		
	Trigofenoside A	HEK293	3.4		
		HepG2	5.6		
	Dichotomin	HEK293	0.58		
		HepG2	0.9		
	Parisyunnanoside B	HEK293	2.5		
	HepG2	1.2			
Pseudoproto-Pb	HEK293	1.8			
	HepG2	1.8			
(–)-OddC (control)	HEK293	0.3			
	HepG2	0.17			
<i>Paris polyphylla</i> var. <i>yunnanensis</i> leaves			IC ₅₀ (μM)	MTT assay	Qin et al. (2016)
	Nuatigenin 3- <i>O</i> -α-L-Rha-(1 → 2)-β-D-Glc	HepG2	2.9 ± 0.5		
		HEK293	5.0 ± 0.6		
	Abutiloside L	HepG2	7.0 ± 0.8		
		HEK293	12.9 ± 2.7		
Troxacitabine (control)	HepG2	0.17 ± 0.02			
	HEK293	0.30 ± 0.03			
<i>Paris quadrifolia</i> rhizomes			IC ₅₀ (μg ml ⁻¹)	MTT assay	Stefanowicz-Hajduk et al. (2011)
	Saponin-rich fractions:	HL-60	13 ± 1.3		
	Solid residue	HeLa	10 ± 0.5		
		MDA-MB-468	27 ± 1.3		
		fibroblasts	28 ± 1.4		
	Butanolic fraction	HL-60	15 ± 2		
		HeLa	24 ± 1.2		
		MDA-MB-468	60 ± 5		
		fibroblasts	60 ± 6		
				IC ₅₀ (μg ml ⁻¹)	MTT assay
<i>Paris quadrifolia</i> rhizomes	Pennogenin 3- <i>O</i> -α-L-Rha-(1 → 4)-β-D-Glc	HL-60	47 ± 2.8		
	Pennogenin 3- <i>O</i> -α-L-Rha-(1 → 2)-β-D-Glc	HL-60	16 ± 0.8		
		HeLa	18 ± 0.9		
		MCF-7	25 ± 1.5		
	Pennogenin 3- <i>O</i> -α-L-Rha-(1 → 4)-[α-L-Rha-(1 → 2)]-β-D-Glc	HL-60	1.0 ± 0.04		

Table 1 continued

Source	Compound	Cell line	Concentration	Assay	References			
<i>Paris quadrifolia</i> rhizomes	Pennogenin 3- <i>O</i> - α -L-Rha-(1 \rightarrow 4)- α -L-Rha-(1 \rightarrow 4)-[α -L-Rha-(1 \rightarrow 2)]- β -D-Glc	HeLa	1.8 \pm 0.072	MTT assay	Stefanowicz-Hajduk et al. (2015)			
		MCF-7	2.4 \pm 0.096					
		HL-60	2.0 \pm 0.08					
	Etoposide (control)	HeLa	2.5 \pm 0.125					
		MCF-7	3.2 \pm 0.128					
		HL-60	0.45 \pm 0.022					
	Mitoxantrone (control)	HeLa	> 50					
		MCF-7	> 50					
		HL-60	0.06 \pm 0.004					
	<i>Paris tibetica</i> rhizomes	PS-1	HeLa			0.93 \pm 0.15	IC ₅₀ (μ mol l ⁻¹)	MTT assay
			HaCaT			0.82 \pm 0.13		
			HeLa			0.55 \pm 0.01		
HaCaT			0.58 \pm 0.04					
<i>Paris vietnamensis</i> rhizomes	PARIS saponin II	BEL-7402	0.48	IC ₅₀ (μ M)	CCK-8 assay			
		U251	2.16 \pm 0.65					
<i>Sansevieria trifasciata</i> aerial parts	25(R)-Diosgenin-3- <i>O</i> - α -L-Rha-(1 \rightarrow 2)- α -L-Rha-(1 \rightarrow 3)- β -D-Glc	U87MG	2.33 \pm 1.03	MTS assay	Teponno et al. (2016)			
		U251	3.14 \pm 1.26					
		U87MG	2.97 \pm 0.94					
	ACNU (control)	U251	0.96 \pm 0.05					
		U87MG	0.88 \pm 0.04					
	<i>Sansevieria cylindrica</i> aerial parts	Trifasciatoside B	HeLa			47.1	IC ₅₀ (μ g ml ⁻¹)	MTT assay
			HeLa			40.7		
HeLa			26.5					
HeLa			26.5					
HeLa			26.5					
(25S)-Ruscogenin-1- <i>O</i> - α -L-Rha-(1 \rightarrow 2)- β -D-Glc	MCF-7	24 \pm 1	HCT 116	23 \pm 1				
	HepG2	21 \pm 1						
	HepG2	21 \pm 1						

Table 1 continued

Source	Compound	Cell line	Concentration	Assay	References			
<i>Sansevieria cylindrica</i> aerial parts	(25S)-Ruscogenin-3- <i>O</i> - α -L-Rha-(1 \rightarrow 4)- β -D-Glc	MCF-7	12 \pm 1	SRB assay	Said et al. (2015)			
		HCT 116	11 \pm 2					
		HepG2	13 \pm 1					
	(25S)-Ruscogenin-3- <i>O</i> - β -D-Glc	MCF-7	> 50					
		HCT 116	> 50					
		HepG2	> 50					
	(25S)-Ruscogenin-1- <i>O</i> - α -L-Rha-(1 \rightarrow 2)- $[\beta$ -D-Xyl-(1 \rightarrow 3)]- α -L-Ara	MCF-7	7 \pm 2					
		HCT 116	4 \pm 2					
		HepG2	9 \pm 2					
	(25R)-26- <i>O</i> - β -D-Glc-furost-5-ene-1 β ,3 β ,22 α ,26-tetrol-1- <i>O</i> - α -L-Rha-(1 \rightarrow 2)- $[\beta$ -D-Xyl-(1 \rightarrow 3)]- α -L-Ara	MCF-7	25 \pm 1					
		HCT 116	19 \pm 1					
		HepG2	21 \pm 1					
	Doxorubicin hydrochloride (control)	MCF-7	13 \pm 1					
		HCT 116	2 \pm 3					
		HepG2	1 \pm 1					
1 β -Hydroxy-kryptogenin-1- <i>O</i> - α -L-Rha-(1 \rightarrow 2)- α -L-Ara	MCF-7	153	IC ₅₀ (μ M)	MTT assay	Sun et al. (2016)			
	PC-3	175						
	HCT 116	90						
Alliospiroside A	MCF-7	69						
	PC-3	99						
	HCT 116	10						
Doxorubicin (control)	MCF-7	6						
	PC-3	4						
<i>Schizocapsa plantaginea</i> tubers	Taccaoside	SMMC-7721				24 h	2.55	
						48 h	1.72	
		BEL-7404				24 h	8.10	
						48 h	5.94	
<i>Smilacina japonica</i> rhizomes and roots	Japonicoside A	SMMC-7221				1.19 \pm 0.03	MTT assay	Liu et al. (2012c)
		DLD-1				1.66 \pm 0.08		
		SMMC-7221	5.40 \pm 0.11					
	Japonicoside B	DLD-1	1.21 \pm 0.05					
		SMMC-7221	3.14 \pm 0.11					
	Japonicoside C	DLD-1	2.16 \pm 0.09					
		SMMC-7221	3.14 \pm 0.11					
	Taxol (control)	SMMC-7221	3.14 \pm 0.11					

Table 1 continued

Source	Compound	Cell line	Concentration	Assay	References
<i>Smilax glauco-china</i> tubers		DLD-1	2.16 ± 0.09	MTT assay	Liu et al. (2017b)
			IC ₅₀ (μM)		
	Glauco-chinaoside A	SGC-7901	2.7		
	Glauco-chinaoside B	SGC-7901	11.5		
	Glauco-chinaoside E	SGC-7901	6.8		
	Cisplatin (control)	SGC-7901	Not specified		
<i>Smilax korthalsii</i> leaves			IC ₅₀ (μM)	MTT assay	Hamid et al. (2016)
	Diosgenin	K562	6.25		
		WRL	14.34		
		MCF-7	38		
		COLO	12.4		
	Tamoxifen (control)	K562	7.26		
		WRL	12.25		
		MCF-7	8.54		
		COLO	10.08		
<i>Smilax ornata</i> roots and rhizomes			Inhibition of cell proliferation (μg ml ⁻¹)	SRB assay	Challinor et al. (2012)
	Sarsaparilloside B	NFF	> 50		
		HeLa	> 50		
		HT29	> 50		
		MCF-7	> 50		
		MM96L	> 50		
		K562	> 50		
	Sarsaparilloside C	NFF	27		
		HeLa	42		
		HT29	4.8		
		MCF-7	24		
		MM96L	23		
		K562	28		
	Sarsaparilloside	NFF	13		
		HeLa	12		
		HT29	5		
		MCF-7	9.5		
		MM96L	14		
		K562	22		
		Δ20(22)-sarsaparilloside	NFF		
		HeLa	40		

Table 1 continued

Source	Compound	Cell line	Concentration	Assay	References
		HT29	14		
		MCF-7	3.4		
		MM96L	3.8		
		K562	4.3		
	Parillin	NFF	> 50		
		HeLa	> 50		
		HT29	> 50		
		MCF-7	> 50		
		MM96L	> 50		
		K562	> 50		
<i>Smilax scobinicaulis</i> rhizomes and roots			IC ₅₀ (μM)	MTT assay	Zhang et al. (2013)
	(25 R)-5α-spirostan-3β, 6β-diol 3-O-β-D-Glc-(1 → 4)-[α-L-Ara-(1 → 6)]-β-D-Glc	A549	3.7		
		LAC	5.7		
		HeLa	3.64		
	Doxorubicin (control)	A549	1.08		
		LAC	0.95		
		HeLa	1.16		
<i>Smilax scobinicaulis</i> rhizomes			IC ₅₀ (μM)	MTT assay	Shu et al. (2017)
	Smilscobinoside D	HCT 116	10.5		
		SGC-7901	21.4		
	Smilscobinoside D	HCT 116	7.8		
		SGC-7901	15.8		
<i>Smilax trinervula</i> rhizomes and roots			IC ₅₀ (μM)	MTT assay	Liang et al. (2016)
	Trinervuloside B	SGC-7901	8.1		
		HCT-116	5.5		
<i>Solanum glabratum</i> var. <i>sepicula</i> aerial parts			IC ₅₀ (μM)	MTT assay	Abdel-Sattar et al. (2015)
	23-β-D-Glc-(23S, 25R)-spirost-5-en-3, 23 diol 3-O-α-L-Rha-(1 → 2)-O-[α-L-Rha-(1 → 4)]-β-D-Glc	PC3	> 32		
		HT29	> 2		
	(25R)-spirost-5-en-3-ol 3-O-α-L-Rha-(1 → 2)-O-[β-D-Glc-(1 → 3)]-β-D-Gal	PC3	14		
		HT29	16.7		
	(23S,25R)-spirost-5-en-3, 23 diol 3-O-α-L-Rha-(1 → 2)-O-[α-L-Rha-(1 → 4)]-β-D-Glc	PC3	> 32		
		HT29	> 32		
	Digitonin (positive control)	PC3	1.8		
		HT29	3		

Table 1 continued

Source	Compound	Cell line	Concentration	Assay	References
<i>Solanum incanum</i> roots/ <i>S. heteracanthum</i> roots	(23S,25R)-spirost-5-en-3 β ,23-diol 3- <i>O</i> - { β -D-Xyl-(1 \rightarrow 2)- <i>O</i> - α -L-Rha-(1 \rightarrow 4)- [<i>O</i> - α -L-Rha-(1 \rightarrow 2)]- β -D-Glc}		IC ₅₀ (μ M)	MTT assay (with HCT 116 and HT-29)	Manase et al. (2012)
		HCT 116	62.42 \pm 0.66	XTT assay (with SW480, DU145 and EMT6)	
		Protodioscin	HT-29		72.24 \pm 20.62
			SW480		> 29.53
			DU145		> 29.53
			EMT6		> 29.53
		Methyl-protodioscin	HCT 116		2.26 \pm 2.29
			HT-29		3.48 \pm 3.01
			SW480		6.68
			DU145		> 28.63
		Indioside D	HCT 116		2.76 \pm 1.93
			HT-29		3.30 \pm 3.00
			SW480		> 28.25
			DU145		> 28.25
		Paclitaxel (control)	HCT 116		3.87 \pm 2.51
			HT-29		5.28 \pm 0.51
			SW480		20.68
			DU145		> 28.20
			EMT6		24.44
Etoposide (control)	HCT 116	2.65 10^{-3}			
	HT-29	2.29 10^{-3}			
	SW480	13.22 \pm 3.79			
	DU145	41.26 \pm 17.57			
	EMT6	> 200			
<i>Solanum procumbens</i> whole plant			IC ₅₀ (μ M)	MTT assay	Hien et al. (2018)
	Solaprocumoside A	HepG2	55.7 \pm 1.5		
	Solaprocumoside B	HepG2	48.1 \pm 2.2		
	Paniculonin B	HepG2	78.3 \pm 2.4		
	Elipticine (control)	HepG2	1.43 \pm 0.17		
<i>Solanum surattense</i> aerial parts			IC ₅₀ (μ M)	MTT assay	Lu et al. (2011)
	(22R,25R)-16 β -H-22 α -N-spirosol-3 β -ol- 5-ene 3- <i>O</i> - α -L-Rha-(1 \rightarrow 2)-[α -L-Rha- (1 \rightarrow 4)]- β -D-Glc	A549	20.3 \pm 1.1		
		MGC-803	45.6 \pm 1.5		
		HepG2	26.1 \pm 0.6		
	(22R,23S,25R)-3 β ,6 α ,23-trihydroxy-5 α - spirostane 6- <i>O</i> - β -D-Xyl-(1 \rightarrow 3)- β -D- Qui	A549	71.2 \pm 2.0		

Table 1 continued

Source	Compound	Cell line	Concentration	Assay	References	
<i>Solanum violaceum</i> whole plant	(22R,23S,25S)-3 β ,6 α ,23-trihydroxy-5 α -spirostane 6-O- β -D-Xyl-(1 \rightarrow 3)-O- β -D-Qui	MGC-803	NA	MTT assay	Yen et al. (2012)	
		HepG2	NA			
		A549	NA			
	(22R,23R,25S)-3 β ,6 α ,23-trihydroxy-5 α -spirostane 6-O- β -D-Xyl-(1 \rightarrow 3)-O- β -D-Qui	MGC-803	63.2 \pm 0.8			
		HepG2	NA			
		A549	62.5 \pm 1.6			
	Khasianine	MGC-803	NA			
		HepG2	88.8 \pm 1.2			
		A549	26.7 \pm 1.5			
	Solamargine	MGC-803	35.4 \pm 0.7			
		HepG2	45.3 \pm 2.1			
		A549	15.7 \pm 0.6			
	Cisplatin (control)	MGC-803	NA			
		HepG2	23.2 \pm 0.8			
		A549	7.6 \pm 1.6			
	<i>Solanum violaceum</i> whole plant	Indioside H	MGC-803			3.5 \pm 0.3
			HepG2			8.7 \pm 0.4
			A549			IC ₅₀ (μ g ml ⁻¹)
		Indioside I	HepG2			2.22 \pm 0.01
			Hep3B			2.95 \pm 0.02
			MCF-7			4.78 \pm 0.02
			A549			3.09 \pm 0.02
			Ca9-22			2.95 \pm 0.07
			MDA-MB-231			6.12 \pm 0.15
		Borassoside D	HepG2			5.33 \pm 0.16
			Hep3B			3.32 \pm 0.42
			MCF-7			11.57 \pm 0.70
A549			7.27 \pm 0.07			
Ca9-22			6.76 \pm 0.15			
MDA-MB-231			8.04 \pm 0.12			
Yamogenin 3-O- α -L-Rha-(1 \rightarrow 2)- β -D-Glc		HepG2	> 20			
		Hep3B	> 20			
		MCF-7	> 20			
		A549	> 20			
		Ca9-22	> 20			
		MDA-MB-231	> 20			
		HepG2	6.48 \pm 0.01			
		Hep3B	6.98 \pm 0.05			
		MCF-7	5.84 \pm 0.04			
		A549	4.26 \pm 0.02			

Table 1 continued

Source	Compound	Cell line	Concentration	Assay	References
		Ca9-22	4.51 ± 0.24		
		MDA-MB-231	7.25 ± 0.15		
	Borassoside E	HepG2	1.83 ± 0.12		
		Hep3B	2.03 ± 0.03		
		MCF-7	2.61 ± 0.10		
		A549	2.34 ± 0.02		
		Ca9-22	2.33 ± 0.02		
		MDA-MB-231	2.75 ± 0.10		
	3- <i>O</i> -chacotriosyl-(25S)-spirost-5-en-3β-ol	HepG2	6.44 ± 0.45		
		Hep3B	2.87 ± 0.04		
		MCF-7	8.84 ± 0.12		
		A549	4.09 ± 0.08		
		Ca9-22	3.77 ± 0.02		
		MDA-MB-231	5.84 ± 0.06		
	Doxorubicin (control)	HepG2	0.18 ± 0.00		
		Hep3B	1.31 ± 0.12		
		MCF-7	0.80 ± 0.03		
		A549	1.40 ± 0.02		
		Ca9-22	0.31 ± 0.01		
		MDA-MB-231	1.39 ± 0.00		
<i>Tacca integrifolia</i> rhizomes			IC ₅₀ (μM)	MTT assay	Shwe et al. (2010)
	(3β,25R)-spirost-5-en-3-yl 6-deoxy-α-L-Man-(1 → 2)-[β-D-Glc-(1 → 4)-6-deoxy-α-L-Man-(1 → 3)]-β-D-Glc	HeLa	72 h	3.0 ± 0.5	
	(3β,25R)-spirost-5-en-3-yl 6-deoxy-α-L-Man-(1 → 2)-[6-deoxy-α-L-Man-(1 → 3)]-β-D-Glc	HeLa	72 h	1.2 ± 0.4	
	(3β,22R,25R)-26-(β-D-Glc)-22-hydroxyfurost-5-en-3-yl 6-deoxy-α-L-Man-(1 → 2)-[6-deoxy-α-L-Man-(1 → 3)]-β-D-Glc	HeLa	72 h	1.5 ± 0.3	
	(3β,22R,25R)-26-(β-D-Glc)-22-methoxyfurost-5-en-3-yl 6-deoxy-α-L-Man-(1 → 2)-[6-deoxy-α-L-Man-(1 → 3)]-β-D-Glc	HeLa	72 h	3.5 ± 0.5	
	(3β,22R,25R)-26-(β-D-Glc)-22-hydroxyfurost-5-en-3-yl 6-deoxy-α-L-Man-(1 → 2)-[β-D-Glc-(1 → 4)-6-deoxy-α-L-Man-(1 → 3)]-β-D-Glc	HeLa	72 h	4.0 ± 0.6	
	Podophyllotoxin (control)	HeLa	72 h	0.1 ± 0.02	
<i>Trillium kamschaticum</i> whole plant			IC ₅₀ (μM)	MTT assay	Qin et al. (2017)
	Trillikamtoside L	HCT 116	17.28 ± 2.69		

Table 1 continued

Source	Compound	Cell line	Concentration	Assay	References		
<i>Trillium tschonoskii</i> rhizomes	Trillikamtoside P	HCT 116	4.92 ± 1.00	Trypan blue dye exclusion assay	Huang and Zou (2015)		
	Trillikamtoside Q	HCT 116	22.48 ± 8.68				
	Trillikamtoside R	HCT 116	5.84 ± 1.05				
	Camptothecin (control)	HCT 116	0.0115 ± 0.0009				
<i>Tupistra chinensis</i> rhizomes	Pennogenin 3- <i>O</i> - α -L-Rha-(1 → 2)[α -L-Rha-(1 → 4)]- β -D-Glc (TTB2)	Rh1	48 h	7.5	MTT assay	Pan et al. (2012)	
			IC ₅₀ (μ M)				
	Tupichinin A	HL-60	18.58				
		SMMC-7721	> 40				
		A549	19.99				
		MCF-7	11.01				
		SW480	10.78				
	3-Epi-neuroscogenin 3- β -D-Glc	HL-60	10.02				
		SMMC-7721	12.76				
		A549	11.4				
MCF-7		5.02					
SW480		28.26					
Cisplatin (control)	HL-60	2.03					
	SMMC-7721	13.54					
	A549	12.56					
	MCF-7	18.65					
	SW480	19.7					
<i>Tupistra chinensis</i> rhizomes			IC ₅₀ (μ M)	MTT assay	Liu et al. (2012b)		
<i>Tupistra chinensis</i> rhizomes	Tupisteroide C	A549	25.9	MTT assay	Liu et al. (2012a)		
	Mitomycin C	A549	not specified				
<i>Tupistra chinensis</i> roots and rhizomes	(25R)-26- <i>O</i> - β -D-Glc-furost-1 β ,3 β ,22 α ,26-tetraol 3- <i>O</i> - β -D-Glc	A549	6.6	MTT assay	Li et al. (2015)		
	(25R)-26- <i>O</i> - β -D-Glc-furost-5-en-1 β ,3 α ,22 α ,26-tetraol 3- <i>O</i> - β -D-Glc	A549	6.7				
	(25R)-26- <i>O</i> - β -D-Glc-furost-1 β ,3 β ,5 β ,22 α ,26-pentaol-3- <i>O</i> - β -D-Glc	A549	29.1				
			IC ₅₀ (μ M l ⁻¹)				
(20S,22R)-Spirost-25 (27)-en-1 β ,3 β ,5 β -trihydroxy-1- <i>O</i> - β -D-Xyl	A549	86.63 ± 2.33					
	NCI-H1299	88.21 ± 1.34					

Table 1 continued

Source	Compound	Cell line	Concentration	Assay	References
<i>Vernonia amygdalina</i> leaves	5-FU (control)	A549	38.65 ± 1.59	MTT assay	Wang et al. (2018)
		NCI-H1299	42.78 ± 1.63		
			Concentration not specified		
			(%) Inhibition		
	Vernoniamyoside A	BT-549	63.61		
		MDA-MB-231	28.97		
		MCF-7	46.54		
	Vernoniamyoside B	HeLa	42.05		
		BT-549	62.17		
		MDA-MB-231	27.78		
		MCF-7	37.07		
	Vernoniamyoside C	HeLa	31.64		
		BT-549	34.18		
		MDA-MB-231	32.74		
		MCF-7	39.38		
	Vernoniamyoside D	HeLa	26.73		
		BT-549	44.00		
		MDA-MB-231	31.53		
		MCF-7	31.36		
	Vernonioside B ₂	HeLa	32.93		
	BT-549	36.41			
	MDA-MB-231	33.61			
	MCF-7	49.72			
Vernoamyoside D	HeLa	21.48			
	BT-549	51.14			
	MDA-MB-231	30.75			
	MCF-7	39.08			
Doxorubicin (control)	HeLa	35.63			
	BT-549	83.79			
	MDA-MB-231	83.39			
	MCF-7	95.32			
	HeLa	92.70			
<i>Ypsilandra thibetica</i> whole plant			IC ₅₀ (μM)	MTT assay	Lu et al. (2010)
	Ypsilandroside H	A549	> 40		
	Ypsilandroside I	HL-60	Not specified		
	Ypsilandroside J	PANC-1			
	Ypsilandroside K	SMMC-7721			
	Ypsilandroside L	SK-BR-3			
	Polyphyllouside III				
	Cisplatin (control)				

Table 1 continued

Source	Compound	Cell line	Concentration	Assay	References
	Polyphyllin I	A549	0.625 ($\mu\text{g ml}^{-1}$)	MTT assay	Kong et al. (2010)
		NCI-H460	24 h		
		SK-MES-1	48 h		
			72 h		
			1.25 ($\mu\text{g ml}^{-1}$)		
			24 h		
			48 h		
			72 h		
			2.5 ($\mu\text{g ml}^{-1}$)		
			24 h		
			48 h		
			72 h		
			5 ($\mu\text{g ml}^{-1}$)		
			24 h		
			48 h		
			72 h		
			10 ($\mu\text{g ml}^{-1}$)		
			24 h		
			48 h		
			72 h		
			0.625 ($\mu\text{g ml}^{-1}$)		
			24 h		
			48 h		
			72 h		
			1.25 ($\mu\text{g ml}^{-1}$)		
			24 h		
			48 h		
			72 h		
			2.5 ($\mu\text{g ml}^{-1}$)		
			24 h		
			48 h		
			72 h		
			5 ($\mu\text{g ml}^{-1}$)		
			24 h		
			48 h		
			72 h		
			10 ($\mu\text{g ml}^{-1}$)		

Table 1 continued

Source	Compound	Cell line	Concentration	Assay	References
			24 h		
			48 h		
			72 h		
			0.625 ($\mu\text{g ml}^{-1}$)		
			24 h		
			48 h		
			72 h		
			1.25 ($\mu\text{g ml}^{-1}$)		
			24 h		
			48 h		
			72 h		
			2.5 ($\mu\text{g ml}^{-1}$)		
			24 h		
			48 h		
			72 h		
			5 ($\mu\text{g ml}^{-1}$)		
			24 h		
			48 h		
			72 h		
			10 ($\mu\text{g ml}^{-1}$)		
			24 h		
			48 h		
			72 h		
			IC ₅₀ ($\mu\text{g ml}^{-1}$)		
				CCK-8 assay	Yu et al. (2018b)
	Polyphyllin I	HCT-116	72 h		

Human cancer cell lines: breast: BT-549, MCF-7, MDA-MB-231, MDA-MB-435, MDA-MB-468, SK-BR-3; cervix: Caski, HeLa, KB; colon: CaCo-2, COLO, DLD-1, HCT 116, HCT-15, HT-29, LOVO SW480, SW620, W480; esophagus: KYSE 510; gingival: Ca9-22; glioblastoma: SF-268, SF-295, U251, U87MG; leukemia: CCRF-CEM, HL-60, Jurkat, K562; larynx: Hep2; liver: BEL-7402, BEL-7403, BEL-7404, HLE, Hep3B, HepG2, SMMC-7721, SMMC-7221, SNU-387, WRL; lung: 95D, A549, LAC, NCI-H1299, NCI-H460, SK-MES-1; melanoma: A375, MM96L, SK-MEL, SK-MEL-2; neuroblastoma: IMR-32, LA-N-2, NB-69; ovary: OVCAR-8, SK-OV-3; pancreas: PANC-1; pharynx: 5-8F, CNE; prostate: DU145, PC-3; sarcoma: MG-63, Rh1; stomach: BGC-823, MGC-803, SGC-7901; urinary bladder: ECV-304

Animal cancer cell lines: breast: EMT6; glioblastoma: C6; lung: LL2; colon: C26; melanoma: B16; sarcoma: WEHI-164, J-774

Human normal cell lines: fibroblasts: HFF, NFF, Hs68; keratinocytes: HaCaT; kidney embryonic: HEK293; lung epithelial: MRS-5; vein endothelial: EA.hy926, HUVEC

Animal normal cell lines: cardiomyoblasts: H9c2; epidermal: JB6 P⁺Cl-41; fibroblasts: 3T3; kidney epithelial: LLC-PK1; kidney fibroblasts: VERO

NA not active

Structure–activity correlation

Despite a vast number of papers that cite the results of cytotoxic activity of steroidal saponins only a relatively small number include some reference to

potential structure–activity relationships. These are usually not fully conclusive statements resulting from the observations made on a very limited number of compounds. In the time-span covered by this review, only a few studies have been specially designed to




Fig. 1 The number of tested substances and number of reports published in the time scope covered by this review (2010–2018)




Fig. 2 The share of experiments on specific types of tumors and normal cell lines in the total number of tests performed on human cell lines

explore structure–activity correlations. These include the one by Pérez-Labrada et al. (2012a, b) who, for the purpose of their study, had synthesized twelve spirostanol glycosides differing mainly in C-ring functional groups, which influenced the lipophilicity and conformational flexibility of compounds (Pérez-Labrada et al. 2012a). These included methylene-, methoxyl-, α,β -unsaturated ketone and lactone. Two

glycosylation pathways led to a series of 3,6-dipivaloylated β -D-glucosides (pivaloyl = 2,2-dimethylpropanoyl) and a series of β -chacotriosides (α -L-Rha-(1 \rightarrow 2)-[α -L-Rha-(1 \rightarrow 4)]- β -D-Glc). The obtained compounds were analysed with respect to their cytotoxicity against the human myeloid leukemia cell line (HL-60) and benign blood cells. The results indicate that among the two glycosidic series, the one

based on a β -chacotrioside moiety was more potent. This activity was however greatly correlated with the rigidity of the aglycone and its hydrophobic character. From among all tested saponins, chacotriosides either with a methylene group at C-12 or no substitution in C-ring showed the highest cytotoxic potential against malignant cell line. However, their selectivity as compared to 3,6-divaloylated spirostanyl glucosides was much lower.

In a subsequent study by the same research group on a larger variety of synthetic spirostanol glycosides, the partially pivaloylated β -D-glucosides of 5α -hydroxy-laxogenin were the most potent (Pérez-Labrada et al. 2012b). Comparison of the results obtained for different β -chacotriosides, has again confirmed that vast differences can be seen with a change in the aglycone part. Hecogenin derivative was highly cytotoxic against the tested HL-60 cell line (IC_{50} 4.3 ± 1.0 μ M) whereas 5α -hydroxy-laxogenin β -chacotrioside showed a complete loss of activity ($IC_{50} > 100$ μ M).

Other studies in which any references to possible structure–activity relationships were made, generally indicate that both structural features of steroidal saponins, that is the nature of the aglycone and the sugar moiety, together determine their cytotoxicity.

Thirteen saponins isolated from the roots of *Liriopis muscari* were analysed in this respect against a fairly wide panel of cancer cell lines (MDA-MB-435, 95D, HepG2, HeLa, MCF-7 and A549) (Wu et al. 2017b). The authors were able to distinguish three groups based on the structural features of the aglycone, namely the (25S)-ruscogenin, (25R)-ruscogenin, and neoruscogenin groups. This allowed to compare the potential contribution to the cytotoxic activity of the specific configuration at C-25, either 25R, 25S or 25,27-double bond. The obtained cytotoxicity results have shown that the impact of this structural feature is related to the nature of the sugar chain. In all saponins bearing β -D-Glc-(1 \rightarrow 2)-[β -D-Xyl-(1 \rightarrow 3)]- β -D-Xyl or β -D-Glc-(1 \rightarrow 2)-[β -D-Xyl-(1 \rightarrow 3)]- β -D-Glc sugar chains at C-1, the configuration at C-25 was of no consequence in all tested cell lines. Interestingly, a different sugar chain composed of β -D-Glc-(1 \rightarrow 2)-[β -D-Ara-(1 \rightarrow 3)]- β -D-Fuc, together with 25R configuration seemed to have a detrimental effect on the cytotoxicity, which was observed against all the tested cell lines. Similar regularity was seen when comparing

compounds with yet another sugar chains, however not in case of the whole spectrum of tested cell lines.

In another study on ten saponins from *Asparagus filicinus* similar results with respect to C-25 configuration were obtained, suggesting that 25S spirostanol aglycone may be a more important structural feature (Wu et al. 2010). Another conclusion drawn from these studies refers to the sugar moiety, clearly indicating that its presence at C-23 significantly reduces the cytotoxic potential of these compounds.

Beit-Yannai et al. (2011) in their study on saponins from *Balanites aegyptiaca* have seen a pronounced difference in cytotoxicity against MCF-7 human breast cancer and HT-29 human colon cancer cells between two compounds differing in only one terminal sugar (dioscin vs SAP-884—diosgenin 3-O- β -D-Glc-(1 \rightarrow 4)-[α -L-Rha-(1 \rightarrow 2)]- β -D-Glc) led the authors to postulate that terminal L-rhamnose seems to be more beneficial than D-glucose (Beit-Yannai et al. 2011). Results of their study also confirmed previous observations with regard to the general aglycone type, that furostane derivatives have lesser cytotoxic effect as compared to spirostanes.

Also Wu et al., who analysed the activity of three new saponins from *Paris polyphylla* var. *yunnanensis* against CNE cells, concluded that the presence of F ring in steroidal saponins may be the structural feature essential for their cytotoxicity (Wu et al. 2017a).

However, a study of Kang et al. showed contradictory results against human CCRF-CEM leukemia cells. From among twenty compounds (including saponins, sapogenins and sterols) isolated from *P. polyphylla*, only furostanols were active and their activity was highly potent. Both spirostanol saponins and sterols lacked any effect on this cell line (Kang et al. 2012).

In some papers included in this review the authors tried to draw conclusions referring solely to the composition and structure of the sugar moieties. This was possible when the isolated saponins differed only with respect to the sugar chain. However, the number of compounds was usually so small that it is hardly possible to consider these observations as contributing to more general statements which would be conclusive. For example, two pennogenyl saponins from *Paris quadrifolia* differing in the length and number of monosaccharides were compared on a single cell line, namely HeLa. Compound bearing a sugar chain at C-3 composed of two rhamnose unit was slightly more

active than the one with single rhamnose unit (Stefanowicz-Hajduk et al. 2015).

Zolfaghari et al. (2013) in their study of four furostane glycosides from *Allium vavilovii* have suggested that xylose instead of galactose and glucose instead of rhamnose seem to enhance cytotoxicity against J-744 (murine macrophage) and WEHI-164 (murine fibrosarcoma) cell lines.

Mechanisms of action

Similarly to what have been published in our previous work (Podolak et al. 2010), most of the steroidal saponins, which are discussed in the present review, triggered cell death by apoptosis stimulation, mainly on its intrinsic pathway. Other effects observed while testing steroidal saponins impact on cancer cells included the stimulation of autophagy, phagocytosis or oncosis, the inhibition of metastatic properties of the tested cells or angiogenesis.

Results of in vitro studies

Apoptosis stimulation

Lin et al. (2018) described the effect of protodioscin on human cervical cancer cells, trying to determine the molecular mechanism of the compound. The authors observed that protodioscin inhibited the viability of cervical cancer cells by stimulating apoptotic process in the cells, expressed by the up-regulation of caspases 8, 3 and 9, but also down-regulation of Bcl-2 expression. Moreover, protodioscin stimulated ROS and ER stress pathway in the examined cells and increased p38 and JNK levels. The authors suggest that protodioscin stimulated ER-stress dependent apoptosis in the human cervical cancer cells and the observed effect could be additionally mediated by the activation of JNK and p38 pathways (Lin et al. 2018). Terrestrosin D (hecogenin 3-*O*- β -D-Gal-(1 \rightarrow 2)-[β -D-Xyl-(1 \rightarrow 3)]- β -D-Glc-(1 \rightarrow 4)- β -D-Gal), isolated from *T. terrestris*, significantly decreased the viability of androgen-independent (DU-145, PC-3, PC-3M) and androgen-dependent (LNCaP, 22RV1) human prostate cancer cells, in dose-dependent manner (Wei et al. 2014). Moreover, the compound induced PC-3 cell cycle arrest in G1 phase and stimulated caspase-

independent apoptosis in the cells. Wang et al. indicated that macrostemonoside A (tigogenin 3-*O*- β -D-Glc-(1 \rightarrow 2)-[β -D-Glc-(1 \rightarrow 3)]- β -D-Glc-(1 \rightarrow 4)- β -D-Gal) stimulated apoptosis in colorectal cancer SW480 cells, manifesting as caspase activation, increase in proapoptotic and decrease of antiapoptotic Bcl-2 family proteins expression. Moreover, the compound induced reactive oxygen species (ROS) production in the examined cells (Wang et al. 2013c). Two studies concern the activity of saponins isolated from *P. polyphylla*. In the first one, four pennogenyl saponins PS1–PS4 were examined on a panel of human cancer and normal cell lines. The results indicated that only saponins PS1 (pennogenin 3-*O*- β -D-Glc-(1 \rightarrow 3)-[α -L-Rha-(1 \rightarrow 2)]- β -D-Glc) and PS2 (pennogenin 3-*O*- α -L-Rha-(1 \rightarrow 4)- α -L-Rha-(1 \rightarrow 4)-[α -L-Rha-(1 \rightarrow 2)]- β -D-Glc) markedly inhibited cell viability in HepG2, MCF-7 and PC-3 cells. The two compounds also induced apoptosis and caused cell cycle arrest in HepG2 cells affecting multiple targets, including mitochondrial caspase-dependent and independent pathway, cyclin-dependent kinase 1 activation or PI3K/Akt signalling (Long et al. 2015). In another study *P. polyphylla* steroidal saponins decreased the viability of human lung cancer A549 cells through both apoptosis and autophagy, with the activation of caspase-8 and 3 and PARP cleavage for the former, and up-regulation of Beclin1 and conversion from LC3 I to LC3 II for the latter process, respectively (He et al. 2014). For the same cell line, A549, an apoptosis inducement was described as the effect of a treatment with novel steroidal saponin cholestanol glucoside CG. The compound had cytotoxic effect also in PC-3 and HepG2 cells, but A549 cell line was most susceptible, with the observed ROS generation inducement and the loss of mitochondrial membrane permeability (Valayil et al. 2016). Similar effect of ROS accumulation was also described for asafilioside B (sarsasapogenin 3-*O*- β -D-Xyl-(1 \rightarrow 4)-[α -L-Ara-(1 \rightarrow 6)]- β -D-Glc), isolated from *Asparagus filicinus*. The compound additionally inhibited both viability and proliferation of HepG2 cells, by arresting the cells in G2 phase and stimulating apoptosis. The underlying mechanism included up-regulation of H-Ras and N-Ras proteins, c-Raf phosphorylation and the activation of ERK and p38. Interesting proapoptotic mechanism was recently proposed for a sapogenin–diosgenin by Chen et al. (2018). The compound was found to inhibit TAZ, one

of the transcription co-activators in Hippo signalling pathway, which may play a role as an oncogenic factor in the cells. Diosgenin also inhibited the growth and migration of human liver cancer cells (Chen et al. 2018). Its widely known glycoside—dioscin exerted rare mechanism of proapoptotic activity by triggering both intrinsic (loss of mitochondrial membrane potential, activation of tBid and Bak proteins) and extrinsic (up-regulation of death ligands and receptors) apoptosis pathways in human leukemia cells. Additionally, the compound induced the differentiation of promyelocytes to granulocytes and monocytes (Chan et al. 2018).

Oncosis stimulation

Oncosis is a non-apoptotic cell death mode, manifested as marked cell swelling, coagulation of the cytoplasm and alterations in cell cytoskeleton elements, noted within a short time after the application of the tested substance. The only report describing oncosis stimulation for steroidal saponins was published by Sun et al. (2011) for solamargine (solasodine 3-*O*- α -L-Rha-(1 \rightarrow 2)-[α -L-Rha-(1 \rightarrow 4)]- β -D-Glc)—a steroidal alkaloid glycoside in human K562 leukemia and KB squamous carcinoma cells. The authors suggested that compound initiated cell membrane blebbing, the increase in cytoplasm volume and also disrupted microtubules and actin filaments within the tested cells (Sun et al. 2011).

Angiogenesis inhibition

Terrestrosin D isolated from *T. terrestris* effectively inhibited viability of HUVEC cells and also induced cell cycle arrest and apoptosis in the cells, which suggests its antiangiogenic potential in vitro (Wei et al. 2014). Similar observations were made for ASC (diosgenin 3-*O*-[2-*O*-acetyl- α -L-Rha-(1 \rightarrow 2)]- β -D-Xyl-(1 \rightarrow 4)]- β -D-Glc), a steroidal saponin from *Ophiopogon japonicus*, which markedly inhibited the proliferation of HUVEC cells and induced G2/M phase arrest in the cells by decreasing the expression of cdc2 and cyclin B1. The compound also significantly inhibited the invasive potential of the examined cells in transwell migration and tube formation assays. Moreover, ASC was found to be a strong inhibitor of Src/Akt/mTOR-dependent metalloproteinases pathway, which may explain its antiangiogenic properties

(Zeng et al. 2015). Antiangiogenic properties were also described for another compound from the *O. japonicus*, ophiopogonin T (26-*O*- β -D-Glc (25R)-furost-5-ene-1 β ,3 β ,22 β ,26-tetraol 1-*O*- β -D-Xyl-(1 \rightarrow 3)-[α -L-Rha-(1 \rightarrow 2)]- β -D-Fuc), which inhibited tube formation of HUVEC cells (Lee et al. 2016).

Metastasis inhibition

Ophiopogonin D (25(R)-ruscogenin 1-*O*- α -L-Rha-(1 \rightarrow 2)-[β -D-Xyl-(1 \rightarrow 3)]- β -D-Fuc) isolated from *O. japonicus* significantly decreased not only the proliferation of MDA-MB-435 melanoma cells, but also decreased the cell invasion properties, probably through the inhibition of the MMP-9 matrix metalloproteinase expression and suppression of the p38/MAPK pathway. The compound inhibited also the adhesion of melanoma cells to human umbilical vascular endothelial cells and fibronectin (Zhang et al. 2015). An interesting explanation for the antiinvasive potential was proposed for dioscin in the experiment on murine B16 melanoma cells. The compound significantly affected the transcription and translation of connexin 43 via retinoid acid signalling pathway and at the same time enhanced the transporting function of connexin 43. Additionally, dioscin increased the secretion of pro-inflammatory interleukines 6 and 1 β and TNF α , but also the increase in phagocytic activity of tumor-associated macrophages was observed (Kou et al. 2017).

Multidrug resistance decreasing

Interesting study was described by Wang et al. on the potential of steroidal saponin from *Trillium tschonoskii* in reversing multidrug resistance (MDR) in hepatocellular carcinoma cells (Wang et al. 2013a). The compound not only reversed MDR in the cells but also enhanced the chemosensitization of the cells to doxorubicin, demonstrated as the significant decrease in IC₅₀ value for the anticancer drug. Moreover, the compound suppressed the P-glycoprotein expression in the drug resistant cells, which led to the accumulation of doxorubicin inside the cells, and also blocked the expression of some genes coding multidrug resistance (Wang et al. 2013a).

Results of in vivo studies

Only a small number of papers describe the in vivo effects of steroidal saponins. In one of them, after 35 days of intraperitoneal administration of 10, 50 or 100 mg kg⁻¹ daily of macrostemonoside A to BALB/c nude mice (with SW480 cells injected s.c.), a significant decrease in tumor volume and weight was noted (Wang et al. 2013c). Similar effect was described by Wei et al. (2014) for terrestrosin D, a steroidal saponin isolated from *T. terrestris*. The compound at the doses of 25 or 50 mg kg⁻¹ was administered 3 times a week for 4 weeks to BALB/c nude mice bearing PC-3 prostate cancer cells and reduced the tumor growth when compared to the control animals. Moreover, no toxic effect was noted during the treatment. Another steroidal saponin, aspafilioside B, significantly inhibited tumor growth in nude mice bearing HepG2 human hepatocellular carcinoma cells, when administered in 5 and 10 mg kg⁻¹ doses. Further analysis indicated the increase in the expression of H-Ras and N-Ras signalling proteins in the tumor cells obtained from aspafilioside B treated animals. Moreover, no side effects were observed during treatment in terms of haematological or histopathological parameters. In a similar study, dioscin revealed significant anti-metastatic effects, activating the expression of a gap junction protein connexin 43 both in metastatic lung nodes and in situ tumor animal models (Kou et al. 2017). An interesting experiment was described by Chen et al. (2016) on the effect of dioscin aglycone–diosgenin on benign prostate hyperplasia in rats (Chen et al. 2016). After 3 weeks of administration the compound at the doses of 50 and 100 mg kg⁻¹ significantly decreased prostate index and PSA level but also improved the pathological changes of the prostate in the treated animals. Moreover, diosgenin down-regulated the expression of Bcl-2 and up-regulated that of Bax and p53 in the treated animals, which suggests the efficacy of the compound in the treatment of prostate enlargement. Interesting antiangiogenic properties of ASC, isolated from *O. japonicus*, were described in matrigel plug in vivo assay. The compound significantly inhibited the formation of new blood vessels and decreased the number of the cells with the expression of PECAM-1, cell adhesion molecule, but also the number of MMP-2, MMP-9 and VEGF positive cells (Zeng et al. 2015).

Compounds with a potential as future anti-cancer therapeutic agents

Several reports indicate that some saponins/sapogenins can be considered as potential candidates for cancer treatment. In many studies on human cancer cell lines of different origin they displayed significant in vitro and in vivo activities through different signaling pathways associated with cell cycle. What is most important, apart from direct cytotoxic effect these compounds revealed also other activities, for example anti-inflammatory, that may be of importance in order to obtain the multidirectional therapeutic effect in cancer treatment. The authors of the present review have chosen five compounds: diosgenin, dioscin, polyphyllin I, paris saponin II, and timosaponin III, which, in our opinion have some interesting features, that make them especially promising for future development as anticancer agents. All selected saponins, except timosaponin AIII, share a common structural feature that is the same sapogenin–diosgenin as well as the presence of one branched sugar chain. It is noteworthy that this sapogenin itself can be considered as a potential lead compound for future development. Below, a short summary of the most interesting data referring to complex mechanisms of action is provided. Moreover, the results of the studies referring to their mechanisms of action at the molecular level, that were published in years 2010–2018 are summarized in details in Table S2 (Tab. S2)—see supplementary material. The structures of selected compounds are presented on Fig. 3.

Diosgenin (3 β ,25R)-spirost-5-en-3-ol, was discovered for the first time in *Dioscorea tokoro* in 1935 (Chen et al. 2015). Since then it has been found in numerous plants of several genera: *Dioscorea*, *Costus*, *Smilax*, *Paris*, *Alteris*, *Allium*, *Helicteres*, *Trillium*, and *Trigonella* (Sethi et al. 2018; Sobolewska et al. 2016; Deshpande and Bhalsing 2014–2015). Diosgenin exerts different pharmacological activities: hypolipemic, neuroprotective, gastro- and hepatoprotective (Jesus et al. 2016; Sethi et al. 2018). Of current interest are its anti-proliferative properties as well as anti-inflammatory effects.

Multiple molecular targets of this sapogenin are noteworthy. It is able to modulate various oncogenic processes (cancer cells proliferation, migration, apoptosis), inhibit angiogenesis, reverse multi-drug resistance in cancer cells and sensitize cancer cells to




Fig. 3 Chemical structures of some of the promising anticancer steroidal saponins/sapogenins

chemotherapy (Stehi et al. 2018; Chen et al. 2015). Diosgenin was suggested to be a good candidate for lung cancer therapy as an inhibitor of hTERT gene expression (Rahmati-Yamchi et al. 2013). Its activity against lung cancer cell line A549 was time- and dose-dependent, with the best effect after 72 h. The compound revealed also antimetastatic potential, which was observed for example on breast cancer cell line MDA-MB-231 (He et al. 2014). A significant suppression of cell migration was seen at the concentration as low as 5 μM , after only 24 h of incubation, without affecting cell proliferation. Moreover, except from downregulation of STAT3 signaling pathway and the inhibition of human hepatocellular carcinoma cells proliferation, diosgenin also potentiated paclitaxel and doxorubicin apoptotic effects (Li et al. 2010). This synergistic effect may be of special importance for further studies of this compound. Diosgenin also downregulated the peroxidation reaction and enhanced the indigenous antioxidant defense system in female rats with NMU-induced mammary cancer (Jagadeesan et al. 2012). As cancer is often related to the hyperactivity of free radicals, this activity profile completes and expands the direct impact of diosgenin on cancer cells.

Dioscin Diosgenin 3-*O*- α -L-Rha-(1 \rightarrow 4)-[α -L-Rha-(1 \rightarrow 2)]- β -D-Glc, is a spirostanol saponin found mostly in *Dioscorea* species; and also in other genera such as *Allium*, *Polygonatum*, and *Smilax* (Sobolewska et al. 2016; Rani et al. 2012; Xu et al. 2016; Wang et al. 2001; Tian et al. 2017). *Dioscorea nipponica* and *Dioscorea zingiberensis* are especially good sources of dioscin and provide raw material for the synthesis of steroid hormone drugs. Many pharmacological studies described antimicrobial, lipid-lowering, hepatoprotective, and anti-allergic activities of dioscin (Cho et al. 2013; Kwon et al. 2003; Tao et al. 2018). A large number of experimental data have confirmed not only its direct cytotoxicity towards cancer cells but also anti-inflammatory and immunoregulatory activities that may contribute to the widely reported anti-tumor effect (Tao et al. 2018; Wu et al. 2015).

Numerous studies were focused on the possible mechanism of antitumor activity of dioscin (Tab. S2). The compound was found to inhibit cancer cell viability via different pathways: G2/M cell arrest, induction of apoptosis and autophagy, downregulation of anti-apoptotic proteins, induction DNA damage

mediated by ROS (Xu et al. 2016). Dioscin treatment increased cellular apoptosis in ovarian cancer SK-OV-3 cells in a dose-dependent manner. At the concentrations of 2.5 or 5 μM it significantly decreased PI3K and phosphorylated (*p*)-AKT, VEGFR2 protein expression compared with the non-treated control group, and induced expression of p-p38 protein (Guo and Ding 2018). Dioscin induced apoptosis in SGC-7901 cells in a dose-dependent manner (Hu et al. 2011). It was more active than hCPT (IC₅₀ of 1.2 $\mu\text{g ml}^{-1}$ and IC₅₀ of 25.2 $\mu\text{g ml}^{-1}$, respectively).

Paris saponin II (PSII, formosanin C) Diosgenin 3-*O*- α -L-Rha-(1 \rightarrow 4)- α -L-Rha-(1 \rightarrow 4)-[α -L-Rha-(1 \rightarrow 2)]- β -D-Glc is one of the main active components of *Paridis rhizoma* obtained from *P. polyphylla* var. *yunnanensis* and *P. polyphylla* var. *chinensis*. This saponin was reported also in other *Paris* sp., as well as in *Cestrum*, *Allium*, *Ypsilandra*, and *Dioscorea* species (Xia et al. 2016; Ribeiro et al. 2016a, b; Sobolewska et al. 2006). With respect to the mechanisms underlying its cytotoxic activity it was found that paris saponin II induced apoptosis via activation of caspase 2, S-phase arrest, and suppressed expression of metalloproteinases MMP-1, -2, and -9 (Li et al. 2014; Man et al. 2011). Intraperitoneal administration of formosanin C at 15 and 25 mg kg⁻¹ in a xenograft mouse model of ovarian cancer led to a 46% and 70% tumor growth inhibition, respectively (Xiao et al. 2012). It is noteworthy that a combination of PSII and curcumin exerted synergic anti-cancer activity on different lung cancer cells, revealed as the increase in the cellular uptake and the bioavailability of both compounds (Man et al. 2018). Additionally, formosanin C showed immunomodulatory activity when given intraperitoneally to mice. The compound activated natural killer cells and induced interferon production (Wu et al. 1990), what can be considered as another aspect of multitargeted anticancer treatment.

Polyphyllin I (PPI) Diosgenin 3-*O*- α -L-Rha-(1 \rightarrow 2)-[β -L-Ara-(1 \rightarrow 4)]- β -D-Glc, is a spirostanol saponin isolated from the rhizomes of *P. polyphylla*. Polyphyllin I significantly suppressed in vitro proliferation of A549, NCI-H460 and SK-MES-1 cell lines with significantly low values of IC₅₀ 1.24, 2.40, and 2.33 $\mu\text{g ml}^{-1}$, respectively and the tumor growth of A549 cells in the nude mice (Kong et al. 2010). PPI inhibited also the vasculogenic mimicry formation in both hepatocellular carcinoma cell lines (HCC) and

xenografts of HCC (Xiao et al. 2018). The activity of PPI against osteosarcoma was examined both in vitro and in vivo, with interesting results. The compound was found to suppress in vitro growth of osteosarcoma 143-B and HOS cells, as well as the primary cells from a osteosarcoma patient and, what is more important, inhibited in vivo intratibial primary tumor growth in xenograft orthotopic mouse model. Moreover, it induced cell apoptosis, cell cycle arrest and inhibited the invasion and migration of osteosarcoma cells (Chang et al. 2017). Other interesting effects were obtained in the studies on concomitant administration of PPI with other compounds, including currently used chemotherapeutics. The combination of polyphyllin I and paris saponin II showed synergistic anti-tumor activity on HepG2 cells. Both compounds inhibited liver cancer growth through the induction of apoptosis, G1 phase arrest and inhibition of the cellular migration (Liu et al. 2016a). It was shown that the combined treatment of PPI and erlotinib resulted in the strengthened drug response and prolonged survival of lung cancer patients (Lou et al. 2017).

Timosaponin AIII (TAIII) Sarsasapogenin 3-*O*- β -D-Glc-(1 \rightarrow 2)- β -D-Gal, was isolated by Kawasaki et al. in 1963 (Kawasaki and Yamauchi 1963; Kawasaki et al. 1963). It is the main active ingredient of the rhizomes of *Anemarrhena asphodeloides*. The compound exerts a wide range of pharmacological effects including anti-inflammatory, antiplatelet, antithrombotic, anti-diabetic, anti-depressant, improving learning and memory deficits activities (Han et al. 2018; Cong et al. 2016). In recent years, it was found that timosaponin AIII is a promising compound that inhibits the growth of a variety of tumor cells. In different studies it was reported that TAIII may induce autophagy in cancer cells followed by apoptotic cell death, cell cycle arrest in the G0/G1 and G2/M phases, and suppresses HGF-induced invasiveness of cancer cells (Sy et al. 2008; Huang et al. 2015).

Summary

A large number of experimental data that are published each year on antitumor potential of steroidal saponins and their interesting results indicate that these natural compounds are considered to be valuable research targets in the process of development of novel chemotherapeutics for human cancers. Similarly to

previous years, the majority of experiments were performed in in vitro conditions with a relatively small number of compounds to enter in vivo studies. Assays performed on human cancer-derived cell lines were definitely predominant over animal cell models. Interestingly several cell lines were used most widely and the pool of experimental data is therefore more conclusive. These include: HepG2—human hepatocellular carcinoma, MCF-7—human breast adenocarcinoma, and A549—human lung adenocarcinoma. Based on the studies summarized in this review (see Tab. 1), it can be seen that the analyzed steroidal saponins/sapogenins revealed a differentiated cytotoxic effect. It is worth noting, however, that tests in which normal cell lines were included in the study accounted for only about 4% of all assays conducted on human cell lines. In addition, simultaneous studies on the cytotoxic activity of a given compound on cancer cells and normal cells derived from the same organ or tissue were extremely rare. Thus, it is difficult to draw more general conclusions with regard to selectivity of steroidal saponins towards cancer cells. Similarly, studies relevant to structure–activity relationships are lacking. It is noteworthy that some species containing steroidal saponins have been more frequently evaluated as sources of cytotoxic compounds in comparison to other, three of them were especially extensively analysed: *A. asphodeloides*, *P. polyphylla* var. *yunanensis*, and *O. japonicus*.

Several compounds, such as diosgenin, dioscin, paris saponin II, polyphyllin I, and timosaponin AIII seem to be specially promising as candidates for future antitumor agents. Not only their activity has been confirmed by numerous studies, but also these compounds are easily accessible for isolation being present in substantial amounts in several plant species. All of them have revealed multidirectional mechanisms of cytotoxicity as well as other effects, e.g. anti-inflammatory, that may contribute to the overall antitumor activity. Moreover, they were effective not only in in vitro assays, but also in animal models and in most cases a significant reduction in tumor size and angiogenesis was seen, especially with respect to prostate, breast, and lung cancers, of which non-small lung cancer seems to be most susceptible. Further studies on steroidal saponins with respect to their anti-cancer potential are certainly needed and worth continuing with more attention paid to compound

selectivity and synergistic effects of combinations with currently applied chemotherapeutics.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Abdel-Sattar E, Farag MA, Mahrous EA (2015) Chemical constituents from *Solanum glabratum* Dunal var. *sepicula*. *Rec Nat Prod* 9(1):94–104
- Acharya D, Mitaine-Offer A-C, Kaushik N et al (2010) Steroidal saponins from *Chlorophytum orchidastrum*. *J Nat Prod* 73(1):7–11
- Barnett D, Dean PW, Hart RJ et al (1988) Determination of contents of steroidal saponins in starfish tissues and study of their biosynthesis. *Comp Biochem Phys Part B Comp Biochem* 90(1):141–145
- Beit-Yannai E, Ben-Shabat S, Goldschmidt N et al (2011) Antiproliferative activity of steroidal saponins from *Balanites aegyptiaca*—an in vitro study. *Phytochem Lett* 4(1):43–47
- Bhuvanlakshmi G, Basappa Rangappa KS et al (2017) Breast cancer stem-like cells are inhibited by diosgenin, a steroidal saponin, by the attenuation of the Wnt β -catenin signaling via the Wnt antagonist secreted frizzled related protein-4. *Front Pharmacol* 8:124
- Böttger S, Hofmann K, Melzig MF (2012) Saponins can perturb biologic membranes and reduce the surface tension of aqueous solutions: a correlation? *Bioorg Med Chem* 20(9):2822–2828
- Challinor VL, De Voss JJ (2013) Open-chain steroidal glycosides, a diverse class of plant saponins. *Nat Prod Res* 30(3):429–454
- Challinor VL, Parsons PG, De Voss JJ (2012) Steroidal saponins from the roots of *Smilax* sp.: structure and bioactivity. *Steroids* 77(5):504–511
- Chan SH, Liang PH, Guh JH (2018) An integrated approach to elucidate signaling pathways of dioscin-induced apoptosis energy metabolism and differentiation in acute myeloid leukemia. *Naunyn Schmiedebergs Arch Pharmacol* 391(6):587–602
- Chang J, Li Y, Wang X et al (2017) Polyphyllin I suppresses human osteosarcoma growth by inactivation of Wnt/ β -catenin pathway in vitro and in vivo. *Sci Rep* 7:7605
- Chen P-S, Shih Y-W, Huang H-C et al (2011a) Diosgenin, a steroidal saponin, inhibits migration and invasion of human prostate cancer PC-3 cells by reducing matrix metalloproteinases expression. *PLoS* 6(5):e20164
- Chen P-Y, Chen C-H, Kuo C-C et al (2011b) Cytotoxic steroidal saponins from *Agave sisalana*. *Planta Med* 77(9):929–933
- Chen Y, Tang Y-M, Yu S-L et al (2015) Advances in the pharmacological activities and mechanisms of diosgenin. *Chin J Nat Med* 13(8):578–587
- Chen J, Zhang H-F, Xiong C-M et al (2016) Inhibitory effect of diosgenin on experimentally induced benign prostatic hyperplasia in rats. *J Huazhong Univ Sci Technolog Med Sci* 36(6):806–810
- Chen Z, Xu J, Wu Y et al (2018) Diosgenin inhibited the expression of TAZ in hepatocellular carcinoma. *Biochem Biophys Res Commun* 503(3):1181–1185
- Cho J, Choi H, Lee J et al (2013) The antifungal activity and membrane-disruptive action of dioscin extracted from *Dioscorea nipponica*. *Biochim Biophys Acta* 1828:1153–1158
- Chu C, Cui T, Li S et al (2018) Structure and activity of a new sapogenin from *Chlorophytum laxum* R. Br. *Chem Res Chin Univ* 34(5):732–735
- Cong Y, Wang L, Peng R et al (2016) Timosaponin AIII induces antiplatelet and antithrombotic activity via Gq-mediated signaling by thromboxane A2 receptor. *Sci Rep* 6:38757
- Deshpande HA, Bhalsing SR (2014–2015) Plant derived novel biomedical: diosgenin. *Int J Pharmacog Phytochem Res* 6(4):780–784
- Dong R, Guo J, Zhang Z et al (2018) Polyphyllin I inhibits gastric cancer cell proliferation by downregulating the expression of fibroblast activation protein alpha (FAP) and hepatocyte growth factor (HGF) in cancer-associated fibroblasts. *Biochem Biophys Res Commun* 497(4):1129–1134
- Duan C-L, Li Y-J, Li P et al (2010) Spirostanol saponins from the fibrous roots of *Ophiopogon japonicus* (Thunb.) Ker-Gawl. *Helv Chim Acta* 93(2):227–232
- Duan C-L, Li Y-J, Wang F-Y et al (2018) New steroidal glycosides from the fibrous roots of *Ophiopogon japonicus*. *J Asian Nat Prod Res* 20(8):744–751
- Eskander J, Sakka OK, Harakat D et al (2013) Steroidal saponins from the leaves of *Yucca de-smetiana* and their in vitro antitumor activity: structure activity relationships through a molecular modeling approach. *Med Chem Res* 22(10):4877–4885
- Esposito D, Munafo JP Jr, Lucibello T et al (2013) Steroidal glycosides from the bulbs of Easter lily (*Lilium longiflorum* Thunb.) promote dermal fibroblast migration in vitro. *J Ethnopharmacol* 148(2):433–440
- Faizal A, Geelen D (2013) Saponins and their role in biological processes in plants. *Phytochem Rev* 12:877–893
- Fouedjou RT, Teponno RB, Quassinti L et al (2014) Steroidal saponins from the leaves of *Cordyline fruticosa* (L.) A. Chev. and their cytotoxic and antimicrobial activity. *Phytochem Lett* 7:62–68
- Gajdus J, Kaczyński Z, Kawiak A et al (2014) Isolation and identification of cytotoxic compounds from the rhizomes of *Paris quadrifolia* L. *Pharmacogn Mag* 10(38):S324–S333

- Gergely JE, Dorsey AE, Dimri GP et al (2018) Timosaponin A-III inhibits oncogenic phenotype via regulation of PcG protein BMI1 in breast cancer cells. *Mol Carcinog* 57(7):831–841
- Ghaly NS, Nabil M, Miyase T et al (2014) Steroidal saponins from the roots of *Dracaena marginata* tam. *Pharm Lett* 6(2):132–141
- Gu L, Feng J, Xu H et al (2013) Polyphyllin I inhibits proliferation of metastasis of ovarian cancer cell line HO-8910PM in vitro. *J Trad Chin Med* 33(3):325–333
- Guo X, Ding X (2018) Dioscin suppresses the viability of ovarian cancer cells by regulating the VEGFR2 and PI3K/AKTR/MAPK signaling pathways. *Oncol Lett* 15(6):9537–9542
- Guo J, Xu C, Xue R (2015) Cytotoxic activities of chemical constituents from rhizomes of *Anemarrhena asphodeloides* and their analogues. *Arch Pharm Res* 38:598–603
- Hamid AA, Aiyelaagbe OO, Negi AS et al (2016) Bioguided isolation and antiproliferative activity of constituents from *Smilax korthalsii* A.D.C. leaves. *J Chin Chem Soc* 63:562–571
- Han W, Hou G, Liu L (2015) Polyphyllin I (PPI) increased the sensitivity of hepatocellular carcinoma HepG2 cells to chemotherapy. *Int J Clin Exp Med* 8(11):20664–20669
- Han F-Y, Song X-Y, Chen J-J et al (2018) Timosaponin AIII: a novel potential anti-tumor compound from *Anemarrhena asphodeloides*. *Steroids* 140:125–130
- He Z, Chen H, Li G et al (2014) Diosgenin inhibits the migration of human breast cancer MDA-MB-231 cells by suppressing Vav2 activity. *Phytomedicine* 21:871–876
- Hien TTT, Tuan HA, Huang DP et al (2018) Two new steroidal saponins from *Solanum procumbens*. *Nat Prod Commun* 13(10):1271–1274
- Hsieh M-J, Tsai T-L, Hsieh Y-S et al (2013) Dioscin-induced autophagy mitigates cell apoptosis through modulation of PI₃K/Akt and ERK and JNK signaling pathways in human lung cancer cell lines. *Arch Toxicol* 87(11):1927–1937
- Huang W, Zou K (2015) Cytotoxicity of the saponin TTB2 on Ewing sarcoma cells. *Exp Ther Med* 10:625–628
- Huang H-C, Lin M-K, Hwang S-Y et al (2013) Two anti-inflammatory steroidal saponins from *Dracaena angustifolia* Roxb. *Molecules* 18(8):8752
- Huang H-L, Chiang W-L, Hsiao P-C et al (2015) Timosaponin AIII mediates caspase activation and induces apoptosis through JNK1/2 pathway in human promyelocytic leukemia cells. *Tumor Biol* 36:3489–3497
- Ivanchina NV, Kicha AA, Stonik VA (2011) Steroid glycosides from marine organisms. *Steroids* 76:425–454
- Jabrane A, Ben Jannet H, Miyamoto T (2011) Spirostane and cholestane glycosides from the bulbs of *Allium nigrum* L. *Food Chem* 125(2):447–455
- Jagadeesan J, Nandakumar N, Rengarajan T et al (2012) Diosgenin, a steroidal saponin, exhibits anticancer activity by attenuating lipid peroxidation via enhancing antioxidant defense system during NMU induced breast carcinoma. *J Environ Pathol Toxicol Oncol* 31(2):121–129
- Jesus M, Martins AP, Gallardo E et al (2016) Diosgenin: recent highlights on pharmacology and analytical methodology. *J Anal Methods Chem* 2016:4156293
- Jiang S, Fan J, Wang Q et al (2016) Diosgenin induces ROS-dependent autophagy and cytotoxicity via mTOR signaling pathway in chronic myeloid leukemia cells. *Phytomedicine* 23:243–252
- Jing S-S, Wang Y, Li X-J et al (2017) Chemical constituents from *Paris thibetica* and their antitumor activities. *Chin Trad Herb Drugs* 48(6):1093–1098
- Jung O, Lee J, Lee YJ et al (2016) Timosaponin AIII inhibits migration and invasion of A549 human non-small-cell lung cancer cells via attenuations of MMP-2 and MMP-9 by inhibitions of ERK1/2, Src/FAK and β -catenin signaling pathways. *Bioorg Med Chem Lett* 26(16):3963–3967
- Kang Y-J, Chung H-J, Nam J-W et al (2011) Cytotoxic and antineoplastic activity of timosaponin A-III for human colon cancer cells. *J Nat Prod* 74(4):701–706
- Kang L-P, Liu Y-X, Eichhorn T et al (2017) Polyhydroxylated steroidal glycosides from *Paris polyphylla*. *J Nat Prod* 75(6):1201–1205
- Kawasaki T, Yamauchi T (1963) Saponins of timo (*Anemarrhena rhizoma*). II. Structure of timosaponin A-III. *Chem Pharm Bull* 11(10):1221–1224
- Kawasaki T, Yamauchi T, Itakura N (1963) Saponins of timo (*Anemarrhena rhizoma*). I *Yakugaku Zasshi* 83:892–896
- Kicha AA, Ivanchina NV, Huang TTT et al (2010) Two new asterosaponins, archasterosides A and B, from the Vietnamese starfish *Archaster typicus* and their anticancer properties. *Bioorg Med Chem Lett* 20(12):3826–3830
- Kim DS, Jeon BK, Lee YE et al (2012) Diosgenin induces apoptosis in HepG2 cells through generation of reactive oxygen species and mitochondrial pathway. *Evid Based Complement Alternat Med* 2012:981675
- Kim CS, Oh JY, Choi SU et al (2013) Chemical constituents from the roots of *Cynanchum paniculatum* and their cytotoxic activity. *Carbohydr Res* 381:1–5
- Kim KM, Im A-R, Kim SH et al (2016) Timosaponin AIII inhibits melanoma cell migration by suppressing COX-2 and in vivo tumor metastasis. *Cancer Sci* 107:181–188
- Kirmizibezkmez H, Masullo M, Festa M et al (2014) Steroidal glycosides with antiproliferative activities from *Digitalis trojana*. *Phytother Res* 28:534–538
- Kong M, Fan J, Dong A et al (2010) Effects of polyphyllin I on growth inhibition of human non-small lung cancer cells and in xenograft. *Acta Biochim Biophys* 42:827–833
- Kou Y, Ji L, Wang H et al (2017) Connexin 43 upregulation by dioscin inhibits melanoma progression via suppressing malignancy and inducing M1 polarization. *Int J Cancer* 141(8):1690–1703
- Kougan GB, Miyamoto T, Tanaka C et al (2010) Steroidal saponins from two species of *Dracaena*. *J Nat Prod* 73(7):1266–1270
- Kwon C-S, Sohn HY, Kim SH et al (2003) Anti-obesity effect of *Dioscorea nipponica* Makino with lipase-inhibitory activity in rodents. *Biosci Biotechnol Biochem* 67(7):1451–1456
- Lanzotti V (2005) Bioactive saponins from *Allium* and *Aster* plants. *Phytochem Rev* 4(2–3):95–100
- Lee SR, Han J-Y, Kang HR et al (2016) A new steroidal saponin from the tubers of *Ophiopogon japonicus* and its protective effect against cisplatin-induced renal cell toxicity. *J Braz Chem Soc* 27(4):706–711
- Lepage C, Liagre B, Cook-Moreau J et al (2010) Cyclooxygenase-2 and 5-lipoxygenase pathways in diosgenin-induced

- apoptosis in HT-29 and HCT-116 colon cancer cells. *Int J Oncol* 36:1183–1191
- Lepage C, Léger DY, Bertrand J et al (2011) Diosgenin induces death receptor-5 through activation of p38 pathway and promotes TRAIL-induced apoptosis in colon cancer cells. *Cancer Lett* 301:193–202
- Li F, Fernandez PP, Rajendran P et al (2010) Diosgenin, a steroidal saponin, inhibits STAT3 signaling pathway leading to suppression of proliferation and chemosensitization of human hepatocellular carcinoma cells. *Cancer Lett* 292(2):197–207
- Li N, Zhang L, Zeng K et al (2013) Cytotoxic steroidal saponins from *Ophiopogon japonicus*. *Steroids* 78(1):1–7
- Li Y, Man S, Li J et al (2014) The antitumor effect of formosanin C on HepG2 cell as revealed by ¹H-NMR based metabolic profiling. *Chem Biol Interact* 220:193–199
- Li Y, Wang X, He H et al (2015) Steroidal saponins from the roots and rhizomes of *Tupistra chinensis*. *Molecules* 20(8):13659
- Li L, Wu J, Zheng F et al (2016) Inhibition of EZH2 via activation of SAPK/JNK and reduction of p65 and DNMT1 as a novel mechanism in inhibition of human lung cancer cells by polyphyllin I. *J Exp Clin Oncol* 5:112–124
- Li G-B, Fu R-Q, Shen H-M et al (2017) Polyphyllin I induces mitophagic and apoptotic cell death in human breast cancer cells by increasing mitochondrial PINK1 levels. *Oncotarget* 8(6):10359–10374
- Liang F, He J-W, Zhu G-H et al (2016) New steroidal saponins with cytotoxic activities from *Smilax trinervula*. *Phytochem Lett* 16:294–298
- Lim W-Ch, Kim H, Kim Y-J et al (2017) Dioscin suppresses TGF- β 1-induced epithelial-mesenchymal transition and suppresses A549 lung cancer migration and invasion. *Bioorg Med Chem Lett* 27(15):3342–3348
- Lin C-L, Lee C-H, Chen C-M et al (2018) Protodioscin induces apoptosis through ROS-mediated endoplasmic reticulum stress via the JNK/p38 activation pathways in human cervical cancer cells. *Cell Physiol Biochem* 46:322–334
- Liu C-X, Guo Z-Y, Xue Y-H et al (2012a) Five new furostanol saponins from the rhizomes of *Tupistra chinensis*. *Fitoterapia* 83(2):323–328
- Liu C-X, Guo Z-Y, Xue Y-H et al (2012b) Tupisteroide A–C, three new polyhydroxylated steroidal constituents from the roots of *Tupistra chinensis*. *Magn Res Chem* 50(4):320–324
- Liu X, Zhang H, Niu X-F et al (2012c) Steroidal saponins from *Smilacina japonica*. *Fitoterapia* 83(4):812–816
- Liu J, Man S, Liu Z et al (2016a) A synergistic antitumor effect of polyphyllin I and formosanin C on hepatocarcinoma cells. *Bioorg Med Chem Lett* 26:4970–4975
- Liu W, Ning R, Chen R-N et al (2016b) Aspafilioside B induces G2/M cell cycle arrest and apoptosis by up-regulating H-Ras and N-Ras via ERK and p38 MAPK signaling pathways in human hepatoma HepG2 cells. *Mol Carcinog* 55(5):440–457
- Liu J, Zhang Y, Chen L et al (2017a) Polyphyllin I induces G2/M phase arrest and apoptosis in U251 human glioma cells via mitochondrial dysfunction and the JNK signaling pathway. *Acta Biochim Biophys Sin* 49(6):1–8
- Liu X, Liang J, Pan L-L et al (2017b) Six new furostanol glycosides from *Smilax glauco-china* and their cytotoxic activity. *J Asian Nat Prod Res* 19(8):754–765
- Liu X, Sun Z, Deng J et al (2018a) Polyphyllin I inhibits invasion and epithelial-mesenchymal transition via CIP2A/PP2A/ERK signaling in prostate cancer. *Int J Oncol* 53(3):1279–1288
- Liu Y, Wang M, Liu K et al (2018b) New steroidal saponins from the rhizomes of *Paris vietnamensis* and their cytotoxicity. *Molecules* 23(3):588
- Long FY, Chen YS, Zhang L et al (2015) Pennogenyl saponins induce cell cycle arrest and apoptosis in human hepatocellular carcinoma HepG2 cells. *J Ethnopharmacol* 162:112–120
- Lou W, Chen Y, Zhu K et al (2017) Polyphyllin I overcomes EMT-associated resistance to erlotinib in lung cancer cells via IL-6/STAT3 pathway inhibition. *Biol Pharm Bull* 40:1306–1313
- Lu Y, Chen C-X, Ni W et al (2010) Spirostanol tetraglycosides from *Ypsilandra thibetica*. *Steroids* 75(12):982–987
- Lu Y, Luo J, Kong L (2011) Steroidal alkaloid saponins and steroidal saponins from *Solanum surattense*. *Phytochemistry* 72(7):668–673
- Luo Q, Yang D, Qi Q et al (2018) Role of the death receptor and endoplasmic reticulum stress signaling pathways in polyphyllin I-regulated apoptosis of human hepatocellular carcinoma HepG2 cells. *Biomed Res Int* 2018:5241941
- Lv L, Zheng L, Dong D et al (2013) Dioscin, a natural steroid saponin, induces apoptosis and DNA damage through reactive oxygen species: a potential new drug for treatment of glioblastoma multiforme. *Food Chem Toxicol* 59:657–669
- Mai NT, Cuc NT, Anh HLT et al (2017) Steroidal saponins from *Datura metel*. *Steroids* 121:1–9
- Man S, Gao W, Zhang Y et al (2011) Formosanin C-inhibited pulmonary metastasis through repression of matrix metalloproteinases on mouse lung adenocarcinoma. *Cancer Biol Ther* 11(6):592–598
- Man S, Zhang L, Cui J et al (2018) Curcumin enhances the anticancer effect of paris saponin II in lung cancer cells. *Cell Prolif* 51(4):e12458
- Manase MJ, Mitaine-Offer A-C, Pertuit D et al (2012) *Solanum incanum* and *S. heteracanthum* as sources of biologically active steroid glycosides: confirmation of their synonymy. *Fitoterapia* 3(6):1115–1119
- Mao Z-J, Tang Q-J, Zhang Z-A et al (2012) Anti-proliferation and anti-invasion effects of diosgenin on gastric cancer BGC-823 cells with HIF-1 α shRNAs. *Int J Mol Sci* 13:6521–6533
- Marella CB, Sharp AE, Shemwell TA et al (2018) *Anemarrhena asphodeloides* Bunge and its constituent timosaponin-AIII induce cell cycle arrest and apoptosis in pancreatic cancer cells. *FEBS Open Bio* 8:1155–1166
- Masullo M, Pizza C, Piacente S (2016) *Ruscus* genus: a rich source of bioactive steroidal saponins. *Planta Med* 82:1513–1524
- Mirunalini S, Arulmozhi V, Shahira R (2011) Effect of diosgenin—a plant steroid on lipid peroxidation and antioxidant status in human laryngeal carcinoma cells (Hep2). *Int J Pharm Pharm Sci* 3(4):94–100

- Mosad RR, Ali MH, Ibrahim MT et al (2017) New cytotoxic steroidal saponins from *Cestrum parqui*. *Phytochem Lett* 22:167–173
- Nakamura O, Mimaki Y, Sashida Y et al (1993) Agapanthussaponins A-D, new potent cAMP phosphodiesterase inhibitors from the underground parts of *Agapanthus inaperatus*. *Chem Pharm Bull* 41(10):1784–1789
- Nie C, Zhou J, Qin X et al (2016) Diosgenin-induced autophagy and apoptosis in a human prostate cancer cell line. *Mol Med Rep* 14:4349–4359
- Ning Z, Li Y, Zhang R (2010) Effects of methyl protodioscin on $[Ca^{2+}]_i$ and ATPase activity in cardiomyocytes and analysis of mechanisms. *Zhongguo Zhong Yao Za Zhi* 35(1):80–83
- Pan Z-H, Li Y, Liu J-L et al (2012) A cytotoxic cardenolide and a saponin from the rhizomes of *Tupistra chinensis*. *Fitoterapia* 83(8):1489–1493
- Park JE, Woo KW, Choi SU et al (2014) Two new cytotoxic spirostane-steroidal saponins from the roots of *Bletilla striata*. *Helv Chim Acta* 97(1):56–63
- Pérez-Labrada K, Brouard I, Estévez S et al (2012a) Effect of C-ring modifications on the cytotoxicity of spirostan saponins and related glycosides. *Bioorg Med Chem* 20(14):4522–4531
- Pérez-Labrada K, Brouard I, Estévez S et al (2012b) New insights into the structure–cytotoxicity relationship of spirostan saponins and related glycosides. *Bioorg Med Chem* 20(8):2690–2700
- Podolak I, Galanty A, Sobolewska D (2010) Saponins as cytotoxic agents: a review. *Phytochem Rev* 9:425–474
- Poudel B, Lim S-W, Ki H-H et al (2014) Dioscin inhibits adipogenesis through the AMPK/MAPK pathway in 3T3-L1 cells and modulates fat accumulation in obese mice. *Int J Mol Med* 34:1401–1408
- Qin X-J, Yu M-Y, Ni W et al (2016) Steroidal saponins from stems and leaves of *Paris polyphylla* var. *yunnanensis*. *Phytochemistry* 121:20–29
- Qin X-J, Si Y-A, Chen Y et al (2017) Cytotoxic steroidal saponins from *Trillium kamtschaticum*. *Bioorg Med Chem Lett* 27:2267–2273
- Qin X-J, Ni W, Chen C-X et al (2018) Seeing the light: shifting from wild rhizomes to extraction of active ingredients from above-ground parts of *Paris polyphylla* var. *yunnanensis*. *J Ethnopharmacol* 224:134–139
- Rahman SU, Ismail M, Khurram M et al (2017) Bioactive steroids and saponins of the genus *Trillium*. *Molecules* 22:2156
- Rahmati-Yamchi M, Ghareghomi S, Haddadchi G et al (2013) Diosgenin inhibits hTERT gene expression in the A549 lung cancer cell line. *Asian Pac J Cancer Prev* 14(11):6945–6948
- Rani AS, Sulakshana G, Patnaik S (2012) *Costus speciosus*, an antidiabetic plant—review. *FS J Pharm Res* 1(3):52–53
- Raslan MA, Melek FR, Said AA et al (2017) New cytotoxic dihydrochalcone and steroidal saponins from the aerial parts of *Sansevieria cylindrica* Bojer ex Hook. *Phytochem Lett* 22:39–43
- Regaldo EL, Tasdemir D, Kaiser M et al (2010) Antiprotozoal steroidal saponins from the marine sponge *Pandaros acanthifolium*. *J Nat Prod* 73(8):1404–1410
- Rezgui A, Mitaine-Offer A-C, Paululat T et al (2014) Cytotoxic steroidal glycosides from *Allium flavum*. *Fitoterapia* 93:121–125
- Ribeiro PR, Araújo AJ, Costa-Lotuf LV et al (2016a) Spirostanol glucosides from the leaves of *Cestrum laevigatum* L. *Steroids* 106:35–40
- Ribeiro PR, Braz-Filho R, Araújo AJ et al (2016b) New epimeric spirostanol and furostanol-type steroidal saponins from *Cestrum laevigatum* L. *J Braz Chem Soc* 27(12):2170–2180
- Said A, Aboutabl EA, Melek FR et al (2015) Steroidal saponins and homoisoflavanone from the aerial parts of *Sansevieria cylindrica* Bojer ex Hook. *Phytochem Lett* 12:113–118
- Salgado RM, Marques-Silva MH, Gonçalves E et al (2017) Effect of oral administration of *Tribulus terrestris* extract on semen quality and body fat index of infertile men. *Andrologia* 49(5):e12655
- Selim S, Al Jaouni S (2015) Anticancer and apoptotic effects on cell proliferation of diosgenin isolated from *Costus speciosus* (Koen.) Sm. *BMC Complem Altern Med* 15(1):301–307
- Sethi G, Shanmugam MK, Warriar S et al (2018) Pro-apoptotic and anti-cancer properties of diosgenin: a comprehensive and critical review. *Nutrients* 10:456–645
- Shen S, Li G, Huang J et al (2012) Steroidal saponins from *Fritillaria pallidiflora* Shrenk. *Fitoterapia* 83:785–794
- Shen H-Y, Zuo W-J, Wang H et al (2014) Steroidal saponins from dragon's blood of *Dracaena cambodiana*. *Fitoterapia* 94:94–101
- Shi Y-M, Yang L, Geng Y-D et al (2015) Polyphyllin I induced-apoptosis is enhanced by inhibition of autophagy in human hepatocellular carcinoma cells. *Phytomedicine* 22(13):1139–1149
- Shi F, Lu L, Li Y et al (2018) Abstract 2871: diosgenin inhibits the proliferation of MCF-7 breast cancer cells through the demethylation of miR-145 gene. *Cancer Res* 78(13 Suppl):2871
- Shu J, Zhu G, Huang G et al (2017) New steroidal saponins with L-arabinose moiety from the rhizomes of *Smilax scobini-caulis*. *Phytochem Lett* 21:194–199
- Shwe HH, Aye M, Sein MM et al (2010) Cytotoxic steroidal saponins from the rhizomes of *Tacca integrifolia*. *Chem Biodiver* 7(3):610–622
- Sobolewska D, Janeczko Z, Kisiel W et al (2006) Steroidal saponins from the underground parts of *Allium ursinum* L. and their cytostatic and antimicrobial activity. *Acta Pol Pharm Drug Res* 63(3):219–223
- Sobolewska D, Michalska K, Podolak I et al (2016) Steroidal saponins from the genus *Allium*. *Phytochem Rev* 15:1–35
- Sohn H-Y, Kum E-J, Ryu H-Y et al (2006) Antifungal activity of fistulosides, steroidal saponins from *Allium fistulosum* L. *J Life Sci* 16(2):310–314
- Stefanowicz-Hajduk J, Kawiak A, Gajdus J et al (2011) Cytotoxic activity of *Paris quadrifolia* extract and isolated saponin fractions against human tumor cell lines. *Acta Biol Cracov Series Bot* 53(2):127–131
- Stefanowicz-Hajduk J, Bartoszewski R, Bartoszevska S et al (2015) Pennogenyl saponins from *Paris quadrifolia* L. induce extrinsic and intrinsic pathway of apoptosis in human cervical cancer HeLa cells. *PLoS ONE* 10(8):e0135993

- Sun L, Zhao Y, Yuan H et al (2011) Solamargine, a steroidal alkaloid glycoside, induces oncosis in human K562 leukemia and squamous cell carcinoma KB cells. *Cancer Chemother Pharmacol* 67(4):813–821
- Sun Y, Qiu H, Ou M et al (2016) Taccaoside induces apoptosis in hepatocellular carcinoma cell lines. *J Int Med Res* 44(6):395–1402
- Sy LK, Yan SC, Lok CN et al (2008) Timosaponin A-III induces preceding mitochondria-mediated in HeLa cancer cells. *Cancer Res* 68(24):10229–10237
- Tabopda TK, Mitaine-Offer A-C, Tanaka C et al (2014) Steroidal saponins from *Dioscorea preusii*. *Fitoterapia* 97:198–203
- Tabopda TK, Mitaine-Offer A-C, Paululat T et al (2016) Steroidal saponins from *Chlorophytum deistelianum*. *Phytochemistry* 126:34–40
- Tang Y, Li N, Duan J-A et al (2013) Structure, bioactivity, and chemical synthesis of OSW-1 and other steroidal glycosides in the genus *Ornithogalum*. *Chem Rev* 113(7):5480–5514
- Tang Y, He X, Ye M et al (2015) Cardioprotective effect of total saponins from three medicinal species of *Dioscorea* against isoprenaline-induced myocardial ischemia. *J Ethnopharmacol* 175:451–455
- Tao X, Xu L, Yin L et al (2017) Dioscin induces prostate cancer cell apoptosis through activation of estrogen receptor- β . *Cell Death Dis* 8:e2989
- Tao X, Yin L, Peng J (2018) Dioscin: a diverse acting natural compound with therapeutic potential in metabolic diseases, cancer, inflammation and infections. *Pharmacol Res* 137:259–269
- Tapondjou LA, Jenett-Siems K, Böttger S et al (2013) Steroidal saponins from the flowers of *Dioscorea bulbifera* var. *sativa*. *Phytochemistry* 95:341–350
- Teponno RB, Tanaka C, Jie B et al (2016) Trifasciatisides A-J, steroidal saponins from *Sansevieria trifasciata*. *Chem Pharm Bull* 64(9):1347–1355
- Teponno RB, Dzoyem JP, Nono RN et al (2017) Cytotoxicity of secondary metabolites from *Dracaena viridiflora* Engl & Krause and their semisynthetic analogues. *Rec Nat Prod* 11(5):421–430
- Tian L-W, Zhang Z, Long H-L et al (2017) Steroidal saponins from the genus *Smilax* and their biological activities. *Nat Prod Bioprospect* 7:283–298
- Timité G, Mitaine-Offer A-C, Miyamoto T et al (2013) Structure and cytotoxicity of steroidal glycosides from *Allium schoenoprasum*. *Phytochemistry* 88:61–66
- Tong Q-Y, Qing Y, Shu D et al (2011) Deltonin, a steroidal saponin, inhibits colon cancer cell growth in vitro and tumor growth in vivo via induction of apoptosis and antiangiogenesis. *Cell Physiol Biochem* 27(3–4):233–242
- Tong Q-Y, He Y, Zhao Q-B et al (2012) Cytotoxicity and apoptosis-inducing effect of steroidal saponins from *Dioscorea zingiberensis* Wright against cancer cells. *Steroids* 77(12):1219–1227
- Tong Q, Qing Y, Wu Y et al (2014) Dioscin inhibits colon tumor growth and tumor angiogenesis through regulating VEGFR2 and AKT/MAPK signaling pathways. *Toxicol Appl Pharmacol* 281(2):166–173
- Valayil JM, Kuriakose GC, Jayabaskaran C (2016) Isolation, purification and characterization of a novel steroidal saponin cholestenol glucoside from *Lasiodiplodia theobromae* that induces apoptosis in A549 cells. *Anticancer Agents Med Chem* 16(7):865–874
- Wang W, Meng H (2015) Cytotoxic, anti-inflammatory and hemostatic spirostane-steroidal saponins from the ethanol extract of the roots of *Bleilla striata*. *Fitoterapia* 101:12–18
- Wang Z, Zhou J, Ju Y et al (2001) Effects of two saponins extracted from the *Polygonatum zanlanscianense* pamp on the human leukemia (HL-60) cells. *Biol Pharm Bull* 24(2):159–162
- Wang K-W, Zhang H, Shen L-Q et al (2011) Novel steroidal saponins from *Liriope graminifolia* (Linn) Baker with anti-tumor activities. *Carbohydr Res* 346(2):253–258
- Wang H, Zhai Z, Li N et al (2013a) Steroidal saponin of *Trillium tschonoskii*. Reverses multidrug resistance of hepatocellular carcinoma. *Phytomedicine* 20(11):985–991
- Wang N, Feng Y, Zhu M et al (2013b) A novel mechanism of XIAP degradation induced by timosaponin AIII in hepatocellular carcinoma. *Biochim Biophys Acta* 1833:2890–2899
- Wang Y, Tang Q, Jiang S et al (2013c) Anti-colorectal cancer activity of macrostemonoside A mediated by reactive oxygen species. *Biochem Biophys Res Commun* 441:825–830
- Wang L, Jiang X-L, Zhang W-M et al (2017a) Homo-*α*-cholestane, furostane and spirostane saponins from the tubers of *Ophiopogon japonicus*. *Phytochemistry* 136:125–132
- Wang Y, Xu L, Lou L-L et al (2017b) Timosaponin AIII induces and autophagy in human melanoma A375-S2 cells. *Arch Pharm Res* 40(1):69–78
- Wang J, Song H, Wu X et al (2018) Steroidal saponins from *Vernonia amygdalina* Del. and their biological activity. *Molecules* 23:579
- Watanabe S, Suzuki T, Hara F et al (2017) Polyphyllin D, a steroidal saponin in *Paris polyphylla*, induces apoptosis and necroptosis cell death of neuroblastoma cells. *Pediatr Surg Int* 33:713–719
- Wei Y, Xu Y, Han X et al (2013) Anti-cancer effects of dioscin on three kinds of human lung cancer cell lines through inducing DNA damage and activating mitochondrial signal pathway. *Food Chem Toxicol* 59:118–128
- Wei S, Fukuhara H, Chen G et al (2014) Terrestrosin D, a steroidal saponin from *Tribulus terrestris* L. inhibits growth and angiogenesis of human prostate cancer in vitro and in vivo. *Pathobiology* 81:123–132
- Wen Y-S, Ni W, Qin X-J et al (2015) Steroidal saponins with cytotoxic activity from the rhizomes of *Paris polyphylla* var. *yunnanensis*. *Phytochem Lett* 12:31–34
- Wu RT, Chiang HC, Fu WC et al (1990) Formosanin-C, an immunomodulator with antitumor activity. *Int J Immunopharmacol* 12(7):777–786
- Wu J-J, Cheng K-W, Zuo X-F et al (2010) Steroidal saponins and ecdysterone from *Asparagus filicinicus* and their cytotoxic activities. *Steroids* 75(10):734–739
- Wu X, Wang L, Wang G-C et al (2012a) New steroidal saponins and sterol glycosides from *Paris polyphylla* var. *yunnanensis*. *Planta Med* 78(15):1667–1675

- Wu X, Wang L, Wang H et al (2012b) Steroidal saponins from *Paris polyphylla* var. *yunnanensis*. *Phytochemistry* 81:133–143
- Wu S, Xu H, Peng J et al (2015) Potent anti-inflammatory effect of dioscin mediated by suppression of TNF- α -induced VCAM-1, ICAM-1 and EL expression via the NF- κ B pathway. *Biochimie* 110:62–72
- Wu X, Chen N-H, Zhang Y-B et al (2017a) A new steroid saponin from the rhizomes of *Paris polyphylla* var. *yunnanensis*. *Chem Nat Comp* 53(1):93–98
- Wu Y, Wang X-M, Bi S-X et al (2017b) Novel cytotoxic steroidal saponins from the roots of *Liriope muscari* (Decne.) L.H. Bailey. *RSC Adv* 7(23):13696–13706
- Wu Y, Bi S-X, Huang Z et al (2018) Novel steroidal saponins with cytotoxic activities from the roots of *Ophiopogon japonicus* (L. f.) Ker-Gawl. *RSC Adv* 8:2498–2505
- Xia L, Ouyang P-Y, Gao W et al (2016) Rapid and sensitive determination of the major steroidal saponins of *Ypsilandra tibetica* Franch by ultra high-performance liquid chromatography coupled with quadrupole mass spectrometry. *J Chromatogr Sci* 54(6):1010–1015
- Xiang S, Zou P, Wu J et al (2018) Crosstalk of NF- κ B/P65 and lncRNA HOTAIR-mediated repression of MUC1 expression contribute to synergistic inhibition of castration-resistant prostate cancer by polyphyllin I-enzalutamide combination treatment. *Cell Physiol Biochem* 47(2):759–773
- Xiao X, Zou J, Bui-Nguyen TM et al (2012) Paris saponin II of *Rhizoma Paridis*—a novel inducer of apoptosis in human ovarian cancer cells. *BioSci Trends* 6(4):201–211
- Xiao T, Zhong W, Zhao J et al (2018) Polyphyllin I suppresses the formation of vasculogenic mimicry via Twist1/VE-cadherin pathway. *Cell Death Dis* 9:906–919
- Xu X-H, Li T, Fong CMV et al (2016) Saponins from Chinese medicines as anticancer agents. *Molecules* 21(10):1326
- Yang M, Zou J, Zhu H et al (2015) Paris saponin II inhibits human ovarian cancer cell-induced angiogenesis by modulating NF- κ B signaling. *Oncol Rep* 33:2190–2198
- Yang B-Y, Zhang J, Liu Y et al (2016a) Steroidal saponins from the rhizomes of *Anemarrhena asphodeloides*. *Molecules* 21(8):1075
- Yang J, Wang P, Wu W et al (2016b) Steroidal saponins in oat bran. *J Agric Food Chem* 64(7):1549–1556
- Yang B-Y, Bi X-Y, Liu Y et al (2017) Four new glycosides from the rhizome of *Anemarrhena asphodeloides*. *Molecules* 22(11):1995
- Yang B, Xu B, Zhao H et al (2018) Dioscin protects against coronary heart disease by reducing oxidative stress and inflammation via Sirt1/Nrf2 and p38 MAPK pathways. *Mol Med Rep* 18(1):973–980
- Yu H, Liu Y, Niu C et al (2018a) Diosgenin increased DDX3 expression in hepatocellular carcinoma. *Am J Transl Res* 10(11):3590–3599
- Yu S, Wang L, Cao Z et al (2018b) Anticancer effect of polyphyllin I in colorectal cancer cells through ROS-dependent autophagy and G2/M arrest mechanism. *Nat Prod Res* 32(12):1489–1492
- Zaki AA, Ali Z, Wang Y-H et al (2017) Cytotoxic steroidal saponins from *Panicum turgidum* Forssk. *Steroids* 125:14–19
- Zeng K-W, Song F-J, Li N et al (2015) ASC, a bioactive steroidal saponin from *Ophiopogon japonicus*, inhibits angiogenesis through interruption of src tyrosine kinase-dependent matrix metalloproteinase pathway. *BCPT* 116(2):115–123
- Zhang C, Feng S, Zhang L et al (2013) A new cytotoxic steroidal saponin from the rhizomes and roots of *Smilax scobinicaulis*. *Nat Prod Res* 27(14):1255–1260
- Zhang Y, Han Y, Zhai K et al (2015) Ophiopogonin D suppresses MDA-MB-435 cell adhesion and invasion by inhibiting matrix metalloproteinase-9. *Mol Med Rep* 12(1):1493–1498
- Zhang L, Man S, Wang Y et al (2016) Paris saponin II induced apoptosis via activation of autophagy in human lung cancer cells. *Chem Biol Interact* 253:125–133
- Zhang Y-W, Zhao Y-F, Wang Y-R et al (2017) Steroidal saponins with cytotoxic activities from the rhizomes of *Anemarrhena asphodeloides* Bge. *Phytochem Lett* 20:102–105
- Zhang S, Pang H, Sun M et al (2018a) Timosaponin AIII inhibits the growth of human leukaemia cells HL-60 by downregulation of PI3K/AKT and Wnt/ β -catenin pathways. *Biotechnol Biochem* 32(1):150–155
- Zhang Y, Huang P, Liu X et al (2018b) Polyphyllin I inhibits growth and invasion of cisplatin-resistant gastric cancer cells by partially inhibiting CIP2A/PP2A/Akt signaling axis. *J Pharmacol Sci* 137:305–312
- Zhang Y-S, Ma Y-L, Thakur K et al (2018c) Molecular mechanism and inhibitory targets of dioscin in HepG2 cells. *Food Chem Toxicol* 120:143–154
- Zhao X, Tao X, Peng J (2016) Dioscin induces apoptosis in human cervical carcinoma HeLa and SiHa cells through ROS-mediated DNA damage and the mitochondrial signaling pathway. *Molecules* 21(6):730
- Zhao Y-Z, Zhang Y-Y, Han H et al (2018) Advances in the antitumor activities and mechanisms of action of steroidal saponins. *Chin J Nat Med* 16(10):732–748
- Zhiyu W, Yue C, Neng W et al (2012) Dioscin induces cancer cell apoptosis through elevated oxidative stress mediated by downregulation of peroxiredoxins. *Cancer Biol Ther* 13(3):138–147
- Zolfaghari B, Sadeghi M, Troiano R et al (2013) Vavilosides A1/A2–B1/B2, new furostane glycosides from the bulbs of *Allium vavilovii* with cytotoxic activity. *Bioorg Med Chem* 21(7):1905–1910
- Zuo S-Q, Liu Y-N, Yang Y (2018) Aspidosaponins A–D, four new steroidal saponins from the rhizomes of *Aspidistra elatior* Blume and their cytotoxicity. *Phytochem Lett* 25:126–131

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.