

Fear Learning Increases the Number of Polyribosomes Associated with Excitatory and Inhibitory Synapses in the Barrel Cortex

Malgorzata Jasinska¹, Ewa Siucinska², Ewa Jasek¹, Jan A. Litwin¹, Elzbieta Pyza³, Malgorzata Kossut^{2,4*}

1 Department of Histology, Jagiellonian University Medical College, Krakow, Poland, **2** Department of Molecular and Cellular Neurobiology, Nencki Institute of Experimental Biology, Warsaw, Poland, **3** Department of Cell Biology and Imaging, Institute of Zoology, Jagiellonian University, Krakow, Poland, **4** Warsaw School of Social Psychology, Warsaw, Poland

Abstract

Associative fear learning, resulting from whisker stimulation paired with application of a mild electric shock to the tail in a classical conditioning paradigm, changes the motor behavior of mice and modifies the cortical functional representation of sensory receptors involved in the conditioning. It also induces the formation of new inhibitory synapses on double-synapse spines of the cognate barrel hollows. We studied density and distribution of polyribosomes, the putative structural markers of enhanced synaptic activation, following conditioning. By analyzing serial sections of the barrel cortex by electron microscopy and stereology, we found that the density of polyribosomes was significantly increased in dendrites of the barrel activated during conditioning. The results revealed fear learning-induced increase in the density of polyribosomes associated with both excitatory and inhibitory synapses located on dendritic spines (in both single- and double-synapse spines) and only with the inhibitory synapses located on dendritic shafts. This effect was accompanied by a significant increase in the postsynaptic density area of the excitatory synapses on single-synapse spines and of the inhibitory synapses on double-synapse spines containing polyribosomes. The present results show that associative fear learning not only induces inhibitory synaptogenesis, as demonstrated in the previous studies, but also stimulates local protein synthesis and produces modifications of the synapses that indicate their potentiation.

Citation: Jasinska M, Siucinska E, Jasek E, Litwin JA, Pyza E, et al. (2013) Fear Learning Increases the Number of Polyribosomes Associated with Excitatory and Inhibitory Synapses in the Barrel Cortex. PLoS ONE 8(2): e54301. doi:10.1371/journal.pone.0054301

Editor: Barbara Bardoni, CNRS UMR7275, France

Received: September 25, 2012; **Accepted:** December 10, 2012; **Published:** February 14, 2013

Copyright: © 2013 Jasinska et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: The study was supported by Iuventus Plus grant IP2010 014970 from the Polish Ministry of Science and Higher Education to MJ. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: kossut@nencki.gov.pl

Introduction

Sensory experience and learning can induce changes in the connections between neurons. Alterations of synaptic and spine density were observed in several learning paradigms [1,2,3,4,5,6] as well as in case of enhanced or decreased sensory stimulation [7,8,9,10,11]. Modifications of the neuronal activity have also been shown to change synaptic efficiency by inducing a remodeling of the existing synapses [12,13,14].

We have investigated microstructure of somatosensory barrel cortex following fear learning-induced plastic change. Short-lasting classical conditioning, in which stimulation of a row of whiskers is paired with the administration of a mild electric shock to the tail, produces plastic modification in layer IV of the barrel cortex. This part of somatosensory cortex offers an excellent model to study the mechanisms of learning and memory because of its clearly defined structure, and the ease of inducing plasticity via a fear learning paradigm [15,16]. 2-deoxyglucose autoradiography revealed enlargement of functional cortical representation of the whiskers stimulated during conditioning [16]. We have shown that this conditioning paradigm changes motor behavior of mice by decreasing the number of head movements in response to the conditioned stimulus (a phenomenon akin to freezing observed in auditory fear conditioning) [17]. It is also associated with an

autonomic response, conditioned bradycardia [16]. The learning-induced plastic change of cortical representation is associated with a number of alterations in the inhibitory neurotransmission. It is paralleled by a rapid increase of GAD67 mRNA, and increased density of GAD and GABA immunoreactive cells in the barrels of the row receiving input from the stimulated whiskers [18,19]. Conditioning led to the enhancement of inhibitory synaptogenesis [20], as well as to increased inhibitory transmission [21] in layer IV of the “trained” barrel. It also affected excitatory transmission by changing subunit composition of NMDA receptors. Increase in mRNA and protein of NMDA2A subunit was found in the barrels receiving the conditioned stimulus [22].

The presence of polyribosomes near the synapses in the dendritic spines or shafts suggests the involvement of local protein synthesis in the process of synapse modification [12,23,24]. The newly synthesized protein molecules enhance the level of the existing proteins or replace them [12,25]. Thus, an increase in the number of polyribosomes near the synapses might indicate either a formation of new synapses or a modification of the existing ones as a result of neuronal activity [14].

In the present study, the polyribosomes in the barrel cortex were investigated following the conditioning, by electron microscopy of serial sections and 3D reconstruction of dendritic spines. We found




Figure 1. Location of polyribosomes in dendrites of B2 barrel. Electron micrographs of B2 barrel hollow showing polyribosomes (arrows) located in the dendritic spine and associated with excitatory synapse (A), located in the dendritic spine and associated with inhibitory synapse (B), located in the dendritic shaft and associated with excitatory synapse (C) and located in the dendritic shaft and associated with inhibitory synapse (D). Scale bar: 0.5 μm . doi:10.1371/journal.pone.0054301.g001

that classical fear conditioning resulted in an increased density of polyribosomes in dendritic spines and shafts associated with inhibitory and excitatory synapses in layer IV barrels receiving the conditioned stimulus. We also observed conditioning-induced enlargement of post-synaptic density (PSD) area of the excitatory and inhibitory synapses on spines containing polyribosomes, which might serve to increase the strength of synapses.

Materials and Methods

We examined the same collection of ultrathin sections which was used in our previous study [20].

Animals

As described previously [20] the experiments were performed on Swiss Webster mice aged 6–7 weeks, raised in standard conditions. All experiments were compliant with the European Communities Council Directive of 24 November 1986 (86/609/EEC) and were approved by the Animal Care and Use Committees of the Polish Academy of Sciences and the Jagiellonian University.

Behavioral training

The mice ($N = 21$) were divided into a trained group ($N = 14$) and a control group ($N = 7$). The trained mice were either conditioned ($N = 7$) or pseudoconditioned ($N = 7$). All animals were habituated in a homemade restrainer which held the mouse neck stationary, leaving the rest of the body, including the head, free. During the habituation period, the mice spent in the restrainer 10 min per day for 3 weeks.

After habituation, the mice from the conditioned group were trained using the classical conditioning paradigm, exactly as described in [16]. Manual stimulation of the selected whiskers (B row; conditioned stimulus, CS) on one side of the snout was paired with a mild electric shock (unconditioned stimulus, UCS) [16]. The pairing procedure included three sweeps along the entire whisker row with a small paintbrush lasting 9 s (CS) and at the last 0.5 s of CS a tail shock (UCS) of 0.5 mA was applied. The CS+UCS pairings were repeated, four times per minute for 10 min. The conditioning sessions were carried out for 3 consecutive days. In the pseudoconditioned animals (random

pairing of CS and UCS), the number and frequency of stimuli applied were the same. The behavior of animals was filmed and their head movements in response to CS were analyzed.

Transmission electron microscopy

Twenty-four hours after the completion of the training, the mice were anesthetized with Vetbutal (25–30 mg/kg body weight; Biowet Pulawy) and perfused through the heart with 20 ml of rinse buffer (0.2% glutaraldehyde and 2% paraformaldehyde in 0.1 M phosphate buffer, pH 7.4) followed by 100–150 ml of fixative (2.5% glutaraldehyde and 2% paraformaldehyde in 0.1 M phosphate buffer, pH 7.4). The brains were removed immediately after perfusion and were left in the same fixative for 24 h at 4°C.

Next, after washing in 0.1 M phosphate buffer (pH 7.4), 60 μm tangential vibratome sections were cut from the barrel cortex. Sections were examined under a stereomicroscope (Nikon Optiphot) and those containing the barrel field cortex were collected for further processing. The sections were washed in 0.1 M cacodylate buffer (pH 7.4), postfixed with 1% osmium tetroxide in 0.1 M cacodylate buffer (pH 7.4) twice (the first time with 1.5% potassium ferrocyanide), washed in 70% ethanol containing 1% uranyl acetate, and after dehydration in a graded series of ethanol, embedded in Epon resin (Polysciences) between two silicon-coated glass slides.

The region of B2 and B3 barrels was identified according to the procedure described previously [20] and trimmed for ultrathin sectioning. A series of 30 to 50 successive sections (60–70 nm thick) were cut from each sample. The sections were collected on formvar-coated copper-palladium slots and contrasted with 1% lead citrate. For an examination of polyribosome density, 6–8 series, consisting of 3 serial electron micrographs each obtained from successive sections containing the B2 barrel central area in which cell bodies are sparse, were taken at 14K under JEOL 100SX transmission electron microscope (JEOL, Japan). Ten to twelve serial electron micrographs were taken from successive sections for a 3D reconstruction of the dendritic spines. The micrographs were initially aligned in Adobe Photoshop CS software, in which stacks of serial images were taken at the final magnification of 30 K.

Quantitative analysis of polyribosomes

The quantitative analysis of the polyribosomes was carried out using NIH Image J Cell Counter software (<http://rsb.info.nih.gov/ij/>) by placing a grid of a two-dimensional sampling frame over the stack of serial sections. The polyribosomes were counted per area unit (μm^2), because their size usually did not exceed the thickness of a single ultrathin section. Each polyribosome was counted only once in the stack and only the polyribosomes located fully within the frame or intersecting the left and the upper borderlines of the frame were included. Synapses and spines were defined according to [8]. The accordance of synapse morphology (asymmetric/symmetric) with the character of synapses (excitatory/inhibitory) was previously confirmed by immunocytochemistry of presynaptic markers in the same material [20]. Astrocytic processes were distinguished from dendrites according to [26]. Since polyribosomes located in cell bodies were omitted from the analysis, the total density of polyribosomes reflects the number of polyribosomes in dendritic spines and shafts counted in a defined area of B2 barrel. If the polyribosomes were located under the synaptic profiles without any intervening organelles and within the distance of $\leq 2 \mu\text{m}$ as measured from the center of the synapse, they were considered as synapse-associated [14,27].

The density of polyribosomes associated with excitatory and inhibitory synapses and not associated with synapses, located in




Figure 2. Density of polyribosomes in dendritic spines. Conditioning increases the density of polyribosomes located in spines and associated with excitatory (ANOVA, $p < 0.001$) and inhibitory synapses (ANOVA, $p < 0.001$). **A.** Total density of polyribosomes located in dendritic spines in: control, conditioned (CS+UCS) and pseudoconditioned (PSEUDO) group. **B.** Density of polyribosomes located in spines and associated with excitatory synapses. **C.** Density of polyribosomes located in spines and associated with inhibitory synapses. All graphs show means \pm SD (** $p < 0.01$, *** $p < 0.001$).

doi:10.1371/journal.pone.0054301.g002

dendritic shafts and dendritic spines was calculated according to the stereological formula $N_A = \Sigma Q^- / A$, where ΣQ^- is the number of polyribosomes counted in the entire area A [28]. The counting was done blind – the observer did not know whether the micrographs were taken from conditioned, pseudoconditioned or control animals.

Morphological analysis of single- and double-synapse spines

Images of 37 single-synapse spines and 41 double-synapse spines from the control, the conditioned group and the pseudoconditioned group were selected for PSD area measurements and 3D reconstruction. Obliquely or longitudinally sectioned dendritic spines were not taken into consideration. To assess PSD area, three area measurements from each micrograph containing profiles of the selected spines were made using NIH Image J software. 3D reconstructions of the spines were performed using 3D Studio Max software (Discreet Logic, Montreal, Canada) and the location of polyribosomes in the dendritic spine areas (head, neck or base) was estimated.

Statistical analysis

All the data were analysed using GraphPad Prism 4.0 software (GraphPad Software Inc., USA).

To compare the effects of conditioning on polyribosome density across the experimental groups, one-way ANOVA with *post hoc* Tukey test was used after pretesting for normality and homogeneity. If the absence of Gaussian distribution was noted

(D'Agostino and Pearson omnibus normality test) or if variances were significantly different (Bartlett's test for equal variances), the data were analysed by use of the Kruskal-Wallis test with *post hoc* Dunn's test. Differences in the location of polyribosomes in dendritic spines between the control, the conditioned and the pseudoconditioned group were compared by chi-square test. Two-way ANOVA test was used to analyse the combined effects of the training and of the polyribosome distribution on PSD area.

Results

In this study, we identified the synapses in B2 barrel of mouse somatosensory cortex and divided them according to their location into synapses on dendritic shafts and on dendritic spines. The synapses placed on dendritic shafts were either symmetric (presumptive inhibitory) or asymmetric (presumptive excitatory). All synapses located on single-synapse spines were excitatory, whereas double-synapse spines always had one excitatory and one inhibitory synapse, as demonstrated in earlier studies [8,29]. We have not found spines with more than two synapses.

Behavioral effects

The behavioral effects of conditioning in mice used in this experiment were described previously [20]. We examined CS-linked changes in head turning towards the stimulating brush. We considered them as a marker of acquiring a predictive value by the CS. We found that the conditioned mice showed a significant decrease ($54 \pm -2.1\%$, $p < 0.05$, Mann Whitney test) in head




Figure 3. Density of polyribosomes in dendritic shafts. Conditioning induces an increase in the density of polyribosomes associated with inhibitory synapses and located in dendritic shafts (ANOVA, $p < 0.001$), while pseudoconditioning decreases the density of polyribosomes not associated with synapses and located in dendritic shafts (ANOVA, $p < 0.01$). **A.** Density of polyribosomes associated with synapses and located in dendritic shafts in: control, conditioned (CS+UCS) and pseudoconditioned (PSEUDO) group. **B.** Density of polyribosomes associated with inhibitory synapses and located in dendritic shafts. **C.** Density of polyribosomes associated with excitatory synapses and located in dendritic shafts. **D.** Density of polyribosomes not associated with synapses and located in dendritic shafts. All graphs show means \pm SD (** $p < 0.01$, *** $p < 0.001$). doi:10.1371/journal.pone.0054301.g003

mobility, that developed during the process of conditioning in response to conditioned stimulus (“minifreezing”, [17]). No decrease of head mobility was found in the pseudoconditioned group ($109 \pm 8.9\%$).

Sampling areas

Polyribosomes were counted in the following total tissue areas: $2510.06 \pm 167.34 \mu\text{m}^2$ (mean area per animal $358.58 \pm 61.73 \mu\text{m}^2$) in the control group; $2210.07 \pm 147.34 \mu\text{m}^2$ (mean area per animal $315.72 \pm 29.78 \mu\text{m}^2$) in the conditioned group; $2147.16 \pm 143.14 \mu\text{m}^2$ (mean area per animal $306.74 \pm 23.60 \mu\text{m}^2$) in the pseudoconditioned group. These sampling areas were not significantly different across the treatment groups (one-way ANOVA, $p = 0.0714$, $F = 3.067$, $N = 21$, total $df = 20$).

Density of polyribosomes

Individual ribosomes were identified as small (18–25 nm) structures with round and opaque centers surrounded by lighter edges [23,25]. Three or more ribosomes forming a spiral or an irregular cluster were identified as polyribosome. Polyribosomes were classified according to their location in dendritic spines (Fig. 1A, B) or shafts (Fig. 1C, D).

Polyribosomes associated with synapses. The distances between synapse-associated polyribosomes and synapses were not significantly different across the groups (asymmetric synapses: control group $0.045\text{--}0.895 \mu\text{m}$, conditioned group 0.055--

$0.940 \mu\text{m}$, pseudoconditioned group $0.038\text{--}0.852 \mu\text{m}$; one-way ANOVA, $p = 0.8956$, $F = 0.1103$, $N = 21$, total $df = 20$; symmetric synapses: control group $0.047\text{--}0.699 \mu\text{m}$, conditioned group $0.025\text{--}0.936 \mu\text{m}$, pseudoconditioned group $0.025\text{--}0.673 \mu\text{m}$; one-way ANOVA, $p = 0.1161$, $F = 2.198$, $N = 21$, total $df = 20$). The vast majority (90%) of synapse-associated polyribosomes located in dendritic shafts and spines was within $0.6 \mu\text{m}$ from the synapses. All polyribosomes located in dendritic spines were not farther than $0.940 \mu\text{m}$ away from the centers of synapses.

Density of polyribosomes located in dendritic spines. The density of polyribosomes located in the dendritic spines increased more than fourfold after conditioning (control group: $0.010 \pm 0.006/\mu\text{m}^2$; conditioned group: $0.046 \pm 0.014/\mu\text{m}^2$; Kruskal-Wallis test, $p = 0.0012$, $H = 13.39$, $N = 21$, total $df = 20$) but not after pseudoconditioning ($0.010 \pm 0.006/\mu\text{m}^2$) (Fig. 2A). Whisker conditioning increased the density of polyribosomes located in spines and associated with excitatory synapses almost fourfold, from $0.010 \pm 0.007/\mu\text{m}^2$ in the control animals to $0.038 \pm 0.011/\mu\text{m}^2$ (one-way ANOVA, $p < 0.0001$, $F = 26.28$, $N = 21$, total $df = 20$) (Fig. 2B) and with inhibitory synapses even more, from $0.001 \pm 0.001/\mu\text{m}^2$ in the control animals to $0.009 \pm 0.005/\mu\text{m}^2$ (one-way ANOVA, $p < 0.0001$, $F = 17.10$, $N = 21$, total $df = 20$) in the conditioned ones (Fig. 2C).

Density of polyribosomes located in dendritic shafts. *Polyribosomes associated with synapses:* The conditioning induced a significant increase in the density of polyribosomes associated with synapses and located in dendritic shafts (control




Figure 4. Location of polyribosomes in heads/necks/bases of dendritic spines. Conditioning and pseudoconditioning do not change the location of polyribosomes in the dendritic spines. **A.** Electron micrograph of B2 barrel hollow showing polyribosomes (arrow) located in the head of single-synapse spine (s). **B.** Electron micrograph of B2 barrel hollow showing polyribosomes (arrows) located in the neck of single-synapse spine (s). Scale bar: 0.5 μm . **C.** Location of polyribosomes in dendritic spines in: control, conditioned (CS+UCS) and pseudoconditioned (PSEUDO) group. Graph shows means \pm SD.

doi:10.1371/journal.pone.0054301.g004

group: $0.006 \pm 0.004/\mu\text{m}^2$; conditioned group: $0.029 \pm 0.016/\mu\text{m}^2$; Kruskal-Wallis test, $p = 0.0026$, $H = 11.89$, $N = 21$, total $df = 20$ (Fig. 3A). An increase was observed in the density of polyribosomes associated with presumptive inhibitory synapses (control group: $0.002 \pm 0.003/\mu\text{m}^2$; conditioned group: $0.015 \pm 0.008/\mu\text{m}^2$; one-way ANOVA, $p = 0.0004$, $F = 12.48$, $N = 21$, total $df = 20$ (Fig. 3B). The density of polyribosomes associated with presumptive excitatory synapses did not significantly

change either in the conditioned or in the pseudoconditioned animals, as compared with the control animals (control group: $0.005 \pm 0.003/\mu\text{m}^2$; conditioned group: $0.011 \pm 0.010/\mu\text{m}^2$; pseudoconditioned group: $0.003 \pm 0.004/\mu\text{m}^2$; Kruskal-Wallis test, $N = 21$, total $df = 20$ (Fig. 3C).

Polyribosomes not associated with synapses: In the conditioned group, the mean density of polyribosomes not associated with synapses did not change ($0.068 \pm 0.021/\mu\text{m}^2$), but in the pseudoconditioned group it was by $\sim 35\%$ lower ($0.049 \pm 0.014/\mu\text{m}^2$) than in the control group ($0.076 \pm 0.009/\mu\text{m}^2$, one-way ANOVA, $p = 0.0119$, $F = 5.719$, $N = 21$, total $df = 20$) (Fig. 3D).

Location of polyribosomes in dendritic spines

Using 3D reconstructions of dendritic spines, we examined the effect of conditioning on distribution of polyribosomes in the spines. Polyribosomes were located in the head, neck or base of the dendritic spine. The base of spine was defined as the area within 150 nm of the neck's origin [12].

Conditioning and pseudoconditioning did not significantly change the location of polyribosomes, as compared to that found in the control animals (chi square test, $\chi^2 = 4.90$, $N = 3$, $df = 2$) (Fig. 4).

Area of postsynaptic density (PSD)

Synapse-associated polyribosomes which increase in density after conditioning might produce proteins necessary for the enhancement of the synaptic response and increase in the synapse size. To test this hypothesis, we measured the area of postsynaptic density (PSD) of both, excitatory and inhibitory synapses. Since the thickness of PSD is quite uniform in the given synapse type, PSD area reflects also the synapse size. We observed a significant difference between the synapses on spines with and without polyribosomes (two-way ANOVA, $N = 21$) (Fig. 5). The synapses on the spines without polyribosomes had the same PSD area in the control (excitatory synapses: $0.066 \pm 0.021 \mu\text{m}^2$, inhibitory synapses: $0.015 \pm 0.007 \mu\text{m}^2$), in the conditioned animals (excitatory synapses: $0.065 \pm 0.021 \mu\text{m}^2$; inhibitory synapses: $0.019 \pm 0.007 \mu\text{m}^2$; two-way ANOVA, $t = 0.0845$, $p > 0.05$, and $t = 0.4048$, $p > 0.05$ for the two synapse types, respectively) and in the pseudoconditioned animals (excitatory synapses: $0.048 \pm 0.021 \mu\text{m}^2$; inhibitory synapses: $0.029 \pm 0.007 \mu\text{m}^2$; two-




Figure 5. Postsynaptic density area. A. Excitatory synapses located on spines containing polyribosomes have larger postsynaptic density area than those on spines without polyribosomes (two-way ANOVA, $p < 0.0001$) and the synapses enlarge during conditioning (ANOVA, $p < 0.01$). Synapses on spines without polyribosomes do not significantly differ in size in the control, in the conditioned group (CS+UCS) and in the pseudoconditioned group (PSEUDO). Graph shows means \pm SD (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$). **B.** Inhibitory synapses located on spines containing polyribosomes have larger postsynaptic density area of than those located on spines without polyribosomes (two-way ANOVA, $p < 0.0001$) and the synapses enlarge during conditioning (ANOVA, $p < 0.05$). Synapses on spines without polyribosomes do not significantly differ in size in the control, in the conditioned group (CS+UCS) and in the pseudoconditioned group (PSEUDO).

doi:10.1371/journal.pone.0054301.g005

way ANOVA, $t = 0.0480$, $p > 0.05$ and $t = 1.346$, $p > 0.05$ for the two synapse types, respectively).

The synapses associated with polyribosomes in all groups had significantly larger PSD area, as compared to those not associated with polyribosomes ($F = 72.29$, $p < 0.0001$ and $F = 21.14$, $p < 0.0001$ for excitatory and inhibitory synapses, respectively). After conditioning PSD area of both excitatory and inhibitory synapses on spines with polyribosomes significantly increased (to $0.141 \pm 0.016 \mu\text{m}^2$ and to $0.061 \pm 0.029 \mu\text{m}^2$, respectively; two-way ANOVA, $t = 3.384$, $p < 0.01$ and $t = 2.665$, $p < 0.05$, respectively).

Discussion

We have found that associative fear learning involving stimulation of whiskers significantly increases the density of polyribosomes associated with inhibitory and excitatory synapses on the dendritic spines and only with inhibitory synapses on the dendritic shafts of the barrels receiving the conditioned stimulus. Interestingly, conditioning did not affect the density of polyribosomes not associated with synapses in the dendritic shafts, suggesting a link between local protein synthesis and synaptic activity.

We used the pseudoconditioned group to test whether our finding was directly associated with the influence of fear learning or whether it resulted only from the application of two kinds of sensory stimuli. From our previous studies we know that pseudoconditioning does not change the functional cortical representation of the “trained” row of whiskers [16]. It also does not bring about the behavioral “minifreezing” effect, an indicator of CS-UCS association. The number of polyribosomes in the neuropil of the barrels of the pseudoconditioned animals was not significantly different when compared with the control group. Thus, the observed changes in the number of polyribosomes are linked to plastic modifications of the barrel cortex induced by conditioning.

In the previous study, we observed the formation of new symmetric (inhibitory) synapses at dendrites during conditioning. An increased concentration of GABA was found in the presynaptic terminals of these synapses. In addition, the number of double-synapse spines was higher after the learning procedure [20]. Therefore it seems likely that the polyribosomes located in double-synapse spines are associated with the experience-dependent formation of new inhibitory synapse at the spines. There is no data available in the literature on local translation of specific proteins of GABA-ergic inhibitory synapses (GABA receptor subunits, gephyrin, GIRK2 channel) or on the presence of mRNAs for these proteins in dendrites. Interestingly, mRNA for GABA α receptor delta subunit was found in dendrites and its content was increased upon mGluR stimulation [30]. Also, Racca et al. [31] reported that mRNA for GIRK2, inwardly rectifying potassium channel that mediates the inhibitory action of GABA β receptor, colocalized with Nova splicing proteins at the synapse. Hence, the increase in the size of inhibitory synapses located on spines containing polyribosomes in the conditioned animals suggests the involvement of polyribosomes in both synaptogenesis and potentiation of the synapses [12].

Since we have not observed an increase in the density of single-synapse spines after fear learning [20], the polyribosomes found in single-synapse spines are most likely involved in the modification of the existing excitatory synapses [14] rather than in building new ones. We found that the area of PSD in single-synapse spines was enlarged after conditioning. On the postsynaptic side, the size of the PSD is proportional to both the total number of ionotropic

glutamate receptors and the AMPA to NMDA receptor ratio [32,33,34]. The coincident increase in polyribosome number and enlargement of PSD in the single-synapse spines suggests a potential involvement of local protein synthesis in enhancement of the strength of the excitatory synapses at the spines [23,24].

A large pool of mRNAs was found in dendrites [35,36,37]. Some of the proteins encoded by these mRNAs could be synthesized in response to stimulation [36,38]. They include a variety of different classes and some of them have been identified, such as the α subunit of calcium/calmodulin – dependent protein kinase II (CaMKII) [36,39]. Some of the newly produced proteins may serve to insert glutamate receptors and rebuild the postsynaptic density scaffolding, to reorganize the chemoarchitecture of the potentiated synapses [23,40]. Our previous studies provided some evidence for the changes of the glutamate receptors in our experimental model. In layer IV of the barrel cortex, within the plastic cortical representation of the whisker row used in the training, we observed an increase in expression of NMDA receptor NR2A subunit mRNA and protein [22]. We also found an increase in the level of PSD95, a locally synthesized scaffolding molecule [41,42], in enriched postsynaptic membrane fraction of the barrel cortex [43]. PSD95 molecules are known to diffuse from spine to spine in the developing neocortex [44]. The conditioning-induced up-regulation of PSD95 can initially result from its transfer to potentiated spines from a diffuse cytoplasmic pool [45] but the long-term persistence of a higher PSD95 level suggests a subsequent local translation of PSD95 mRNA.

In conclusion, this study provides the first evidence that fear learning, which modifies cortical whisker representations, results in mobilization of organelles responsible for local protein synthesis in both inhibitory and excitatory synapses in the layer IV of the barrel cortex. This effect is accompanied by increased inhibitory synaptogenesis and enlargement of PSD area of polyribosome-associated excitatory and inhibitory synapses on spines. Remodeling of cortical circuits by excitatory and inhibitory interactions was observed at the electrophysiological level in other experimental models of learning, in which the neuronal excitability was found to be increased together with an enhancement of inhibitory effects [46,47,48,49]. Similarly, in our experimental model, we have demonstrated that conditioning is followed not only by an increase in inhibitory efficiency [20,21] but also by an enhanced responsiveness of the excitatory neurons [50]. Enhanced inhibition in layer IV produces feedforward inhibition which sharpens the tuning of excitatory barrel neurons [51,52]. Mutual regulation of excitation and inhibition may prevent neurons in the cortical representation of the trained whiskers to become hypo- or hyperactive and to maintain the ability to produce action potentials in response to fluctuating inputs. It can also contribute to an increase in the selectivity to the “trained” input within the conditioned cortical representation and lead to improved perception of stimuli of a novel biological significance.

Our results show some elements of a mechanism responsible for an adjustment of excitatory-inhibitory balance during fear learning-dependent plasticity in the barrel cortex and present yet more evidence that selective regulation of dendritic spines is central to the mechanisms underlying the plasticity of the neuronal circuits.

Author Contributions

Conceived and designed the experiments: MJ MK. Performed the experiments: MJ ES. Analyzed the data: MJ EJ. Wrote the paper: MJ JAL EP MK. Participated in data interpretation: EP JAL. Revised the manuscript critically for important intellectual content: ES EJ.

References

- Bailey CH, Chen M (1988) Long-term sensitization in *Aplysia* increases the number of presynaptic contacts onto the identified gill motor neuron L7. *Proc Natl Acad Sci USA* 85: 9356–9359.
- Black JE, Isaacs KR, Anderson BJ, Alcantara AA, Greenough WT (1990) Learning causes synaptogenesis, whereas motor activity causes angiogenesis, in cerebellar cortex of adult rats. *Proc Natl Acad Sci USA* 87: 5568–5572.
- Kleim JA, Lussnig E, Schwarz ER, Comery TA, Greenough WT (1996) Synaptogenesis and Fos expression in the motor cortex of the adult rat after motor skill learning. *J Neurosci* 16: 4529–4535.
- Hunter A, Stewart MG (1993) Long-term increases in the numerical density of synapses in the chick lobus parolfactorius after passive avoidance training. *Brain Res* 605: 251–255.
- Knafo S, Ariav G, Barkai E, Libersat F (2004) Olfactory learning-induced increase in spine density along the apical dendrites of CA1 hippocampal neurons. *Hippocampus* 14: 819–825.
- Xu T, Yu X, Perlik AJ, Tobin WF, Zweig JA, et al. (2009) Rapid formation and selective stabilization of synapses for enduring motor memories. *Nature* 462: 915–919.
- Jones MS, Barth DS (1997) Sensory-evoked high-frequency (γ -band) oscillating potentials in somatosensory cortex of the unanesthetized rat. *Brain Res* 768: 167–176.
- Knott GW, Quairiaux C, Genoud C, Welker E (2002) Formation of dendritic spines with GABAergic synapses induced by whisker stimulation in adult mice. *Neuron* 34: 265–273.
- Majewska AK, Newton JR, Sur M (2006) Remodeling of synaptic structure in sensory cortical areas in vivo. *J Neurosci* 26: 3021–3029.
- Holtmaat A, De Paola V, Wilbrecht L, Knott GW (2008) Imaging of experience-dependent structural plasticity in the mouse neocortex in vivo. *Behav Brain Res* 192: 20–25.
- Yu H, Majewska AK, Sur M (2011) Rapid experience-dependent plasticity of synapse function and structure in ferret visual cortex in vivo. *Proc Natl Acad Sci USA* 108: 21235–21240.
- Ostroff LE, Cain CK, Bedont J, Monfils MH, Ledoux JE (2010) Fear and safety learning differentially affect synapse size and dendritic translation in the lateral amygdala. *Proc Natl Acad Sci USA* 107: 9418–9423.
- Lushnikova I, Skibo G, Muller D, Nikonenko I (2011) Excitatory synaptic activity is associated with a rapid structural plasticity of inhibitory synapses on hippocampal CA1 pyramidal cells. *Neuropharmacology* 60: 757–764.
- Bourne JN, Harris KM (2011) Coordination of size and number of excitatory and inhibitory synapses results in a balanced structural plasticity along mature hippocampal CA1 dendrites during LTP. *Hippocampus* 21: 354–373.
- Woolsey TA, Van Der Loos, H (1970) The structural organization of layer IV in the somatosensory region (SI) of mouse cerebral cortex: The description of a cortical field composed of discrete cytoarchitectonic units. *Brain Res* 17: 205–242.
- Siucinska E, Kossut M (1996) Short-lasting classical conditioning induces reversible changes of representational maps of whiskers in mouse SI cortex – a 2DG study. *Cereb Cortex* 6: 506–513.
- Cybulska-Klosowicz A, Zakrzewska R, Kossut M (2009) Brain activation patterns during classical conditioning with appetitive or aversive UCS. *Behav Brain Res* 204: 102–111.
- Siucinska E, Kossut M, Stewart MG (1999) GABA immunoreactivity in mouse barrel field after aversive and appetitive classical conditioning training involving facial vibrissae. *Brain Res* 843: 62–70.
- Gierdalski M, Jablonska B, Siucinska E, Lech M, Skibinska A, et al. (2001) Rapid regulation of GAD67 mRNA and protein level in cortical neurons after sensory learning. *Cereb Cortex* 11: 806–815.
- Jasinska M, Siucinska E, Cybulska-Klosowicz A, Pyza E, Furness DN, et al. (2010) Rapid, learning-induced inhibitory synaptogenesis in murine barrel field. *J Neurosci* 30: 1176–1184.
- Tokarski K, Urban-Ciecko J, Kossut M, Hess G (2007) Sensory learning-induced enhancement of inhibitory synaptic transmission in the barrel cortex of the mouse. *Eur J Neurosci* 26: 134–141.
- Skibinska A, Lech M, Kossut M (2005) Differential regulation of cortical NMDA receptor subunits by sensory learning. *Brain Res* 1065: 26–36.
- Ostroff LE, Fiala JC, Allwardt B, Harris KM (2002) Polyribosomes redistribute from dendritic shafts into spines with enlarged synapses during LTP in developing rat hippocampal slices. *Neuron* 35: 535–545.
- Bourne JN, Sorra KE, Hurlburt J, Harris KM (2007) Polyribosomes are increased in spines of CA1 dendrites 2 h after the induction of LTP in mature rat hippocampal slices. *Hippocampus* 17: 1–4.
- Steward O, Schuman EM (2001) Protein synthesis at synaptic sites on dendrites. *Annu Rev Neurosci* 24: 299–325.
- Witcher MR, Park YD, Lee MR, Sharma S, Harris KM, et al. (2010) Three-dimensional relationships between perisynaptic astroglia and human hippocampal synapses. *Glia* 58: 572–587.
- Steward O, Ribak CE (1986) Polyribosomes associated with synaptic specializations on axon initial segments: localization of protein-synthetic machinery at inhibitory synapses. *J Neurosci* 6: 3079–3085.
- Fiala JC, Harris KM (2001) Extending unbiased stereology of brain ultrastructure to three-dimensional volumes. *J Am Med Inform Assoc* 8: 1–16.
- Jasińska M, Siucińska E, Glazewski S, Pyza E, Kossut M (2006) Characterization and plasticity of the double synapse spines in the barrel cortex of the mouse. *Acta Neurobiol Exp (Wars)* 66: 99–104.
- Dictenberg JB, Swanger SA, Antar LN, Singer RH, Bassell GJ (2008) A direct role for FMRP in activity-dependent dendritic mRNA transport links filopodial-spine morphogenesis to fragile X syndrome. *Dev Cell* 14: 926–939.
- Racca C, Gardiol A, Eom T, Ule J, Triller A, et al. (2010) The Neuronal Splicing Factor Nova Co-Localizes with Target RNAs in the Dendrite. *Front Neural Circuits* 4: 5.
- Takumi Y, Ramirez-León V, Laake P, Rinvik E, Ottersen OP (1999) Different modes of expression of AMPA and NMDA receptors in hippocampal synapses. *Nat Neurosci* 2: 618–624.
- Ganeshina O, Berry RW, Petralia RS, Nicholson DA, Geinisman Y (2004) Synapses with a segmented, completely partitioned postsynaptic density express more AMPA receptors than other axospinous synaptic junctions. *Neuroscience* 125: 615–623.
- Ganeshina O, Berry RW, Petralia RS, Nicholson DA, Geinisman Y (2004) Differences in the expression of AMPA and NMDA receptors between axospinous perforated and nonperforated synapses are related to the configuration and size of postsynaptic densities. *J Comp Neurol* 468: 86–95.
- Steward O, Levy WB (1982) Preferential localization of polyribosomes under the base of dendritic spines in granule cells of the dentate gyrus. *J Neurosci* 2: 284–291.
- Steward O, Schuman EM (2003) Compartmentalized synthesis and degradation of proteins in neurons. *Neuron* 40: 347–359.
- Cajigas JJ, Tushev G, Will TJ, Dieck S, Fuerst N, et al. (2012) The local transcriptome in the synaptic neuropil revealed by deep sequencing and high-resolution imaging. *Neuron* 74: 453–66.
- Martin KC, Zukin RS (2006) RNA trafficking and local protein synthesis in dendrites: an overview. *J Neurosci* 26: 7131–7134.
- Scheetz AJ, Nairn AC, Constantine-Paton M (2000) NMDA receptor-mediated control of protein synthesis at developing synapses. *Nat Neurosci* 3: 211–216.
- Wells DG, Richter JD, Fallon JR (2000) Molecular mechanisms for activity-regulated protein synthesis in the synapto-dendritic compartment. *Curr Opin Neurobiol* 10: 132–137.
- Todd PK, Mack KJ, Malter JS (2003) The fragile X mental retardation protein is required for type-I metabotropic glutamate receptor-dependent translation of PSD-95. *Proc Natl Acad Sci U S A* 100: 14374–14378.
- Lee CC, Huang CC, Wu MY, Hsu KS (2005) Insulin stimulates postsynaptic density-95 protein translation via the phosphoinositide 3-kinase-Akt-mammalian target of rapamycin signaling pathway. *J Biol Chem* 280: 18543–18550.
- Skibinska A, Lech M, Kossut M (2001) PSD95 protein level rises in murine somatosensory cortex after sensory training. *Neuroreport* 12: 2907–2910.
- Gray NW, Weimer RM, Bureau I, Svoboda K (2006) Rapid redistribution of synaptic PSD-95 in the neocortex in vivo. *PLoS Biol* 4: e370.
- Li Z, Sheng M (2003) Some assembly required: the development of neuronal synapses. *Nat Rev Mol Cell Biol* 4: 833–41.
- Froemke RC, Merzenich MM, Schreiner CE (2007) A synaptic memory trace for cortical receptive field plasticity. *Nature* 450: 425–429.
- Galindo-Leon EE, Lin FG, Liu RC (2009) Inhibitory plasticity in a lateral band improves cortical detection of natural vocalizations. *Neuron* 62: 705–716.
- Brosh I, Barkai E (2009) Learning-induced enhancement of feedback inhibitory synaptic transmission. *Learn Mem* 16: 413–416.
- Saar D, Reuveni I, Barkai E (2011) Mechanisms underlying rule learning-induced enhancement of excitatory and inhibitory transmission. *J Neurophysiol* 107: 1222–1229.
- Bekisz M, Garkun Y, Wabno J, Hess G, Wrobel A, et al. (2001) Increased excitability of cortical neurons induced by associative learning: an ex vivo study. *Eur J Neurosci* 32: 1715–1725.
- Sun QQ, Huganard JR, Poirine DA (2006) Barrel cortex microcircuits: thalamic inhibition in spiny stellate cells is mediated by a small number of fast-spiking neurons. *J Neurosci* 26: 1219–1230.
- Miller KD, Pinto DJ, Simons DJ (2001) Processing in layer 4 of the neocortical circuit: New insights from visual and somatosensory cortex. *Curr Opin Neurobiol* 11: 488–497.