

A role for *GLABRA1* in dark-induced senescence*

Aleksandra Eckstein^{1,2}, Joanna Grzyb³, Paweł Hermanowicz⁴, Justyna Łabuz⁴
and Agnieszka Katarzyna Banaś¹✉

¹Department of Plant Biotechnology, Faculty of Biochemistry, Biophysics and Biotechnology, Jagiellonian University, Kraków, Poland; ²Department of Plant Physiology and Biotechnology, Faculty of Biology, University of Gdańsk, Gdańsk, Poland; ³Department of Biophysics, Faculty of Biotechnology, University of Wrocław, Wrocław, Poland; ⁴Laboratory of Photobiology, The Malopolska Centre of Biotechnology, Jagiellonian University, Kraków, Poland

The *GLABRA (GL1)* gene, belonging to the transcription factor-encoding *myb* gene family, is responsible for trichome formation in *Arabidopsis thaliana* (L.) Heynh. The leaves and stems of *glabra1* mutant plants are devoid of trichomes. Having an easily observable phenotype, the *gl1* mutation was one of the first markers established for genetic mapping of *Arabidopsis thaliana*. Since then, the *GL1* gene has been assigned roles in other processes, also related to leaf structure. In this study we present some previously undescribed effects of the *gl1* mutation on dark-induced senescence. This process was induced by covering selected mature leaves of Columbia wild-type and *gl1* *Arabidopsis* with black paper for 4 days, while the plants remained growing in a normal photoperiod. While no visible differences in the external symptoms of senescence could be observed in the darkened leaves, the expression of senescence-associated genes was significantly lower in *gl1* plants as compared to the wild type. The darkening of leaves led to a decrease in photosynthetic activity and the expression of photosynthesis-associated genes, in comparison to the control leaves. This effect was much less pronounced in *gl1* than in the wild type plants. Therefore, *gl1* plants seem to be less susceptible to dark-induced aging, suggesting a possible role for the *GL1* gene in controlling the onset and progress of senescence. This result is also of practical importance, since *gl1* is the genetic background of many other mutants. It may therefore be advisable to revise some of the results obtained with such mutants in light of findings presented here.

Key words: *GLABRA1*, senescence, darkness, photosynthesis, photosynthetic pigments, trichomes

Received: 11 April, 2019; revised: 23 May, 2019; accepted: 29 May, 2019; available on-line: 30 June, 2019

✉e-mail: a_katarzyna.banas@uj.edu.pl

Acknowledgements of Financial Support: This research was supported by the Polish National Science Centre, grant no. UMO-2011/03/D/NZ3/00210.

*A preliminary report on this subject was presented at the 46th Winter School of the Faculty of Biochemistry, Biophysics and Biotechnology, Jagiellonian University in Krakow, Zakopane, Poland, February 11th–15th 2019

Abbreviations: CAB, CHLOROPHYLL A/B BINDING PROTEIN; *gl1*, *glabra1*; F_v/F_m , maximum quantum yield of photosystem II; PAL1, PHENYLALANINE AMMONIA-LYASE 1; PSI/PSII, photosystem I/II; *RbcS1A*, RIBULOSE BIPHOSPHATE CARBOXYLASE SMALL CHAIN 1A, *SAG*, SENESCENCE ASSOCIATED GENE; *SEN1*, SENESCENCE1; WT, wild type

INTRODUCTION

Arabidopsis thaliana (L.) Heynh trichomes, the so called leaf hairs, are branched, non-glandular structures originating from single epidermal cells. Their formation is controlled by a complex interplay of several transcription factors, belonging to different families. *GLABRA (GL1)* was one of the first genes identified as being responsible for trichome development (Herman & Marks, 1989; Marks & Feldmann, 1989; Oppenheimer *et al.*, 1991). It encodes an R2R3 MYB transcription factor, which along with one of the two redundantly acting basic helix-loop-helix (bHLH) transcription factors *GLABRA3 (GL3)* or *ENHANCER OF GLABRA3 (EGL3)*, and *TRANSPARENT TESTA GLABRA1 (TTG1, WD40 family)*, forms the trichome activator complex (Hauser, 2014; Dai *et al.*, 2016; Matías-Hernández *et al.*, 2016). Activation of this complex (also called MBW: MYB-bHLH-WD40), triggers the trichome formation pathway in a given epidermal cell, while inhibits this process in surrounding cells. This complex induces transcription of *GLABRA2 (GL2)* and other downstream genes required for trichome development (Dai *et al.*, 2016). At the same time it also initiates expression of several R3 MYBs acting as trichome repressors: *CAPRICE (CPC)*, *ENHANCER OF TRIPTYCHON AND CAPRICE1 (ETC1)*, *ETC3* and *TRIPTYCHON (TRY)* (Matías-Hernández *et al.*, 2016). These negative regulators move into adjacent cells, where they inhibit the formation of the activator complex by competing with *GL1* for *GL3* or *EGL3* binding. Other R3 MYB trichome repressors are controlled by an independent pathway mediated by *microRNA156* and *SQUAMOSA PROMOTER BINDING PROTEIN LIKE (SPL)*, regulating temporal trichome patterning (Yu *et al.*, 2010). Moreover, the phytohormones: gibberellins and cytokinins, act as positive regulators of trichome formation (Matías-Hernández *et al.*, 2016).

Although they may seem inconspicuous, trichomes have a wide array of important physiological functions. Trichomes increase the light reflectance of leaves, protecting them against excessive heat and photoinhibition, as well as UV-B damage. The presence of trichomes affects such leaf properties as wettability, water repellence and boundary layer thickness, thus regulating transpiration, water uptake and gas exchange (Bickford, 2016). Trichomes also provide protection against biotic stresses, especially those caused by herbivorous insects (Hauser, 2014). The leaves and stems of *glabra1 (gl1)* mutant

plants are devoid of trichomes, while some remain present on the leaf margins. Having an easily observable phenotype and exhibiting no other apparent abnormalities, the natural *gl1* mutation was one of the first markers established for the genetic mapping of *Arabidopsis* in the 1960's (Redei & Hirono, 1964; Koornneeff *et al.*, 1982). Later, the *gl1* mutation was introduced into the widely used Landsberg and Columbia ecotypes. Since then, *gl1* has been used as a genetic background for generating other mutants. Recently, some other effects of the *gl1* mutation, also related to leaf structure, were reported. *gl1* Leaves exhibit a lower stomatal density and a less ordered pattern of stomatal distribution than wild type (WT) (Tsuji & Coe, 2013). The *gl1* mutation has been also found to affect cuticle formation (Xia *et al.*, 2010). The defective structure of *gl1* cuticle was caused by a reduced content of several cuticular wax components. This in turn compromised the systemic acquired resistance (SAR) of *gl1* plants, as shown by their enhanced susceptibility to fungal pathogens.

In this study, we found a novel function of the GL1 transcription factor. The motivation came from some ambiguous results obtained previously using *Arabidopsis* mutants with a *gl1* background. We found differences in the dark-induced chlorophyll degradation between WT Columbia and mutants with a *gl1* background and *gl1* plants (unpublished data). To clarify whether the lack of GLABRA1 is involved in the observed effect, we compared the process of dark-induced senescence in *Arabidopsis thaliana* WT Columbia and the *gl1* mutant. We used an experimental protocol involving the darkening of individual leaves, while the plants remained in a normal photoperiod (Weaver & Amasino, 2001; Keech *et al.*, 2007). This model shares many common features with natural, age-dependent senescence, including leaf yellowing, chlorophyll degradation and induction of senescence associated genes (*SAGs*) (van der Graaff *et al.*, 2006). To assess the progress of senescence we used several well-established markers, such as photosynthetic efficiency and photosynthetic pigment content in the leaves (Weaver & Amasino, 2001; Sztatelman *et al.*, 2015). We also assessed expression of several genes involved in senescence (*SENESCENCE ASSOCIATED GENES – SAG13*, *SAG12* and *SENESCENCE1 – SEN1*), photosynthesis (*RIBULOSE BIPHOSPHATE CARBOXYLASE SMALL CHAIN 1A – RbcS1A* and *CHLOROPHYLL A/B BINDING PROTEINS – CABs*, including *CAB1* and *CAB2*) and anthocyanin biosynthesis (*PHENYLALANINE AMONIA-LYASE 1 – PAL1*). All genes selected for investigation in this study have been previously used in our work on darkening of detached *Arabidopsis* leaves (Sztatelman *et al.*, 2015).

MATERIALS AND METHODS

Plant material. *Arabidopsis thaliana* WT Columbia-0 and *gl1* (CS28174) seeds were purchased from the Nottingham *Arabidopsis* Stock Center (NASC, Nottingham, UK). After sowing in Jiffy-7 pots (Jiffy International AS), the seeds were stratified at 4°C for 2 days. Plants were cultured in a Sanyo MLR 350 H growth chamber, at 23°C, 80% humidity, in a 14 h light/10 h dark photoperiod and illuminated with fluorescent lamps (OSRAM L36W/77 and PHILIPS Master TL-D 36/W/840) with an average photosynthetic photon flux density of 70 $\mu\text{mol m}^{-2}\text{s}^{-1}$. Plants used for experiments were 5–6 weeks old.

Light/dark treatment. Two or three adult, healthy leaves from each plant were darkened by wrapping in black paper. The plants remained in the growth chamber and after 4 days the darkened and non-darkened leaves were collected from each plant. Both darkened and non-darkened samples were chosen from leaves number 5 to 10. Whole plants were dark-adapted for 16 h before leaf collection (starting with the onset of darkness in the growth chamber). The harvested leaves were either directly used for chlorophyll fluorescence measurements or frozen in liquid nitrogen and stored at –80°C for further experiments (pigment extraction, RNA and protein isolation). Frozen samples consisted of 1–3 leaves (*gl1*) or 3 leaves (WT) pooled from different plants.

Pigment extraction/HPLC measurement. Leaf samples (50–200 mg of fresh mass) were ground in a mortar in 1–2 ml of methanol with a few grains of CaCO_3 . The homogenate was collected quantitatively and centrifuged, and the extraction of the pellet was repeated. Supernatants from both extractions were pooled and adjusted to known volume with methanol. Pigment separation, detection and quantitative analysis were performed according to (Sztatelman *et al.*, 2015).

Measurement of chlorophyll fluorescence. The maximum quantum yield of photosystem II (PSII) (F_v/F_m) was measured using an Open FluorCam FC 800-O/1010 imaging fluorometer. Detached leaves (darkened and non-darkened) were placed on water-soaked paper and dark-adapted for 30 min. After recording the basal fluorescence (F_0) for 2 s, samples were illuminated with saturating white light ($2000 \mu\text{mol m}^{-2} \text{s}^{-1}$) for 960 ms. The average F_v/F_m value for each leaf was calculated using the FluorCam7 software.

RNA isolation and real-time PCR. Total RNA was extracted using the Spectrum Plant Total Kit (Sigma-Aldrich) and digested with DNaseI (Sigma-Aldrich) during purification on columns. RNA concentration was measured using a NanoDrop Lite (Thermo Fisher Scientific) spectrophotometer and its integrity was checked by electrophoresis in 1% agarose gel. First-strand cDNA synthesis was performed using the RevertAid M-MuLV Reverse Transcriptase Kit (Thermo Fisher Scientific) with 1000 ng of RNA and oligo(dT)₁₈ primers. Real-time PCR was performed using SYBR Green JumpStart Taq ReadyMix (Sigma Aldrich) and a Rotor-Gene 6000 thermal cycler (Corbett Research, Australia) according to (Łabuz *et al.*, 2012), with the following modifications: reactions were run in triplicates, and *UBC*, *UBQ10* and *SAND* were used as the reference genes. Primer sequences and annealing temperatures were based on (Sztatelman *et al.*, 2015) and are listed in Table S1 (at <https://ojs.ptbioch.edu.pl/index.php/abp/>). cDNA pooled from six different samples (darkened and non-darkened leaves) was used as a reference for calculating relative gene expression levels. Normalization was performed using normalization factors based on reference gene levels, calculated using geNorm v3.4 (Vandesompele *et al.*, 2002).

Protein extraction and Western blot. Leaf samples were homogenized and adjusted to an equal mass. Proteins were extracted as described by Sakamoto and Briggs (Sakamoto & Briggs, 2002). SDS-PAGE was performed on 12% polyacrylamide gels and was followed by a semi-dry protein transfer (BioRad) to a PVDF membrane. The membrane was blocked overnight with 5% (w/v) non-fat powdered milk in phosphate-buffered saline (PBS), 0.5% Tween-20. Primary antibodies, anti PsbaA (D1 protein, Agrisera AS05084A), were applied at a dilution of 1:10000




Figure 1. (a) Maximum quantum yield of PSII (F_v/F_m) and (b) relative PSII D1 protein levels in the control and 4-day-darkened leaves of WT and *gl1* *A. thaliana* plants.

(a) Each bar corresponds to an average of 40–43 biological replicates. Error bars = S.E. Asterisks indicate statistically significant differences of means (P values adjusted with the Hankel method: * $P=0.01$ –0.05; ** $P=0.001$ –0.01, *** $P<0.001$). (b) Each bar corresponds to an average of 6 biological replicates. The D1 protein level in wild type leaves kept in light was set to 1. Error bars = S.E. Asterisks indicate statistically significant differences of means of the log-transformed values (P values adjusted with the Hankel method: * $P=0.01$ –0.05; ** $P=0.001$ –0.01, *** $P<0.001$).

(Agrisera) at room temperature for 1 h. After washing, secondary antibodies (goat anti-rabbit horseradish peroxidase (HRP) conjugated IgG, Agrisera) were applied at a dilution of 1:10 000, at room temperature for 1 h. The signal was detected with Clarity Western ECL Blotting Substrate (Bio-Rad) and the Bio-Spectrum Imaging System (UVP Ultra-Violet Products Ltd). Then, the membranes were stripped with Restore Plus Western Blot Stripping Buffer (Thermo Scientific) and probed again with anti-actin antibody (AS132640, Agrisera) at a dilution of 1:2500, at room temperature for 1 h, followed by incubation with secondary antibodies and ECL detection. Densitometric quantification was performed using ImageJ. Intensities of the chemiluminescent signals were normalized to actin levels in each sample.

Statistical analysis. The significance of the effects of the plant line (WT, *gl1*) and light conditions (control, darkened) was calculated with two-way ANOVA using the R software. The significance of the four pairwise differences of means (WT control *vs* WT darkened, *gl1* control *vs* *gl1* darkened, WT control *vs* *gl1* control and WT darkened *vs* *gl1* darkened) was tested using the *glbt* command of the *multcomp* package. Holm's correction was used to adjust the p -values for multiple comparisons. In general, variance was not uniform across groups. Thus, the heteroskedasticity-consistent variance – covariance matrix was used in the pairwise comparison procedure, calculated using the sandwich library. The results of the expression ex-

periments (mRNA and protein levels) were log – transformed before statistical analysis.

RESULTS

Photosynthetic efficiency

Photosynthetic efficiency, measured as F_v/F_m , was comparable in WT and *gl1* plants grown under the control conditions. In the darkened leaves of both genotypes, the F_v/F_m parameter values were significantly lower. However, the decrease in F_v/F_m was smaller in *gl1* and the difference between both genotypes was statistically significant at the level of 0.001 (Fig. 1a). Additionally, we assessed the levels of the photosynthetic D1 protein, a core subunit of PSII (Fig. 1b). In control plants the amount of D1 was visibly higher in *gl1* than in WT. After leaf darkening the amount of D1 increased in the WT plants and it slightly decreased in *gl1*, resulting in equal amounts of protein in both genotypes. However, all observed differences in the D1 protein levels between WT and *gl1* plants were statistically non-significant (Fig. 1b).

Photosynthetic pigments

After four days of darkening, the leaves of both, the WT and *gl1*, showed partial yellowing indicating a change in photosynthetic pigment composition (Fig. S1 at <https://ojs.ptbioch.edu.pl/index.php/abp/>). The contents of all analyzed photosynthetic pigments (chlorophyll a and b, neoxanthin, violaxanthin, β -carotene and lutein) per g of fresh weight were comparable in the WT and *gl1* light-grown plants. After leaf darkening, the content of all pigments per fresh weight was visibly reduced, especially in the case of both chlorophylls and β -carotene (by up to 50%) (Fig. 2a–c). This decrease was found to be statistically significant in the case of almost all pigments (apart from violaxanthin) in the WT plants (Fig. 2b). Although in *gl1* plants the differences between control and darkened leaves were not statistically significant, the decrease in pigment content was also visible, suggesting the same process occurred as in the WT plants (Fig. 2). The seemingly minor differences between both genotypes were more pronounced when the pigment content was expressed per amount of total chlorophyll (Fig. S2b–c <https://ojs.ptbioch.edu.pl/index.php/>




Figure 2. Photosynthetic pigment content: (a) chlorophyll a and chlorophyll b, (b) neoxanthin and violaxanthin, (c) lutein and β -carotene, in the control and 4-day-darkened leaves of WT and *gl1* *A. thaliana* plants.

Each bar corresponds to an average of 4 biological replicates. Asterisks indicate statistically significant differences of means (P values adjusted with the Hankel method: * $P=0.01$ –0.05, ** $P=0.001$ –0.01, *** $P<0.001$).




Figure 3. Relative expression of genes associated with (a) senescence (*SAG13*, *SAG12*, *SEN1*), (b) photosynthesis (*RbcS1A*, *CABs*) and anthocyanin synthesis (*PAL1*), at the mRNA level in the control and 4-day-darkened leaves of WT and *gl1* *A. thaliana* plants. Each bar corresponds to an average of 12 biological replicates. Asterisks indicate statistically significant differences of means of the log-transformed values (P values adjusted with the Hankel method: * $P=0.01$ – 0.05 ; ** $P=0.001$ – 0.01 , *** $P<0.001$).

abp/). In the light-grown WT and *gl1* plants all pigment content ratios were comparable (Fig. S2a–c at <https://ojs.ptbioch.edu.pl/index.php/abp/>). After darkening, the content of carotenoids (neoxanthin, violaxanthin and lutein) in relation to chlorophyll had increased, while the β -carotene:chlorophyll ratio remained unchanged (Fig. S2 <https://ojs.ptbioch.edu.pl/index.php/abp/>). The dark-induced increase in carotenoid:chlorophyll ratios was slightly higher in the WT plants than in *gl1* (Fig. S2b–c at <https://ojs.ptbioch.edu.pl/index.php/abp/>). In the case of violaxanthin and lutein, the differences between the control and darkened plants were statistically significant for both, the WT and *gl1* (Fig. S2b–cat at <https://ojs.ptbioch.edu.pl/index.php/abp/>).

Expression of genes involved in photosynthesis and senescence

To check whether the observed changes in pigment content and photosynthetic efficiency were associated with the dark-induced process of senescence, the expression of three senescence marker genes was assessed: *SAG13*, *SAG12* and *SEN1* (Oh *et al.*, 1996; Weaver *et al.*, 1998) (Fig. 3a). In control plants only *SAG13* and *SEN1* were expressed, the transcript levels being similar in the WT and *gl1* plants (Fig. 3a). After leaf darkening, both genes were strongly upregulated (12- and 15-fold respectively) in the WT plants, whereas in *gl1* only a 2-fold increase in *SAG13* expression was observed (Fig. 3a). Dark treatment also induced the expression of *SAG12*, with an almost 6-fold higher transcript level in WT as compared to *gl1* (Fig. 3a). In the case of *SAG13* and *SAG12*, expression levels in the darkened WT leaves were significantly higher than in *gl1*.

We also tested the expression of genes associated with photosynthesis (*RbcS1A*, *CABs*) or known to be induced by light (*PAL1*) (Jones, 1984; Kimura *et al.*, 2003) (Fig. 3b). In the light-grown plants the levels of *RbcS1A* and *CABs* transcripts were slightly lower in *gl1* than in the WT, and the expression of *PAL1* was comparable in both genotypes (Fig. 3b). Leaf darkening caused the downregulation of all three genes. In most cases the effect was statistically significant, except for *PAL1* expression in *gl1* where the decrease was relatively small (Fig. 3b). The dark-induced downregulation of the tested genes was visibly stronger in the WT plants, the effect being statistically significant in the case of *RbcS1A* (Fig. 3b). Thus, for all three genes, the differences between transcript levels in the

control and darkened plants were greater in WT than in *gl1*.

DISCUSSION

Until recently, physiological and molecular studies of the *A. thaliana glabra1* mutant were restricted to the process of trichome formation and some other effects related to the leaf structure (Xia *et al.*, 2010; Tsuji & Coe, 2013). To our knowledge, other aspects of the mutant's physiology, including the progress of senescence, have not been investigated in more detail to date.

The experimental scheme of darkening individual leaves on growing plants applied here has been established as a good model of natural senescence (Weaver & Amasino, 2001; van der Graaff *et al.*, 2006; Keech *et al.*, 2007). While whole darkened plants seem to enter a 'stand-by' mode, aimed to preserve the activity of crucial metabolic pathways for as long as possible, the darkening of individual leaves triggers the degradation of cellular components and nutrient retrieval (Keech *et al.*, 2007). This is reflected in a rapid decline in photosynthetic activity. Similarities between natural and dark-induced senescence in individual leaves have been confirmed by a transcriptome study, showing at the same time that the former process is much more complex (van der Graaff *et al.*, 2006).

WT and *gl1* plants, when grown under control conditions, were comparable by all parameters tested. The only pronounced difference was the lower expression of *RbcS1A* and *CABs* in *gl1* (Fig. 3b), which did not affect the photosynthetic efficiency measured as F_v/F_m (Fig. 1a). This allows us to assume that all the differences observed between both genotypes after leaf darkening are truly linked to differences in the process of senescence. The significantly higher value of the F_v/F_m parameter in the darkened *gl1* leaves as compared to WT was the first hint to a slower progress of senescence in mutant plants. The decline in photosynthetic activity in darkened leaves was not reflected in the levels of the D1 protein which increased in the WT and only slightly decreased in *gl1* (Fig. 1b). Lower photosynthetic efficiency in darkened leaves was correlated with the downregulation of *RbcS1A* and *CABs* photosynthetic genes (Fig. 3b), which is consistent with results of previous similar studies (Weaver & Amasino, 2001; Sztatelman *et al.*, 2015). This suggests that although the biosynthesis of new components has already been inhibited, the degradation of the photosynthetic ma-

chinery might not be significantly advanced yet. Higher *RbcS1A* and *CABs* transcript levels in the darkened *gl1* leaves as compared with the WT ones further point to a slower progress of senescence in these plants. This effect was statistically significant in the case of *RbcS1A*. Degradation of the D1 protein is also known to be induced by abiotic stresses, such as excessive light, UV-B radiation or high salinity (Kato & Sakamoto, 2009; Sakuraba *et al.*, 2014; Sztatelman *et al.*, 2015). The lack of an evident decrease in D1 levels and the absence of degradation products may therefore mean that the applied dark treatment specifically triggers senescence, without being an additional source of stress. The *PAL1* gene is involved in anthocyanin biosynthesis and induced by light (Jones, 1984; Kimura *et al.*, 2003). Downregulation of this gene in darkness probably reflects the reduced need for the production of photoprotective pigments under these conditions. The small decrease in *PAL1* transcript level, observed in the darkened *gl1* leaves, may therefore be another indicator of slower senescence in these plants (Fig. 3b).

Natural senescence is invariably connected with gradual leaf yellowing. This could be also observed in our experimental model (Fig. S1 at <https://ojs.ptbioch.edu.pl/index.php/abp/>). This leaf yellowing is due to the degradation of photosynthetic pigments, which contributes to the decline in photosynthetic activity. This was confirmed by the analysis of photosynthetic pigment content. As expected, the levels of all pigments had decreased substantially after dark treatment (Fig. 2). No major, significant differences could be observed in the reaction to darkening between *gl1* and WT. In both genotypes, an increase in the amount of carotenoids (neoxanthin, violaxanthin and lutein) as compared to total chlorophyll content was observed after leaf darkening, which is consistent with the usually faster degradation of chlorophylls (Young *et al.*, 1991; Lu *et al.*, 2001) (Fig. S2b–c at <https://ojs.ptbioch.edu.pl/index.php/abp/>). This parameter revealed some subtle differences between the WT and *gl1* plants. A slightly higher amount of chlorophylls after darkening in *gl1* leaves as compared to WT (Fig. 2a) is also reflected in lower violaxanthin, neoxanthin and lutein ratio to total chlorophyll in the darkened *gl1* leaves (Fig. S2b–c at <https://ojs.ptbioch.edu.pl/index.php/abp/>). This points to a slower decay of chlorophylls as compared with other photosynthetic pigments in *gl1*, thus supporting the assumption of delayed senescence in this mutant. An increase in the chlorophyll *a*:chlorophyll *b* ratio in the darkened leaves, typical for senescence, was comparable in the WT and *gl1* plants (Fig. S2a at <https://ojs.ptbioch.edu.pl/index.php/abp/>) (Pružinská *et al.*, 2005). It indicates that in both genotypes, in darkness chlorophyll *b* was degraded faster than chlorophyll *a*. As the chlorophyll *a*:chlorophyll *b* ratio reflects the PSI/PSII antennae stoichiometry (Pfannschmidt *et al.*, 2001), we may also postulate that there is no selective degradation of one photosystem in both genotypes tested.

To confirm the observed slower senescence in darkened *gl1* leaves we assessed the expression of three senescence marker genes: *SAG12*, *SAG13* and *SEN1* (Fig. 3a). All of them were upregulated (*SAG13*, *SEN1*) or induced (*SAG12*) during leaf darkening (Oh *et al.*, 1996; Weaver *et al.*, 1998). Low expression of *SAG13* and the absence of *SAG12* transcript in control plants is consistent with the respective roles of these genes as early and late senescence markers (Weaver *et al.*, 1998). *SAG12* is also considered to be the most specific marker for natural senescence, which is in line with the observed expression pattern (transcript detected only in the darkened leaves). The *SEN1* gene, apart from being an

other senescence marker, is also thought to be involved in a direct response to darkness (Oh *et al.*, 1996). A significantly lower expression of all three genes in the darkened *gl1* leaves when compared to WT provides overwhelming evidence for the delayed senescence in this mutant.

The results obtained in this work provide ample and robust evidence that the onset and progress of dark-induced senescence is delayed in the *A. thaliana glabra1* mutant. Taking into account previous reports comparing different models of senescence (Weaver & Amasino, 2001; van der Graaff *et al.*, 2006), it is highly probable that the same holds true for the process of natural senescence in these plants. The lack of trichomes makes the *gl1* mutant more susceptible to numerous environmental stresses, both abiotic and biotic, which may contribute to the induction of senescence in these plants (Hauser, 2014; Bickford, 2016). Nevertheless, trichome production comes at a cost (Sletvold *et al.*, 2010; Hauser, 2014). Although the presence of trichomes may be beneficial to the plant, they are developed at the expense of other traits. As demonstrated in *Arabidopsis lyrata*, trichome density (and the resulting resistance to herbivores) was negatively correlated with plant size and fitness (Sletvold *et al.*, 2010). Perhaps in the case of the *gl1* mutant, the resources which would otherwise be used for trichome production can be allocated to other processes. This could explain a higher PSII efficiency and slower progress of senescence in the darkened *gl1* leaves. On the other hand, taking into account the nature of *GLABRA1* as a transcription factor, its direct role in regulating the expression of genes involved in senescence cannot be excluded. It is also possible that both of the proposed (and perhaps also other) mechanisms contribute to the observed delayed senescence in the *gl1* mutant.

The demonstrated effect of the *gl1* mutation on the process of senescence should be taken into account in any experiments using other mutants with a *gl1* genetic background and the results should be interpreted with caution. The *gl1* example shows that a single mutation can affect very different, seemingly unrelated processes. It is therefore advisable to avoid using such “cryptic” multiple mutants. If this is impossible, an appropriate control, that is the *gl1* mutant, should be used.

Conflict of Interest Disclosures

The authors declare no conflict of interest.

REFERENCES

- Bickford CP (2016) Ecophysiology of leaf trichomes. *Funct Plant Biol* 43: 807–814. <https://doi.org/10.1071/FP16095>
- Dai X, Zhou L, Zhang W, Cai L, Guo H, Tian H, Schiefelbein J, Wang S (2016) A single amino acid substitution in the R3 domain of *GLABRA1* leads to inhibition of trichome formation in *Arabidopsis* without affecting its interaction with *GLABRA3*. *Plant Cell Environ* 39: 897–907. <https://doi.org/10.1111/pce.12695>
- Hauser MT (2014) Molecular basis of natural variation and environmental control of trichome patterning. *Front Plant Sci* 5: 320. <https://doi.org/10.3389/fpls.2014.00320>
- Herman PL, Marks MD (1989) Trichome development in *Arabidopsis thaliana*. II. Isolation and complementation of the *GLABROUS1* gene. *Plant Cell* 1: 1051–1055. <https://doi.org/10.1105/tpc.1.11.1051>
- Jones DH (1984) Phenylalanine ammonia-lyase: regulation of its induction, and its role in plant development. *Phytochemistry* 23: 1349–1359. [https://doi.org/10.1016/S0031-9422\(00\)80465-3](https://doi.org/10.1016/S0031-9422(00)80465-3)
- Kato Y, Sakamoto W (2009) Protein quality control in chloroplasts: a current model of D1 protein degradation in the photosystem II repair cycle. *J Biochem* 146: 463–469. <https://doi.org/10.1093/jb/mvp073>
- Keech O, Pesquet E, Ahad A, Askne A, Nordvall DAG, Vodnala SM, Tuominen H, Hurry V, Dizengremel P, Gardestrom P (2007) The

- different fates of mitochondria and chloroplasts during dark-induced senescence in *Arabidopsis* leaves. *Plant Cell Environ* **30**: 1523–1534. <https://doi.org/10.1111/j.1365-3040.2007.01724.x>
- Kimura M, Yamamoto YY, Seki M, Sakurai T, Sato M, Abe T, Yoshida S, Manabe K, Shinozaki K, Matsui M (2003) Identification of *Arabidopsis* genes regulated by high light-stress using cDNA microarray. *Photochem Photobiol* **77**: 226–233. [https://doi.org/10.1562/0031-8655\(2003\)0770226IOAGRB2.0.CO;2](https://doi.org/10.1562/0031-8655(2003)0770226IOAGRB2.0.CO;2)
- Koornneef M, Dellaert LWM, Van der Veen JH (1982) EMS-and relation-induced mutation frequencies at individual loci in *Arabidopsis thaliana* (L.) Heynh. *Mut Res-Fund Mol M* **93**: 109–123. [https://doi.org/10.1016/0027-5107\(82\)90129-4](https://doi.org/10.1016/0027-5107(82)90129-4)
- Lu C, Lu Q, Zhang J, Kuang T (2001) Characterization of photosynthetic pigment composition, photosystem II photochemistry and thermal energy dissipation during leaf senescence of wheat plants grown in the field. *J Exp Bot*, **52**: 1805–1810. <https://doi.org/10.1093/jxb/52.362.1805>
- Labuz J, Sztatelman O, Banaś AK, Gabrys H (2012) The expression of phototropins in *Arabidopsis* leaves: developmental and light regulation. *J Exp Bot* **63**: 1763–1771. <https://doi.org/10.1093/jxb/ers061>
- Marks MD, Feldmann KA (1989) Trichome development in *Arabidopsis thaliana*. I. T-DNA tagging of the *GLABROUS1* gene. *Plant Cell* **1**: 1043–1050. <https://doi.org/10.1105/tpc.1.11.1043>
- Matías-Hernández L, Aguilar-Jaramillo AE, Cigliano RA, Sanseverino W, Pelaz S (2016) Flowering and trichome development share hormonal and transcription factor regulation. *J Exp Bot* **67**: 1209–1219. <https://doi.org/10.1093/jxb/erv534>
- Oh SA, Lee SY, Chung IK, Lee CH, Nam HG (1996) A senescence-associated gene of *Arabidopsis thaliana* is distinctively regulated during natural and artificially induced leaf senescence. *Plant Mol Biol* **30**: 739–754. <https://doi.org/10.1007/BF00019008>
- Oppenheimer DG, Herman PL, Sivakumaran S, Esch J, Marks MD (1991) A *myb* gene required for leaf trichome differentiation in *Arabidopsis* is expressed in stipules. *Cell* **67**: 483–493. [https://doi.org/10.1016/0092-8674\(91\)90523-2](https://doi.org/10.1016/0092-8674(91)90523-2)
- Pfannschmidt T, Schütze K, Brost M, Oelmüller R (2001) A novel mechanism of nuclear photosynthesis gene regulation by redox signals from the chloroplast during photosystem stoichiometry adjustment. *J Biol Chem* **276**: 36125–36130. <https://doi.org/10.1074/jbc.M105701200>
- Pružinská A, Tanner G, Aubry S, Anders I, Moser S, Müller T, Ongania H-K, Kräutler B, Youn J-Y, Liljegren SJ, Hörtensteiner S (2005) Chlorophyll breakdown in senescent *Arabidopsis* leaves. Characterization of chlorophyll catabolites and of chlorophyll catabolic enzymes involved in the degreening reaction. *Plant Physiol* **139**: 52–63. <https://doi.org/10.1104/pp.105.065870>
- Redei GP, Y Hirono (1964) Linkage studies. *Arab Inf Serv*, **1**: 9–10.
- Sakamoto K, Briggs WR (2002) Cellular and subcellular localization of phototropin 1. *Plant Cell* **14**: 1723–1735. <https://doi.org/10.1105/tpc.003293>
- Sakuraba Y, Lee SH, Kim YS, Park OK, Hörtensteiner S, Paek NC (2014) Delayed degradation of chlorophylls and photosynthetic proteins in *Arabidopsis* autophagy mutants during stress-induced leaf yellowing. *J Exp Bot* **65**: 3915–3925. <https://doi.org/10.1093/jxb/eru008>
- Sletvold N, Huttunen P, Handley R, Kärkkäinen K, Ågren J (2010) Cost of trichome production and resistance to a specialist insect herbivore in *Arabidopsis lyrata*. *Evol Ecol* **24**: 1307–1319. <https://doi.org/10.1007/s10682-010-9381-6>
- Sztatelman O, Grzyb J, Gabrys H, Banaś AK (2015) The effect of UV-B on *Arabidopsis* leaves depends on light conditions after treatment. *BMC Plant Biol* **15**: 281. <https://doi.org/10.1186/s12870-015-0667-2>
- Tsuji J, Coe L (2013) The *glabra1* mutation affects the stomatal patterning of *Arabidopsis thaliana* rosette leaves. *Bios* **84**: 92–98. <https://doi.org/10.1893/0005-3155-84.2.92>
- van der Graaff E, Schwacke R, Schneider A, Desimone M, Flüge U, Kunze R (2006) Transcription analysis of *Arabidopsis* membrane transporters and hormone pathways during developmental and induced leaf senescence. *Plant Physiol* **141**: 776–792. <https://doi.org/10.1104/pp.106.079293>
- Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, De Paep A, Speleman F (2002) Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biol* **3**: research0034. <https://doi.org/10.1186/gb-2002-3-7-research0034>
- Weaver LM, Amasino RM (2001) Senescence is induced in individually darkened *Arabidopsis* leaves, but inhibited in whole darkened plants. *Plant Physiol* **127**: 876–886. <https://doi.org/10.1104/pp.010312>
- Weaver LM, Gan S, Quirino B, Amasino RM (1998) A comparison of the expression patterns of several senescence-associated genes in response to stress and hormone treatment. *Plant Mol Biol* **37**: 455–469. <https://doi.org/10.1023/A:1005934428906>
- Xia Y, Yu K, Navarre D, Seebold K, Kachroo A, Kachroo P (2010) The *glabra1* mutation affects cuticle formation and plant responses to microbes. *Plant Physiol* **154**: 833–846. <https://doi.org/10.1104/pp.110.161646>
- Young AJ, Wellings R, Britton G (1991) The fate of chloroplast pigments during senescence of primary leaves of *Hordeum vulgare* and *Avena sativum*. *J Plant Physiol* **137**: 701–705. [https://doi.org/10.1016/S0176-1617\(11\)81225-3](https://doi.org/10.1016/S0176-1617(11)81225-3)
- Yu N, Cai WJ, Wang S, Shan CM, Wang LJ, Chen XY (2010) Temporal control of trichome distribution by microRNA156-targeted SPL genes in *Arabidopsis thaliana*. *Plant Cell* **22**: 2322–2335. <https://doi.org/10.1105/tpc.109.072579>