

# Mesenchymal Stem Cells in Treatment of Perianal and Rectovaginal Fistulas

## Abstract

Perianal fistulas are still a major challenge in everyday clinical practice. Particularly difficult fistulas, like those in Crohn's Disease, often require multiple aggressive surgical interventions, with high risk of sphincter damage and incontinence. Intensive trials have been conducted for over 10 years on the treatment of difficult fistulas with mesenchymal stem cells (MSC). This paper presents a short review of basic mechanisms of mesenchymal stem cells in healing of damaged tissues and results of major clinical trials. MSC in perianal fistulas, regardless of research protocol has a 70% healing rate in recurrent fistulas and in fistulas in Crohn's Disease. Further research is needed to precisely define the indications for MSC-based treatment of tissue damage, and particularly perianal and recto-vaginal fistulas.

**Keywords:** Fistula-in-ano; Recurrent fistula; Complex fistula; Crohn disease; Adipose-derived mesenchymal stem cells; Clinical trials

## Mini Review

Volume 5 Issue 4 - 2016

**Piotr Wałęga<sup>1\*</sup>, Marcin Piejko<sup>1,2,3</sup>, Michal Romaniszyn<sup>1</sup> and Justyna Drukala<sup>2</sup>**

<sup>1</sup>3<sup>rd</sup> Department of General Surgery, Medical College Jagiellonian University, Poland

<sup>2</sup>Department of Cell Biology, Faculty of Biochemistry, Poland

<sup>3</sup>Department of Analytical Biochemistry, Biophysics and Biotechnology, Jagiellonian University, Poland

**\*Corresponding author:** Piotr Wałęga, 3<sup>rd</sup> Department of General Surgery, Medical College Jagiellonian University, ul. Pradnicka 35-37, 31-202 Krakow, Poland, Tel: +48 12 633 19 95; Fax: +48 12 633 31 05; Email: gabinet.walega@gmail.com

**Received:** July 29, 2016 | **Published:** October 19, 2016

**Abbreviations:** ASC: Adipose-derived mesenchymal Stem Cells; MSC isolated from adipose tissue; bFGF: Basic Fibroblast Growth Factor; MSC: Mesenchymal Stem Cell; HGF: Hepatocyte Growth Factor; ILGF-1: Insulin-Like Growth Factor 1; VEGF: Vascular-Endothelial Growth Factor

## Introduction

Perianal and rectovaginal fistulas are pathological connections between the rectum and adjacent structures in the rectoanal region. About 80% of perianal fistulas are of cryptoglandular origin the remaining 20% develop in Crohn's Disease, after trauma, or have iatrogenic etiology, etc.

Standard in perianal fistula treatment are cutting (fistulotomy, fistulectomy) or non-cutting (seton) surgical procedures, to remove granulation tissue, necrosis and translocated mucosal cells. The success rate of primary surgical treatment of perianal fistulas reaches about 70% [1]. In the remaining cases, there is no healing of the fistula after the procedure, or the fistula reoccurs after initial healing. Fistula recurrence rate in Crohn's Disease is relatively high (34%) [2] which is due to constant inflammation of the intestines and soft tissues. Therefore surgical approach is preceded by immunosuppressive treatment, including anti-TNF $\alpha$ , which is necessary for proper soft tissues healing.

Regardless of the fistula etiology, the disease greatly reduces the patients' quality of life both due to recurrence and incontinence [3]. The metaanalysis by Blumetti et al. [4] shows, that aggressive surgical approach (eg. fistulectomy) has a greater chance of successful healing, at the same time increasing the risk of fecal incontinence. Therefore, there is a constant search for better alternatives for fistula treatment.

In 2003 Prof. Garcia-Olmo made the first attempt to treat a rectovaginal fistula in a 33-year old female patient with Crohn's

Disease, using the mesenchymal stem cells derived from the fatty tissue of the patient (Adipose-derived mesenchymal Stem Cells) [5]. By implantation of about 9mln of autologous stem cells he achieved complete closure and lack of recurrence of the fistula. From that time, many approaches to soft tissue treatment with MSC were made. This paper presents a short review of clinical research on the MSC treatment of perianal and rectovaginal fistulas so far.

## Discussion

### Mechanism of action

Mesenchymal stem cells exist in the perivascular niche of almost all tissues [6], however, the most useful source of these cells are bone marrow, adipose tissue and umbilical cord [7]. In classic approach, MSC are pluripotent cells that support the natural healing process by increasing the overall pool of cells (self-renew) and differentiation in the direction of mesenchymal cells (fibroblasts producing collagen, adipocytes, myocytes, osteocytes, etc.) [8]. In addition, MSC modulate the healing process by stimulating dermal fibroblasts to divide, migrate toward the injury, and the expression of collagens [9]. In hypoxic conditions which occur on the edges of tissue damage, MSC secrete growth factors, such as vascular endothelial growth factor (VEGF), basic fibroblast growth factor (bFGF), hepatocyte growth factor (HGF), insulin-like growth factor, and other [10] therefore inducing revascularisation [11]. Mesenchymal stem cells also possess immunosuppressive properties [12,13]. In a mouse model of inflammatory bowel disease MSC suppress Th1 lymphocytes, which leads to a reduction in clinical and histological symptoms of inflammatory bowel disease [13]. Multifactorial mechanism of action of MSC responds to the need for proper healing of soft tissue, but we still do not know which of the effects described by basic science have clinical significance. What's more, there are no

direct studies on the characteristic of the microenvironment of the damaged tissue, and in particular its “weak spots” in relation to the properties of the MSC, which could allow the development of better treatment strategies.

### Clinical outputs

Since 2005 at least 9 clinical trials with MSC treatment of

the perianal fistula were conducted [14-24]. The summary of prospective studies are shown in Table 1. In all studies, 225 cell-treated patients were completed the protocol, 107 were healed in 8-24 weeks of follow-up. In all trial administration of the cells was combined with surgical closure of the fistula (one-stage procedure). Patients received 9 to 139 million cells suspended in fibrin glue [15,16,23] or in saline solution [15,19-22]. The cells were administered into the walls of the fistula and into its lumen.

**Table 1:** Summary of clinical trials and compassionate-use programmes with mesenchymal stem cells therapy in treatment of fistulas.

	Published Clinical Trial	Etiology of Fistula	Type and Amount of Cells	N / n	H / h
1	Garcia-Olmo et al. [14]	Crohn	Autogenic ASC, 3-30 mln	9	6
2	Garcia-Olmo et al. [23]	both	Autogenic ASC, 20 mln	17 / 7*	12 / 5*
3	Ciccocioppo et al. [18]	Crohn	Autogenic MSC, 54-139 mln	10	7
4	de la Portilla et al. [20]	Crohn	Allogenic ASC, 20-40 mln	16	9
5	Herreros et al. [15]	non-Crohn	Autogenic ASC, 20 mln	124	51
		Results in multicenter trial excluded a pioneering center		101	35
		Results in the pioneering center (Garcia-Olmo)		23	16
6	Lee et al. [16]	Crohn	Autogenic ASC, 9-42 mln	33	27
7	Cho et al. [17]	Crohn	Autogenic ASC, 30-40 mln	9	3
8	Garcia-Olmo et al. [22]**	Both	Auto- autogenic ASC, no data	7 / 3*	4 / 2*
9	Garcia-Arranz et al. [24]	Crohn	Allogenic ASC, 20-40 mln	5	3

N: number of cell-treated patients who completed the protocol, H – number of cell-treated patients with healed fistula in 8-24 weeks follow-up; \* - Number of non-Crohn/Crohn patients; \*\* - Compassionate-use programme.

It is difficult to determine the efficacy of MSC / ASC as each of these studies had a slightly different protocol. For this reason, the total numbers of patients were counted - patients who received mesenchymal stem cells (isolated from bone marrow and adipose tissue, auto and allogeneic, alone or with fibrin glue) in clinical trials or registered treatment programs (compassionate use). Only prospective studies were taken into account, and the endpoint (ie. “healed”) was considered as clinical healing in the period from 8 to 24 weeks after administration of the cells (the status of “partially healed” was classified as “not healed” unless the healing was achieved up to 12 months after the cells were administered).

Among 225 (or 154) patients who received the MSC or ASC status of “healed” were reported in 138 (54%) (or 103 (67%) - excluding non-pioneering center from the largest clinical trial conducted by Herreros et al. [15]) cases. The healing rate in non-Crohn fistulas were 45% (n=148 cell-treated patients) and (or 68% (n=47) excluding non-pioneering centers from Herreros et al.). We decided to separately present a results from Herreros et al. because of only in the pioneering center outcomes of cell-treatment patients were significantly better than in the rest

centers (the healing rate in 19 centers were 50% (n=124) but in pioneering center reached 70% (n=23)). This fact suggest major influence of experience in cell-based products management on the final results.

In the case of Crohn’s Disease overall healing rate was 66% (n=107) which include Lee et al. [16] where reported healing rates even up to 82%. There are several different strategies reported in the literature (use of auto or allogenic cells, the administration of from 9 to 139 million cells, the administration of cells in fibrin glue, or alone, administration of cells at once or sequentially until healed, and various ways to provide the fistula before administration of the cells). Therefore it is very difficult to compare results between studies. However, achieved results show, that there is a positive effect of the MSC on soft tissue healing.

In the presented studies some of AE (adverse effects) were reported but only in Herreros et al. [15] there were reporting in systematic condition. The most frequent adverse effects [15] were a proctalgia (43%) (that was also reported by our team (data not shown)) and abscess drainage (22.4%). Pain, perianal abscess, pyrexia, swelling and pruritus were also reported in around 10%

or less patients who were included in protocol [15]. Frequency of these effects were no statistically differences within groups which received the cells or fibrin glue. In longer observation the most probability AE, directly associated with cell transplantation - uncontrolled cellular growth were no reported in Herreros et al. and another publication as well.

Lack of differences between groups in occurring of the AE and lack of the uncontrolled cellular growth suggest that reported adverse effects were more related to the act of administration an ascetical products risky perianal location than the products were a mesenchymal stem cells. It must be stressed here that cells cannot be sterilized as typical pharmaceuticals (ex. gamma sterilization) even more a precise bacteriological status of cell-living product is unknown at the time of cells administration (all methods of microbiological assessment have to be incubated at least a couple of hours to observe a pre-result; but an aseptic examination that have to be conduct to precise assess of the final product require a 2 weeks of incubation (accordingly with European Pharmacopeia). From that reasons using of cell-living products demand a high quality of cell production in cell banks, trained medical staff as well as perfect coordination between themselves.

## Conclusion

In clinical trials so far, the healing rates of MSC fistula treatment was about 70%. Pleiotropic effect of action of the stem cells seems to be particularly beneficial in the process of perianal fistula healing, but we still don't know, whether the cells are permanently built into the tissue, and what clinical value have the properties of the cells we described. From that reason there is still grate necessity to explore a mechanism of soft tissue healing driven by autogenic as well as allogenic MSC. Developing of spin-labelling technics and magnetic resonance imagining seems to be useful tools in investigation of "fate of cells" after implantation and can be directly implemented in clinical trials protocols. The explanation of these aspects will help to develop better strategies to introduce the cells and thus contribute to the definition of precise indications for such treatment.

The cell-based therapies that are still in investigational phase require from all scientific and medical society to treat the patients only in restricted conditions of registered clinical trials or compassionate use programs to decrease chance of serious adverse effects. In over 10-year long research on safety and effectiveness of the mesenchymal stem cells in treatment of soft tissues give hope for a new, safer therapeutic option, especially useful in difficult fistula cases but still need more effort to better understand a nature and consequences of cell-based products.

## References

- Rojanasakul A (2009) LIFT procedure: a simplified technique for fistula-in-ano. *Tech Coloproctol* 13(3): 237-240.
- Schwartz DA, Loftus EV Jr, Tremaine WJ, Panaccione R, Harmsen WS, et al. (2002) The natural history of fistulizing Crohn's disease in Olmsted County, Minnesota. *Gastroenterology* 122(4): 875-880.
- García-Aguilar J, Davey CS, Le CT, Lowry AC, Rothenberger DA (2000) Patient satisfaction after surgical treatment for fistula-in-ano. *Dis Colon Rectum* 43(9): 1206-1212.
- Blumetti J, Abcarian A, Quinteros F, Chaudhry V, Prasad L, et al. (2012) Evolution of treatment of fistula in ano. *World J Surg* 36(5): 1162-1167.
- García-Olmo D, García-Arranz M, García LG, Cuellar ES, Blanco IF, et al. (2003) Autologous stem cell transplantation for treatment of rectovaginal fistula in perianal Crohn's disease: a new cell-based therapy. *Int J Colorectal Dis* 18(5): 451-454.
- da Silva Meirelles L, Chagastelles PC, Nardi NB (2006) Mesenchymal stem cells reside in virtually all post-natal organs and tissues. *J Cell Sci* 119(pt 11): 2204-2213.
- Kern S, Eichler H, Stoeve J, Klüter H, Bieback K (2006) Comparative analysis of mesenchymal stem cells from bone marrow, umbilical cord blood, or adipose tissue. *Stem Cells* 24(5): 1294-1301.
- Zuk PA, Zhu M, Mizuno H, Huang J, Futrell JW, et al. (2001) Multilineage cells from human adipose tissue: implications for cell-based therapies. *Tissue Eng* 7(2): 211-228.
- Kim WS, Park BS, Sung JH, Yang JM, Park SB, et al. (2007) Wound healing effect of adipose-derived stem cells: a critical role of secretory factors on human dermal fibroblasts. *J Dermatol Sci* 48(1): 15-24.
- Salgado AJ, Reis RL, Sousa NJ, Gimble JM (2010) Adipose tissue derived stem cells secretome: soluble factors and their roles in regenerative medicine. *Curr Stem Cell Res Ther* 5(2): 103-110.
- Lee EY, Xia Y, Kim WS, Kim MH, Kim TH, et al. (2009) Hypoxia-enhanced wound-healing function of adipose-derived stem cells: Increase in stem cell proliferation and up-regulation of VEGF and bFGF. *Wound Repair Regen* 17(4): 540-547.
- Keyser KA, Beagles KE, Kiem HP (2007) Comparison of mesenchymal stem cells from different tissues to suppress T-cell activation. *Cell Transpl* 16(5): 555-562.
- González MA, Gonzalez-Rey E, Rico L, Büscher D, Delgado M (2009) Adipose-derived mesenchymal stem cells alleviate experimental colitis by inhibiting inflammatory and autoimmune responses. *Gastroenterology* 136(3): 978-989.
- García-Olmo D, García-Arranz M, Herreros D, Pascual I, Peiro C, et al. (2005) A phase I clinical trial of the treatment of Crohn's fistula by adipose mesenchymal stem cell transplantation. *Dis Colon Rectum* 48(7): 1416-1423.
- Herreros MD, Garcia-Arranz M, Guadalajara H, De-La-Quintana P, Garcia-Olmo D, et al. (2012) Autologous expanded adipose-derived stem cells for the treatment of complex cryptoglandular perianal fistulas: a phase III randomized clinical trial (FATT 1: fistula Advanced Therapy Trial 1) and long-term evaluation. *Dis Colon Rectum* 55(7): 762-72.
- Lee WY, Park KJ, Cho YB, Yoon SN, Song KH, et al. (2013) Autologous adipose tissue-derived stem cells treatment demonstrated favorable and sustainable therapeutic effect for Crohn's fistula. *Stem Cells* 31(11): 2575-2581.
- Cho YB, Lee WY, Park KJ, Kim M, Yoo HW, et al. (2013) Autologous adipose tissue-derived stem cells for the treatment of Crohn's fistula: a phase I clinical study. *Cell Transplant* 22(2): 279-285.
- Ciccocioppo R, Bernardo ME, Sgarella A, Maccario R, Avanzini MA, et al. (2011) Autologous bone marrow-derived mesenchymal stromal cells in the treatment of fistulizing Crohn's disease. *Gut* 60(6): 788-798.
- Molendijk I, Bonsing BA, Roelofs H, Peeters KC, Wasser MN, et al. (2015) Allogeneic Bone Marrow-Derived Mesenchymal Stromal Cells Promote Healing of Refractory Perianal Fistulas in Patients With Crohn's Disease. *Gastroenterology* 149(4): 918-927.

20. de la Portilla F, Alba F, García-Olmo D, Herreras JM, González FX, et al. (2013) Expanded allogeneic adipose-derived stem cells (eASCs) for the treatment of complex perianal fistula in Crohn's disease: results from a multicenter phase I/IIa clinical trial. *Int. J Colorectal Dis* 28(3): 313-323.
21. Garcia-Olmo D, Herreros D, Pascual M, Pascual I, De-La-Quintana P, et al. (2009) Treatment of enterocutaneous fistula in Crohn's Disease with adipose-derived stem cells: a comparison of protocols with and without cell expansion. *Int J Colorectal Dis* 24(1): 27-30.
22. Garcia-Olmo D, Guadalajara H, Rubio-Perez I, Herreros MD, de-la-Quintana P, et al. (2015) Recurrent anal fistulae: limited surgery supported by stem cells. *World J Gastroenterol* 21(11): 3330-3336.
23. Garcia-Olmo D, Herreros D, Pascual I, Pascual JA, Del-Valle E, Zorrilla J, et al. (2009) Expanded adipose-derived stem cells for the treatment of complex perianal fistula: A phase ii clinical trial. *Dis Colon Rectum* 52(1): 79-86.
24. García-Arranz M, Dolores Herreros M, González-Gómez C, de la Quintana P, Guadalajara H, et al. (2016) Treatment of Crohn's-Related Rectovaginal Fistula With Allogeneic Expanded-Adipose Derived Stem Cells: A Phase I–IIa Clinical Trial. *Stem Cells Transl Med* 2015-0356.